

The effect of intracameral, per-operative antibiotics on microbial contamination of anterior chamber aspirates during phacoemulsification

BIJAN BEIGI, WILLIAM WESTLAKE,
BERNARD CHANG, CATHERINE MARSH,
JOHN JACOB, TERRY RIORDAN

Abstract

Purpose To assess the effect of per-operative antibiotics on contamination of anterior chamber (AC) aspirates during phacoemulsification.

Methods Two hundred and twenty patients undergoing phacoemulsification of cataract were randomly allocated to receive an irrigation infusion fluid containing either balanced salt solution (BSS) alone or BSS with vancomycin (20 mg/l) and gentamicin (8 mg/l) during surgery. Conjunctival swabs were obtained from all patients immediately before pre-operative preparation. At the end of surgery 20 ml of the AC aspirate was sent for direct and enrichment cultures. Qualitative and quantitative microbiological studies were undertaken. The chi-squared test was used to compare differences between the two groups.

Results There was no significant difference between the positive culture rates of the conjunctival swabs (28 vs 27; $p > 0.8$). In the group that received BSS alone there were 22 (20%) positive AC aspirate cultures, 18 of which were from enrichment cultures. There were 3 positive (2.7%) cultures from the group that received antibiotics added to the BSS ($p < 0.0001$).

Conclusion The addition of gentamicin and vancomycin to the irrigation fluid during phacoemulsification results in a highly significant reduction in the microbial contamination of AC aspirates.

Key words Anterior chamber contamination, Endophthalmitis, Intraocular antibiotics, Phacoemulsification

Endophthalmitis is a small but serious risk of intraocular surgery. The incidence has been found to be lower for extracapsular cataract

extraction (0.1%) than intracapsular extraction (0.7%).¹ Recent studies in developed countries have reported an incidence as high as 0.3–0.7%.^{2,3} The vast number of operations performed worldwide and the serious visual morbidity arising from endophthalmitis demand that every effort be made to minimise the incidence further.

Endophthalmitis usually arises from contamination of the operating field during surgery by commensals of the ocular surface or airborne bacteria.⁴ These contaminate the anterior chamber (AC) either directly or by adhering to the intraocular lens.⁵ Endogenous infections are felt to be uncommon.⁶ Most, if not all, post-operative intraocular infections are caused by organisms introduced at the time of the surgery rather than subsequently acquired.⁷

The commonest Gram-positive causative organisms are coagulase-negative staphylococci (CNS), but a significant number of endophthalmitis cases arise from Gram-negative organisms such as *Proteus* sp.⁸ Efforts have been made to reduce per-operative intraocular microbial contamination by meticulous pre-operative ocular preparation, such as the pre-operative use of povidone-iodine⁹ and the administration of per-operative broad spectrum antibiotics.

We designed a prospective, randomised double-masked study to assess the effect of per-operative broad spectrum antibiotics on contamination of the AC during phacoemulsification.

Materials and methods

The conjunctival flora and microbial contamination of AC aspirates from 220 patients undergoing routine cataract extraction with phacoemulsification were studied. Patients were randomly allocated to receive one of two

B. Beigi
W. Westlake
B. Chang
C. Marsh
J. Jacob
West of England Eye Unit
Royal Devon & Exeter
Hospital
Exeter
Devon EX2 5DW, UK

B. Beigi ✉
Bristol Eye Hospital
Lower Maudlin Street
Bristol BS1 2LX, UK
T. Riordan
Public Health Laboratories
Royal Devon & Exeter
Hospital
Exeter
Devon EX2 5DW, UK

infusion fluid solutions. The theatre staff in charge of the list were responsible for randomisation by drawing colour-coded cards for each case. In group 1 (110 patients) balanced salt solution (BSS) with 1.2 ml/l of adrenaline 1/1000 was used for infusion during surgery. In group 2 (110 patients) 20 mg/l vancomycin and 8 mg/l gentamicin were added to the BSS containing adrenaline.

Patients with ocular or general infectious disease and those who had a history of previous intraocular surgery were excluded from the study. A bacterial filter was not used with the infusion tubing. Both the surgeon and the microbiologist were masked to the allocation of each patient.

Pre-operatively guttae phenylephrine 2.5% and cyclopentolate 1% were instilled in the conjunctival sac every 15 min for 1 h. Eighty per cent of operations were performed with the patient under local anaesthesia, using a peribulbar block containing a mixture of equal proportions of xylocaine 2% and bupivacaine 0.5%.

Conjunctival swabs were taken from each patient prior to skin preparation. The skin was cleaned with povidone-iodine 10% and a 5% solution instilled into the conjunctival sac. An adhesive drape was used to protect the surgical field from the lashes and lid margins.

Cataract surgery was performed through a scleral tunnel using a frown incision. A 1.5 mm paracentesis was formed for insertion of the second instrument. After continuous circular capsulorhexis and hydrodissection, the crystalline lens was removed using the divide-and-conquer phacoemulsification technique. The scleral incision was extended to 5.5 mm and a 5 × 6 mm one-piece PMMA phaco intraocular lens (IOLAB) placed in the capsular bag. The incision site was covered by cauterising the conjunctiva at the limbus; no suture was used. Directly after each procedure 20 ml of fluid was poured from the phaco aspiration collection cassette into a sterile specimen bottle. At the end of each operating list the AC aspirates and swabs that had been collected were sent for microbiological studies. Fifty control swabs (sterile) and 50 specimen bottles containing sterile sodium chloride were sent among the study specimens.

Microbiological studies

Conjunctival cultures

Conjunctival swabs were spread onto a blood agar plate (5% Horseblood Columbia Agar base, Unipath), chocolate agar plate and a Sabouraud agar plate with gentamicin. The chocolate agar plate was incubated at 37 °C in 5% CO₂ for 48 h. The blood agar plate was incubated in an anaerobic cabinet for 7 days and the Sabouraud plate at 37 °C for 7 days. An incubation temperature of 37 °C was felt to be more suitable for detection of intraocular contaminants than the 32 °C commonly used for the study of corneal infections.

AC aspirate

AC aspirate (0.1 ml) was inoculated by pipette onto the same types of plates as were used for the swabs, and spread with a spreader. The plates were then cultured as for conjunctival swabs (direct culture). In addition, the fluid was centrifuged and the deposit resuspended in 2 ml of fluid (enrichment culture). This was then inoculated into a Pedi-bact blood culture broth (Organon-Technika), supplemented with 1 ml of Fildes extract. The bottles were incubated in a BacTAlert machine for 7 days. This is as effective as thioglycollate broth for the detection of anaerobes. Any bottles flagging positive were unloaded and subcultured both aerobically and anaerobically.

Identification of isolates

Any growth, even if scanty, was recorded and isolates were identified to genus/species level using standard methods (API kit). CNS were not identified to the species level.

Statistical method

After a pilot study, it was calculated that our sample size would have an 80% chance of producing a significant result at $p < 0.05$ (EPI INFO Stat Calc). The chi-squared test was used to test for statistical significance between the two groups.

Results

Conjunctival cultures

Group 1 (BSS alone) had 28 (25.4%) positive conjunctival swab cultures. Group 2 (BSS with antibiotics) had 27 (24.3%) positive cultures. The two groups had a similar microbial flora (Tables 1, 2). There was no significant difference between the positive culture rates of the conjunctival swabs in the two groups ($\chi^2 = 0.02$; $p > 0.8$). None of the 50 control swabs was positive.

AC aspirate cultures

There were 22 (20%) positive AC aspirate cultures in group 1. Six of the direct and 21 of the enriched AC aspirate cultures grew a microorganism. In group 2 there were only 3 (2.7%) positive cultures, 2 of which were identified by enriched cultures. Intraocular bacterial contamination during surgery was significantly lower in this group ($\chi^2 = 16.29$; $p < 0.0001$). The types of microorganism grown in the two groups are shown in Tables 1 and 2. None of the 50 control bottles cultured positive.

Discussion

The risk of endophthalmitis is best minimised by eliminating the commensals of the ocular surface and preventing the entry of microbes into the AC during

Table 1. Bacterial growth in group 1 (BSS alone)

| Patient no. | Conjunctival swabs | AC culture | |
|-------------|----------------------------------|--------------------------------|--------------------------------|
| | | Direct | Enrichment |
| 112 | | | Group B streptococci |
| 113 | CNS | | CNS |
| 114 | <i>Acinetobacter lwoffii</i> | | |
| 115 | | CNS | |
| 116 | | | <i>Acinetobacter</i> sp. |
| 118 | CNS | | |
| 123 | | | α-haemolytic streptococci |
| 128 | CNS | CNS | CNS |
| 129 | | | CNS |
| 130 | <i>Morganella morganii</i> , CNS | CNS | CNS |
| 136 | CNS | α-haemolytic streptococci, CNS | α-haemolytic streptococci, CNS |
| 138 | CNS | | |
| 143 | CNS | | |
| 148 | | | CNS |
| 152 | | | CNS |
| 156 | Oxidase +ve Gram -ve rod | | |
| 157 | CNS | | |
| 158 | | | CNS |
| 166 | | | CNS |
| 168 | CNS, oxidase +ve Gram -ve rod | | |
| 169 | CNS | | |
| 170 | CNS | | |
| 174 | CNS | | |
| 176 | <i>Candida</i> sp. | | CNS |
| 181 | CNS | | <i>Acinetobacter</i> sp. |
| 184 | <i>Staphylococcus aureus</i> | | |
| 185 | CNS | | |
| 187 | CNS | | |
| 189 | <i>Haemophilus influenzae</i> | | |
| 193 | α-haemolytic streptococci | | Oxidase -ve Gram +ve rod |
| 197 | <i>Staphylococcus aureus</i> | | |
| 198 | CNS | | CNS |
| 201 | | <i>Candida</i> sp., CNS | <i>Candida</i> sp., CNS |
| 202 | | | CNS |
| 207 | | | CNS |
| 208 | <i>Staphylococcus aureus</i> | | |
| 209 | Coliforms | | |
| 210 | | | <i>Moraxella</i> sp. |
| 212 | CNS | | |
| 216 | | | CNS |
| 218 | CNS | | |
| 219 | α-haemolytic streptococci | | |

CNS, coagulase-negative staphylococci.

surgery. Per-operative intracameral antibiotics provide a second line of defence and aid the elimination of microbes that have breached other preventive measures – for example those in the 2–3% of irrigation solutions that have previously been found to be contaminated.⁷

Although we have noticed a poor identification rate of anaerobes, conjunctival culture of the two groups in this study showed a comparable bacterial flora pre-operatively. Probably more anaerobes would have been isolated if plates had been incubated for 14 instead of 7 days. Pre-operative skin preparation with 10% povidone-iodine and instillation of a 5% solution in the conjunctival sac is effective at reducing the number of commensal organisms within the operating field^{9,10} and was used in both groups. Pre-operative topical antibiotics, such as norfloxacin, gentamicin and chloramphenicol, are not effective,^{11,12} and were not used

in this study. They can produce corneal toxicity and inhibit epithelial healing.¹³ Sub-conjunctival antibiotics were not used; these have been shown to have no impact on post-operative inflammation and infection.¹⁴

Gentamicin and vancomycin were added to the BSS in group 2 of this study. These are broad spectrum antibiotics and provide good coverage against Gram-negative and Gram-positive organisms respectively. The estimated concentration in the AC was 8 µg/ml for gentamicin and 20 µg/ml for vancomycin. These concentrations are within the maximum inhibitory concentration (MIC) of most Gram-negative and Gram-positive organisms likely to contaminate the operative field.¹⁵ Antibiotics at these concentrations are not toxic to the retina and corneal endothelial cells.

Gills *et al.*¹⁶ used antibiotics in the irrigation solution during cataract surgery in a large but poorly controlled

Table 2. Bacterial growth in group 2 (BSS with antibiotics)

| Patient no. | Conjunctival swabs | AC culture | |
|-------------|------------------------------------|-----------------------------|------------|
| | | Direct | Enrichment |
| 3 | CNS | | |
| 10 | CNS | | CNS |
| 11 | CNS | | |
| 12 | <i>Proteus mirabilis</i> | | |
| 14 | CNS | | |
| 17 | CNS | | |
| 24 | CNS | | |
| 27 | <i>Moraxella</i> sp. | | |
| 29 | | | CNS |
| 31 | CNS | | |
| 37 | <i>Acinetobacter lwoffii</i> , CNS | | |
| 41 | CNS | | |
| 43 | CNS | | |
| 46 | CNS | | |
| 50 | CNS | | |
| 52 | CNS | | |
| 54 | CNS | | |
| 59 | CNS | | |
| 66 | CNS | | |
| 67 | CNS | | |
| 77 | <i>Proteus</i> sp. | | |
| 79 | α -haemolytic streptococci | | |
| 86 | CNS | <i>Pseudomonas stutzeri</i> | |
| 95 | CNS coryneforms | | |
| 102 | CNS | | |
| 103 | CNS | | |
| 106 | CNS | | |
| 109 | CNS | | |

CNS, coagulase-negative staphylococci.

series. They reported a 1 in 20 000 post-operative intraocular infection rate using gentamicin in the irrigating fluid, 1 in 9928 using vancomycin and none in 25 000 when both vancomycin and gentamicin were used. Gimbel *et al.*¹⁷ reported no cases of endophthalmitis in a group of 4684 cataract extractions in which both vancomycin was injected into the capsular bag at the end of the procedure and gentamicin added to the irrigation solution. They showed no endothelial cell loss due to the use of antibiotics. There were no other reported complications such as retinal toxicity. These studies inspired us to select gentamicin and vancomycin for per-operative use.

Whilst these reports have encouraged the per-operative use of antibiotics during intraocular surgery, they have not addressed the potential complications adequately. Mistakes can be made during the mixing of the antibiotic solutions. It has been suggested that pharmacists are less liable to such errors than surgeons and should prepare the infusion solutions.¹⁸ A single-dose commercial preparation of the antibiotics for this purpose would also be useful. Induced resistance of microorganisms because of the widespread use of antibiotics in a large population is a further concern. To quantify the benefits of per-operative antibiotic use against its potential complications a large multi-centre study is required.

Our study was not designed to assess the effect of using per-operative antibiotics on the incidence of post-operative endophthalmitis. The aim was to identify,

quantify and compare the microorganisms left *in situ* at the end of phacoemulsification for each group.

Nevertheless, the majority of cases of panophthalmitis are caused by the growth of bacterial colonies introduced into the AC or capsular bag at surgery,⁷ so it would be logical to expect a lower infection rate following a reduction of AC contamination. None of our patients had post-operative endophthalmitis. Case 86 in group 2 developed post-operative membranous uveitis with posterior synechiae. The per-operative AC aspirate had grown *Pseudomonas stutzeri* on the enrichment plate that was resistant to gentamicin. This organism is not usually a pathogen in humans.

We have demonstrated nearly a 7-fold reduction (95% CI: 4.47–9.92-fold reduction) in bacterial contamination of the AC as a result of the per-operative use of two antibiotics (Fig. 1). This was a highly significant reduction ($p < 0.0001$). In contrast Henry *et al.*,¹⁹ in a much smaller study using gentamicin in the infusion fluid, failed to show a significant difference.

Endophthalmitis has been reported despite the use of antibiotics in the irrigation field.²⁰ Indeed before the onset of this study we experienced a case of post-operative panophthalmitis from which no microorganism was isolated following the use of per-operative antibiotics. This patient was treated with broad spectrum antibiotics and subsequently achieved a visual acuity of 6/9.




Fig. 1. Comparison of sample size, conjunctival culture and anterior chamber culture results of group 1 (no antibiotic; dark columns) and group 2 (with antibiotic; light columns).

Conclusion

The majority of cases of post-operative panophthalmitis result from the introduction of microorganisms during intraocular surgery.⁷ Meticulous pre-operative preparation of the ocular surface and conjunctiva by povidone-iodine, the use of adhesive sterile drapes to keep the eyelashes away from the operating field and standard protocols regarding instrument sterilisation and BSS preparation will reduce the introduction of microbes into the AC. In this study, addition of vancomycin and gentamicin to irrigation fluids reduced the AC contamination significantly (Fig. 1). The peri-operative use of intracameral antibiotics is gaining popularity, although there are potential problems associated with their use. Large multi-centre studies are needed to quantify the benefits against these possible risks. On balance, the decision regarding the peri-operative use of antibiotics will depend on the answer to the question: 'What price are we prepared to pay to reduce the risk of endophthalmitis to the absolute minimum?'²¹

The authors would like to thank all the theatre staff at the West of England Eye Unit and Mr L. Joslin in the Microbiology Department for their help during the study. We would also like to thank Mr A. Alaghebandian for his help and support.

References

- Javitt JC, Vitale S, Canner JK, Street DA, *et al.* National outcomes of cataract extraction. *Arch Ophthalmol* 1991;109:1085-9.
- Kattan HM, Flynn HW Jr, Pflugfelder SC, *et al.* Nosocomial endophthalmitis survey: current incidence of infection after intraocular surgery. *Ophthalmology* 1991;98:227-38.
- Verbracken H. Treatment of postoperative endophthalmitis. *Ophthalmologica* 1995;209:165-71.
- Sherwood DR, Rich WJ, Jacob JC, Hart RJ, Fairchild YL. Bacterial contamination of intraocular and extraocular fluids during extracapsular cataract extraction. *Eye* 1989;3:308-12.
- Doyle A, Beigi B, Early A, Blake A, Eustace P, Hone R. Adherence of bacteria to intraocular lenses, a prospective study. *Br J Ophthalmol* 1995;79:347-9.
- Spencer SR, Dealler SP, Hassett PDA, Todd NJ, Hawkey PM, Noble BA. Bacterial contamination of intraocular lenses: the source of the bacteria. *Eye* 1989;3:685-9.
- Assalin A, Thompson P, St Antonie P, Lemire J, *et al.* Anterior chamber fluid contamination after uncomplicated phacoemulsification. *J Cataract Refract Surg* 1995;21:539-42.
- Heaven CJ, Mann PJ, Boase DL. Endophthalmitis following extracapsular cataract surgery: a review of 32 cases. *Br J Ophthalmol* 1992;76:419-23.
- Apt L, Isenberg SJ, Yoshimor R, Spierer A. Outpatient topical use of povidone-iodine in preparing the eye for surgery. *Ophthalmology* 89;96:289-92.
- Klie F, Boge I, Jensen OL. The effect of polyvinylpyrrolidone iodine as a disinfectant in eye surgery. *Acta Ophthalmol (Copenh)* 1986;64:67-71.
- Maeck CR, Eckardt C, Holler C. Bacteriocidal effect of preoperative use of gentamicin in comparison with PVP-iodine solution. *Fortschr Ophthalmol* 1991;88:848-51.
- Chitkara DK, Manners T, Chapman F, Stoddart MG, Hill D, Jenkins D. Lack of effect of the pre-operative norfloxacin on bacterial contamination of anterior aspirates after cataract surgery. *Br J Ophthalmol* 1994;78:772-4.
- Caronia RM, Obstbaum SA. Drug therapy before, during and after cataract surgery. *Curr Opin Ophthalmol* 1993;4:61-74.
- Sanders R, MacEwen CJ, Haining WW. A comparison of prophylactic, topical and subconjunctival treatment in cataract surgery. *Eye* 1992;6:105-10.
- Kucers A, Bennett NM. Genatmicin. In: Kurcers A, Bennett NM, editors. *The use of antibiotics: a comprehensive review with clinical emphasis*. 4th ed. Philadelphia: JB Lippincott, 1987:629.
- Gills JP, *et al.* Filters and antibiotics in irrigating solution for cataract surgery. *J Cataract Refract Surg* 1991;17:385.
- Gimbel HV, De Brof RS, De Brof BM. Prophylactic intracameral antibiotics during cataract surgery: the incidence of endophthalmitis and corneal endothelial cell loss. *Eur J Implant Ref Surg* 1994;6:280-5.
- Jeglun EL, Rosenberg SB, Benson WE. Preparation of intra-vitreous drug doses. *Ophthalmic Surg* 1981;12:355-9.
- Henry JC, Rozas D, Bryant E, Memphis W. Bacterial growth from anterior chamber fluid aspirates using different irrigating solution in phacoemulsification. *Invest Ophthalmol Vis Sci* 1994;34:884.
- Alfonso EC, Flynn HW Jr. Controversies in endophthalmitis prevention: the risk of emerging resistance to vancomycin. *Arch Ophthalmol* 1995;113:1369-70.
- Towler H. Aqueous contamination during small incision cataract surgery: a lesson in study design. *Br J Ophthalmol* 1995;79:873.