

ultraviolet light.<sup>2</sup> It is certainly worthwhile to identify high-risk groups and offer screening, but it would also be helpful to identify the relevant environmental factor(s) as well.

#### References

1. Houlston R, Damato B. Genetic predisposition to ocular melanoma. *Eye* 1999;13:43–6.
2. Dolin PJ, Foss AJE, Hungerford JL. Uveal melanoma: is solar ultraviolet radiation a risk factor? *Ophthalmic Epidemiol* 1994;1:27–30.
3. Foss AJE, Dolin PJ. Trends in eye cancer mortality among adults in the USA and England and Wales. *Br J Cancer* 1996;74:1687–9.
4. Foss AJE, Cree IA, Dolin PJ, Hungerford JL. Modelling uveal melanoma. *Br J Ophthalmol* 1999;83:760–70.
5. Bercher L, Munier F, Zografos L, Schorderet D, Egger E, Chamot L. Familial uveal melanoma. *Klin Monatsbl Augenheilkd* 1995;206:384–7.
6. Singh AD, Shields CL, Shields JA, De Potter P. Bilateral primary uveal melanoma. Bad luck or bad genes? *Ophthalmology* 1996;103:256–62.
7. Young LH, Egan KM, Walsh SM, Gragoudas ES. Familial uveal melanoma. *Am J Ophthalmol* 1994;117:516–20.

Alexander Foss ✉  
Department of Ophthalmology  
Queen's Medical Centre  
Nottingham NG7 2UH, UK

Sir,

The purpose of our article<sup>1</sup> was to highlight the increasing evidence that some cases of uveal melanoma arise as a result of a genetic predisposition. We do not think that we overstated the role of constitutional gene mutations in the development of uveal melanoma. Although as Foss points out only 2% of cases are familial as defined by having a relative affected with uveal melanoma, indirect evidence suggests that a greater number, around 7% of cases, are likely to be caused by mutations in genes with pleiotropic effects such as *BRCA2* and those causing the atypical naevus syndrome.

To assess the contribution of germline mutations to the development of uveal melanoma we have made a systematic collection of family histories, blood samples and tumour material from over 400 patients attending the Ocular Oncology Service in Liverpool. Using this resource we are currently investigating the contribution of mutations in *BRCA2* and *CDKN2A* to

**Table 1.** Post-operative refraction and best corrected acuity

Post-operative time (weeks)	Refraction	Best corrected acuity
4	+2.00/—	6/18
8	0.00/–1.50 × 90°	6/9
20	+7.00/–2.00 × 150°	6/18

uveal melanoma by screening both these genes in this series of blood samples from these patients.

Clearly the identification of genetic factors does not detract from the potential role of specific environmental factors in the aetiology of uveal melanoma, and in this respect we concur with the view of Foss.

#### Reference

1. Houlston R, Damato B. Genetic predisposition to ocular melanoma. *Eye* 1999;13:43–6.

Richard Houlston ✉  
Section of Cancer Genetics  
Institute of Cancer Research  
Sutton  
Surrey SM2 5NG, UK

Bertil Damato  
Ocular Oncology Service  
Royal Liverpool University Hospital  
Liverpool L7 8XP, UK

Sir,

I have read with interest the paper by Zambarakji *et al.*,<sup>1</sup> and subsequent correspondence,<sup>2</sup> concerning capsulorhexis phymosis following phacoemulsification. My own experience illustrates another possible parameter in the estimation of capsulorhexis phymosis.

A healthy 70-year-old man underwent phacoemulsification via a 5.0 mm capsulorhexis. The procedure was uncomplicated apart from a small (2 clock-hours) zonular dehiscence. A 23.0 D acrylic intraocular lens (Acrygel, Corneal Laboratoire) was implanted into the capsular bag.

Post-operative acuity was 6/9, and further changes were as shown in Table 1 (periods of subnormal best corrected visual acuity were assumed to be due to cystoid macular oedema, not proven by fluorescein angiography, but which improved with the use of oral acetazolamide and topical

betamethasone). At this point capsulorhexis phymosis was noted, and nine radial relaxing incisions were made in the anterior capsular ring with a YAG laser. The posterior capsule was left intact. The refraction and best corrected acuities following YAG laser are shown in Table 2.

The only logical explanation for this remarkable variation in post-operative refraction is posterior displacement, or posterior bowing, of the flexible intraocular lens caused by capsulorhexis phymosis and relieved by YAG laser relaxing incisions. Previous literature has alluded to the fact that capsulorhexis phymosis can alter refraction,<sup>3</sup> and Shamma<sup>4</sup> has measured such changes in refraction. In one case, he reports +1.25 D, and in another, +0.75 D of induced hyperopia.

My experience indicates that accurate refraction may help in monitoring some cases of capsulorhexis phymosis, especially if a foldable intraocular lens is used.

#### References

1. Zambarakji HJ, Rauz S, Reynolds A, Joshi N, Simcock PR, Kinnear PE. Capsulorhexis phymosis following uncomplicated phacoemulsification surgery. *Eye* 1997;11:635–8.
2. Walsh LM, Pande M. [Correspondence]. *Eye* 1998;12:1036–7.
3. Hansen SO, Crandall AS, Olson RJ. Progressive constriction of the anterior capsular opening following intact capsulorhexis. *J Cataract Refract Surg* 1993;19:77–82.
4. Shamma HJ. Relaxing the fibrosed capsulorhexis rim to correct induced hyperopia after phacoemulsification. *J Cataract Refract Surg* 1995;21:228–9.

Robert Murray, FRCS, FRCOphth ✉  
Borders General Hospital  
Melrose  
Roxburghshire TD6 9BS, UK  
Tel: +44 (0)1896 754333, ext 1033  
Fax: +44 (0)1896 823476

**Table 2.** Post-YAG laser refraction and best corrected acuity

Post-YAG laser time (weeks)	Refraction	Best corrected acuity
2	+4.00/–1.50 × 180°	6/18
7	+1.75/–0.25 × 180°	6/6
26	+1.00/—	6/6