

Reference

1. Choong YF, Irfan S, Menage MJ. Acute angle closure glaucoma: an evaluation of a protocol for acute treatment. *Eye* 1999;13:613–6.

Elizabeth Patsoura ✉  
J.M. Butcher  
Ophthalmology Department  
Countess of Chester Hospital  
Liverpool Road  
Chester CH2 1UL, UK

Sir,

The maximum effect of oral Diamox occurs at 2 h but the intravenous Diamox is much faster. If no reduction in intraocular pressure has occurred an hour and a half after the intravenous Diamox then it is not unreasonable to consider further treatment at that stage. After all this condition is extremely painful with potential to damage the optic nerve, and a rapid reduction in intraocular pressure from the very high levels exhibited by some patients is desirable.

Mr M.J. Ménage, MRCP, FRCS, FRCOphth ✉  
Department of Ophthalmology  
The General Infirmary at Leeds  
Great George Street  
Leeds  
West Yorkshire LS1 3EX, UK  
Tel: +44 (0)113 243 2799  
Fax: +44 (0)113 392 6336

Sir,

We read with interest the editorial<sup>1</sup> and article<sup>2</sup> on the immediate management of acute angle-closure glaucoma (AACG). We would, however, like to propose an alternative: argon laser peripheral iridoplasty (ALPI).<sup>3,4</sup> ALPI was traditionally employed only when medications failed to control intraocular pressure (IOP),<sup>5</sup> but we found it effective and safe as an initial treatment.

Immediate ALPI may be more effective than conventional medications in lowering IOP. Employing the same criteria of 'satisfactory IOP control'

proposed by Choong *et al.*<sup>2</sup> we achieved satisfactory IOP control in 83.3% of AACG patients, at just 15 min after ALPI.<sup>3,4</sup> At 120 min after ALPI we achieved satisfactory IOP control in 100% of patients. We have, however, excluded patients presenting more than 48 h from the onset of symptoms.

We have also documented, using ultrasound biomicroscopy, re-opening of the closed drainage angle immediately following ALPI.<sup>3</sup> By opening up the angle promptly, we would expect a lower chance of a patient developing peripheral anterior synechiae, and subsequently chronic angle-closure glaucoma.

Immediate ALPI in the management of AACG appears to be a safe procedure. It will certainly spare patients the systemic side effects of acetazolamide, glycerol or mannitol. We have followed up the 18 reported cases for 2 years by now, and so far we have not come across any patient with corneal decompensation or scarring, iris atrophy or necrosis, or other complications that could have arisen from the ALPI. We have also shown, in another study, that halving the quantity of ALPI still yields IOP-lowering results comparable to the conventional 360° treatment.<sup>6</sup> By applying ALPI to only 180° of the peripheral iris, the risk of complication from ALPI is further lowered.

After ALPI, all patients would still require a peripheral iridotomy to break the pupillary block, and to avoid recurrent attack of acute glaucoma. Most commonly, the peripheral iridotomy has to be delayed until the cornea recovers from the acute attack. With the use of immediate ALPI, all corneas clear within 2 h,<sup>3,4</sup> allowing a very early peripheral iridotomy to be performed.

Supported in part by the Dr Yu-Tung Cheng Eye Foundation.

References

1. Salmon JF. The management of acute angle-closure glaucoma. *Eye* 1999;13:609–10.

2. Choong YF, Irfan S, Menage MJ. Acute angle-closure glaucoma: an evaluation of a protocol for acute treatment. *Eye* 1999;13:613–6.
3. Lam DSC, Lai JSM, Tham CCY. Immediate argon laser peripheral iridoplasty (ALPI) as treatment for acute attack of primary angle closure glaucoma (PACG): a preliminary study. *Ophthalmology* 1998;105:2231–6.
4. Tham CCY, Lai JSM, Lam DSC. Immediate argon laser peripheral iridoplasty for acute attack of PACG [letter]. *Ophthalmology* 1999;106:1042–3.
5. Ritch R. Argon laser treatment for medically unresponsive attacks of angle-closure glaucoma. *Am J Ophthalmol* 1982;94:197–204.
6. Lai JSM, Tham CCY, Lam DSC. Limited argon laser peripheral iridoplasty as immediate treatment for an acute attack of primary angle closure glaucoma: a preliminary study. *Eye* 1999;13:26–30.

Dennis S.C. Lam  
Department of Ophthalmology & Visual Sciences  
The Chinese University of Hong Kong  
Hong Kong Eye Hospital  
Kowloon, Hong Kong

Jimmy S.M. Lai  
Clement C.Y. Tham  
Department of Ophthalmology & Visual Sciences  
The Chinese University of Hong Kong  
Prince of Wales Hospital  
Shatin, Hong Kong

Professor Dennis S.C. Lam ✉  
Department of Ophthalmology & Visual Sciences  
The Chinese University of Hong Kong  
Hong Kong Eye Hospital  
147K Argyle Street, Kowloon  
Hong Kong  
Tel: +852 2762 3157  
Fax: +852 2711 0464  
e-mail: dennislam@cuhk.edu.hk