5. Mapstone R. Precipitation of angle closure. *Br J Ophthalmol* 1974;58:46–54.

William Westlake ✉
McCusker Glaucoma Unit
The Lions Eye Institute
Nedlands
6019 Western Australia
e-mail: westlake@networx.net.au

Sir,

We read with great interest the paper by Sadiq and Downes¹ and the letter by Ziakes *et al.*² on dysversion of lateral eyelashes in children. They described a few cases where this diagnosis was made for recurrent sore eyes in children. Two of our cases support this diagnosis.

Case 1

A 5-year-old white girl was referred to the eye clinic by her general practitioner for bilateral recurrent red and sore eyes. On the day of examination her eyes were white. The lateral eyelashes of both upper lids seemed to be quite long and were intumed and tucked under the

lower lid. The roots of the lashes were normally situated. The lashes were manually repositioned using a cotton bud. Her mother was shown the procedure and advised to do it two times a day. The patient was reviewed in the clinic 3 months later and there had been no further episode of soreness and conjunctivitis. The lateral eyelashes were nicely directed down and outwards. On discharge, her mother was told to watch the lateral eyelashes and repeat the procedure when needed. It is 6 months since we last saw them.

Case 2

An 8-year-old boy was referred to the eye clinic by his general practitioner for bilateral recurrent conjunctivitis. On examination his lateral eyelashes of both upper lids were intumed and tucked under the lower lid. The rest of the anterior segments were unremarkable. His grandmother was shown and advised to do the manual repositioning of the lashes using a cotton bud twice a day. Three months later all symptoms had settled and he had no further episodes of conjunctivitis.

Comment

Our experience with the two cases supports the diagnosis of dysversion of lateral eyelashes in children. This condition can indeed cause a great deal of discomfort in patients and can be overlooked by the general practitioner. Ziakes *et al.*² treated their patients with electrolysis. We support the treatment by Sadiq and Downes,¹ of manual repositioning of the lateral eyelashes. This is very simple and avoids general anaesthetic and surgery.

References

1. Sadiq SA, Downes RA. Dysversion of lateral eyelashes in children: a new diagnosis. *Eye* 1996;10:473–5.
2. Ziakes NG, Clarke MP, Strong NP. Dysversion of lateral eyelashes in children: a new diagnosis [letter]. *Eye* 1997;11:572–3.

B.B. Patil ✉
J.R. Clarke
North Riding Infirmary
Newport Road
Middlesbrough TS1 5JE, UK