blade (Swann Morton, Sheffield, UK) to make the initial incision in the nasal mucosa and lacrimal sac. The stab incision is then enlarged with the Werb's scissors. The no. 12 blade is not ideally employed for this purpose due to its large size and long cutting edge. In an effort to improve technique other instruments have been used to incise the mucosa, for example the no. 11 Bard-Parker blade, crescent knife,¹ or a Graefe's knife. We have found the phacoemulsification keratome ideally designed for the purpose of incising the lacrimal sac and nasal mucosa. To our knowledge there are no previous reports of this technique in the literature.

Comment

We have employed the use of the disposable stainless steel Spectrum keratome (72-2561) to make the initial incision into the lacrimal sac and the extension of the incision. The keratome is ideally designed for creating the mucosal flaps in dacryocystorhinostomy. The 2.5 mm keratome is held so that the cutting edge is perpendicular to the surface of the lacrimal sac. A Bowman's 00 probe is inserted into the sac and the initial stab incision is made through the tented portion of the sac wall. Gentle rotation of the instrument allows extension of the initial incision in a controlled manner (Fig. 1). The 60° bevel-up angulation facilitates this manoeuvre. In the relatively confined space available the features of the keratome

Fig. 1. Phacoemulsification keratome used to make an incision in the lacrimal sac wall.

allow the creation of the mucosal flaps to be performed with ease. The keratome is then used to make the incision of the nasal mucosa in the same manner.

To date, we have employed the use of the keratome in 40 dacryocystorhinostomies and have achieved a success rate of 95%. We would therefore advocate the use of the keratome above the no. 12 blade when making the mucosal flaps in dacryocystorhinostomy.

Reference

1. Fong KS, Koh AH, Choo CT. Using the phacoemulsification crescent knife in dacryocystorhinostomy. *Ophthalmic Surg Lasers* 1998;29:343-4.

S. Brannan
T.Q. Reuser
Birmingham and Midland Eye Centre
Birmingham UK

Ms Suzanne O. Brannan ✉
43 Vicarage Road
Yardley
Birmingham B33 8PH, UK

Tel: +44 (0)121 783 4982
Tel: +44 (0) 831 327 384 mobile
e-mail: suzanne@brannan43.fsnet.co.uk

Sir,

An unusual case of cavernous sinus syndrome

Cranial nerve palsies are a documented complication of penetrating intracranial and orbital trauma¹⁻⁴ and Horner's syndrome may result from trauma anywhere along the efferent sympathetic pathway.⁵ Patients who are fortunate enough to survive such severe trauma commonly also have globe or extraocular muscle injury and vascular complications such as carotidocavernous fistula (CCF). We describe a patient with multiple cranial neuropathies, an intact globe and without a CCF or cavernous sinus thrombosis, following an orbital/intracranial gunshot injury.

Case report

A 25-year-old man was shot in the face from a range of 0.6 m with a handgun of uncertain calibre. The bullet passed posterolaterally from an entry wound below the right medial canthus lodging by the apex of the petrous bone. His Glasgow Coma Scale was normal and vital signs stable, so he was managed conservatively by the neurosurgical team for a basal skull fracture with cerebrospinal fluid leak. Two days after admission he reported diplopia and was seen by an ophthalmologist.

The visual acuity was 6/6 each eye. Both globes were intact and the fundi normal. The 3 × 4 mm entry wound was inferomedial to the inferior rim of the right orbit, and involved the lacrimal sac exposing the upper end of the nasolacrimal duct. There was a right inferior canaliculus laceration but the lids were otherwise intact. He had right Horner's syndrome, with 1 mm of ptosis

Fig. 1. Photograph of the patient showing the entry wound.

and anisocoria (the right pupil being relatively miosed) but no anhidrosis. There was no evidence of a CCF or cavernous sinus thrombosis (Fig. 1).

Ocular movements showed deficient abduction (−4) of the right eye with a normal forced duction test excluding restrictive strabismus. Third and fourth cranial nerves were intact. There was normal touch sensation in the mandibular trigeminal dermatome and no trigeminal motor deficit, but the maxillary division of trigeminal had altered light touch sensation, and there was loss of corneal sensation. There were no left-sided cranial nerve abnormalities and no other neurological deficits.

Skull radiographs showed a bullet lodged in the right petrous temporal bone, with the lateral radiograph (Fig. 2) showing a shrapnel trail from point of entry. A CT scan confirmed the position of the bullet, but showed no damage to the extraocular muscles or orbital contents. The metal of the bullet caused signal artefact. Angiography showed thrombosis of the cervical portion of the right internal carotid artery, secondary to proximal dissection from the petrous portion.

The sixth nerve palsy was managed initially by botulinum toxin injection to the right medial rectus, and

Fig. 2. Lateral skull radiograph showing the shrapnel trail from the point of entry of the bullet.

dacryocystorhinostomy with canalicular intubation is planned. Three months after the injury no CCF had developed.

Comment

Pathology in the region of the cavernous sinus and petrous bone may result in a range of cranial nerve palsies. Both cavernous sinus thrombosis and CCF can cause ophthalmoplegia either through the mechanical effects of orbital vascular engorgement or via cranial nerve palsies, but in each case the signs of vascular obstruction are prominent: proptosis, lid swelling, conjunctival and retinal vessel engorgement. Trauma to the internal carotid artery in the cavernous sinus may result in delayed CCF,^{6,7} but our patient had no evidence of increased cavernous sinus venous pressure either at initial presentation or 3 months later. It is possible that the internal carotid artery thrombosis, demonstrated on the angiogram, protected the cavernous sinus from transmission of increased pressure. He fortunately has an intact circle of Willis enabling blood flow to the right ophthalmic, anterior and middle cerebral arteries. The sixth cranial nerve is joined in the anterior cavernous sinus by sympathetic branches; these pass to the iris dilator muscle via the ophthalmic division of the fifth nerve. This anatomy may in part account for the clinical signs present in this patient.

This patient shows an unusual clinical presentation after surviving a gunshot to his head in that the sixth cranial nerve, the maxillary and ophthalmic division of the trigeminal nerve, the sympathetic root and the carotid artery are all affected, but the mandibular division of trigeminal and the third and fourth cranial nerves were spared, the cavernous sinus was not thrombosed, nor did a carotico-cavernous fistula develop.

References

1. Lagreze W. Neuro-ophthalmology of trauma. *Curr Opin Ophthalmol* 1998;9:33–9.
2. Lepore FE. Disorders of ocular motility following head trauma. *Arch Neurol* 1995;52:924–6.
3. Keane JR. Third nerve palsy due to penetrating trauma. *Neurology* 1993;43:1523–7.
4. Sarvesvaran E. Fatal penetrating orbital injuries. *Med Sci Law* 1991;31:261–3.
5. Abad J, Alvarez F, Blazquez MG. An unrecognised neurological syndrome: sixth-nerve palsy and Horner's syndrome due to traumatic intracavernous carotid aneurysm. *Surg Neurol* 1981;16:140–4.
6. Bickerstaff ER. Mechanism of presentation of carotico-cavernous fistulae. *Br J Ophthalmol* 1970;54:186–90.
7. Keane JR. Cavernous sinus syndrome: analysis of 151 cases. *Arch Neurol* 1996;53:967–71.

D.C. Harrison
Groote Schuur Hospital
Cape Town, South Africa

C.J. Roberts ✉
Moorfields Hospital
City Road
London EC1V 2PD, UK