Sir,
The minim technique for diagnostic anterior chamber paracentesis

We describe a technique of aqueous collection by using a minim of saline (Chauvin Pharmaceuticals, Essex, UK) attached to a 25-gauge needle. Benoxinate 0.4% and betadine 5% antiseptic solutions are instilled into the conjunctival sac. The patient is positioned on the slit lamp and a lid speculum is inserted. Sterile gloves are worn and a 25-gauge needle is attached firmly to the end of a saline minim (Figure 1). A 'click' is heard when the needle is inserted correctly. The minim is held between the thumb and forefinger and the saline is passed out through the 25-gauge needle. The minim remains depressed from this point. The eye is held steady either with a forceps or a cotton tip applicator on the nasal conjunctiva. The needle is inserted bevel up through the paralimbal peripheral clear cornea, anterior to the iris plane. Once the eye is entered, compression of the minim is slowly relaxed. The aqueous now enters the minim. Once a sufficient quantity of aqueous has been drained, without anterior chamber (AC) collapse, the minim is withdrawn from the eye. To avoid excessive vacuum, the minim may be depressed only half way before entering the eye. The needle is discarded and the cap is reattached to the minim. The needle-free sample is sent safely to the laboratory. The AC depth and pressure is checked 30 min after the paracentesis. The patient is started on a 3-day course of topical chloramphenicol TDS.

The complications of AC paracentesis are well known and include endophthalmitis, corneal abscess, hyphaema, and inadvertent lens touch.^{1,2} Traditionally, AC paracentesis has been performed with a tuberculin syringe, with or without a plunger attached to a hypodermic needle. When using a syringe without a plunger, there is often loss of aqueous when transferring the sample to a needle-free container. The minim technique does not eliminate these risk factors and potential complications, but does eliminate the awkward, often cumbersome manipulation associated with using an attached syringe for aspiration. The O'Rourke pipette (DORC) for aqueous sampling consists of a shorter $\frac{1}{4}$ inch 30-gauge needle attached to a deformable pipette, and is similar in its methodology of a vacuum-assisted aspiration of sample from the AC.³ Although

Figure 1 A 25-gauge needle securely attached to a saline minim.

acknowledging the increased length of the needle attached to the minim, and the potential for damage to the intraocular structures may be disadvantageous, the ability to cap the sample is clearly advantageous, both in eliminating loss of sample and in its future processing. Each minim contains 0.5 ml of volume. The volume of the AC is 0.3 ml. It is important therefore, that the minim is slowly released between thumb and forefinger, and that the entire volume of the AC is not lost. The use of 25-, 27-, and 30-gauge needles has been described earlier.²⁻⁷ The equipment needed for the minim technique can be found in any ophthalmology outpatient department. Furthermore, the ease of subsequent sample manipulation and aliquoting may prevent post-retrieval loss during processing.

References

- 1 Helbig H, Noske W, Kleineidam M, Kellner U, Foerster MH. Bacterial endophthalmitis after anterior chamber paracentesis. *Br J Ophthalmol* 1995; **79**(9): 866.
- 2 Cheung CM, Durrani OM, Murray PI. The safety of anterior chamber paracentesis with uveitis. *Br J Ophthalmol* 2004; **88**(4): 582-583.
- 3 O'Rourke J, Taylor DM, Wang Y. Compact bulb pipette simplifies paracentesis. *Ophthalmic Surg Lasers Imaging* 2004; **35**: 172-173.
- 4 Finger PT, Papp C, Latkany P, Kurli M, Iacob CE. Anterior chamber paracentesis cytology (cytospin technique) for the diagnosis of intraocular lymphoma. *Br J Ophthalmol* 2006; **90**(6): 690-692.
- 5 May DR, Noll FG. An improved approach to paracentesis. *Ophthalmic Surg* 1988; **19**(11): 821-822.
- 6 O'Rourke J, Taylor DM, Mc Donald P, Kreutzer DL. An aqueous paracentesis pipette. *Ophthalmic Surg* 1991; **22**(3): 166-167.
- 7 Van der Lelij A, Rothav A. Diagnostic anterior chamber paracentesis in uveitis: a safe procedure? *Br J Ophthalmol* 1997; **81**(11): 976.

MS Wertheim, PP Connell, MA Majid and AD Dick

Bristol Eye Hospital, Bristol, Avon, UK
 E-mail: drwertie@hotmail.com

Eye (2009) **23**, 1491; doi:10.1038/eye.2008.231;
 published online 8 August 2008

Sir,
Effect of systemic administration of valsartan, an angiotensin II type 1 receptor blocker, on retinal circulation in healthy humans

Recent studies have reported that inhibition of the renin-angiotensin system can reduce the progression of diabetic retinopathy.¹ We reported earlier that an angiotensin II type 1 receptor blocker can inhibit increased leukocyte entrapment in the retina of diabetic rats.² In addition, treatment with either an angiotensin-converting enzyme inhibitor or an angiotensin II type 1 receptor blocker normalized retinal blood flow in diabetic rats.³ These results suggest that renin-angiotensin system inhibition blocks a diabetes-induced

Table 1 Changes in systemic and retinal circulatory parameters in both groups

	Baseline		90 min		7 days	
	Valsartan	Placebo	Valsartan	Placebo	Valsartan	Placebo
Systolic BP (mmHg)	111.0 ± 2.9	108.5 ± 2.5	103.3 ± 2.5	107.8 ± 2.6	105.5 ± 2.5	104.8 ± 2.6
Diastolic BP (mmHg)	58.3 ± 2.6	56.8 ± 2.5	55.8 ± 2.2	60.3 ± 1.9	56.4 ± 1.5	58.9 ± 2.2
Mean BP (mmHg)	75.9 ± 2.4	75.2 ± 2.0	71.6 ± 2.1	74.5 ± 2.3	72.8 ± 1.6	76.3 ± 2.5
Intraocular pressure (mmHg)	17.4 ± 0.5	16.6 ± 0.5	15.8 ± 0.5	16.2 ± 0.6	15.3 ± 0.8*	16.8 ± 0.6
Ocular perfusion pressure (mmHg)	33.2 ± 1.8	31.6 ± 1.5	31.9 ± 1.5	33.3 ± 1.5	33.2 ± 1.6	31.4 ± 1.1
Vessel diameter (μm)	108.3 ± 3.6	107.7 ± 3.4	110.0 ± 3.8	109.3 ± 3.7	109.8 ± 4.2	110.5 ± 4.5
Blood velocity (mm/s)	34.9 ± 1.7	37.1 ± 2.4	33.7 ± 2.5	36.5 ± 1.9	33.2 ± 2.7	35.8 ± 2.2
Retinal blood flow (μl/min)	9.8 ± 0.8	10.3 ± 1.1	9.7 ± 0.9	10.5 ± 0.9	10.3 ± 1.3	10.5 ± 1.0

Values are expressed as the mean ± standard error of the mean.

* $P < 0.05$ compared with baseline.

BP = blood pressure.

abnormality in the retinal vasculature, thereby preserving the retinal hemodynamics. However, no study has evaluated the effect of any angiotensin II type 1 receptor blockers on the retinal circulation in humans. Therefore, we evaluated how systemic valsartan affects the retinal circulation in healthy subjects.

The effects of systemic administration of valsartan (160 mg) or placebo on the retinal circulation of 12 healthy men (mean age, 22.0 ± 0.7 years (mean ± SE); range, 20–24 years) after 90 min and 7 days were studied in a placebo-controlled, double-masked clinical trial. We used laser Doppler velocimetry (Canon Laser Blood Flowmeter; CLBF model 100, Canon, Tokyo, Japan), which has been described in detail elsewhere⁴ to measure vessel diameter and blood velocity in the retinal arterioles and estimate the retinal blood flow.

In the valsartan group, the systemic blood pressure tended to decrease, but the difference did not reach significance. (Table 1) The intraocular pressure decreased slightly but significantly from 17.4 ± 0.5 mmHg at baseline to 15.3 ± 0.8 mmHg 7 days ($P < 0.01$) after administration of valsartan. However, the calculated ocular perfusion pressure did not change significantly after the administration of valsartan in either phases of the study. Ninety minutes and 7 days after the administration of valsartan, the diameter, velocity, and retinal blood flow did not change significantly compared with the baseline values in the retinal arteries. The reduction of intraocular pressure (2 mmHg) induced by valsartan seemed to have little effect on the results of the retinal circulation in this study because the calculated ocular perfusion pressure did not change significantly after administration of valsartan.

The current results indicate for the first time that short-term and long-term systemic administration of valsartan has little effect on the retinal blood flow in healthy humans. The current results from the retinal circulation seem comparable with the results from the choroidal and retrobulbar circulation using another angiotensin II type 1 receptor blockers losartan, whereas valsartan has a much higher affinity for the angiotensin II type 1 receptor than losartan.⁵ Further study is needed to elucidate the effect of angiotensin II type 1 receptor blockers on the retinal circulation in patients with disease states, ie, hypertension, diabetes mellitus, or both.

Acknowledgements

This work was supported by a Grant-in-Aid for Young Scientists (B) 16791037 and (A) 19689035 (TN); Grant-in-Aid for scientific Research (C) 18591904 (AY). We indicate no financial conflict of interest. TN and AY are involved in the design of study, ES in the analysis of interpretation, TN in writing the article, AY in the final approval of the article, and TN, AT, and HY in data collection. This study adhered to the tenets of the Declaration of Helsinki and the Good Clinical Practice guidelines.

References

- 1 UK Prospective Diabetes Study Group. Efficacy of atenolol and captopril in reducing risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 39. *BMJ* 1998; **317**: 713–720.
- 2 Mori F, Hikichi T, Nagaoka T, Takahashi J, Kitaya N, Yoshida A. Inhibitory effect of losartan, an AT1 angiotensin II receptor antagonist, on increased leucocyte entrapment in retinal microcirculation of diabetic rats. *Br J Ophthalmol* 2002; **86**: 1172–1174.
- 3 Horio N, Clermont AC, Abiko A, Abiko T, Shoelson BD, Bursell SE *et al.* Angiotensin AT(1) receptor antagonism normalizes retinal blood flow and acetylcholine-induced vasodilatation in normotensive diabetic rats. *Diabetologia* 2004; **47**: 113–123.
- 4 Nagaoka T, Yoshida A. Noninvasive evaluation of wall shear stress on retinal microcirculation in humans. *Invest Ophthalmol Vis Sci* 2006; **47**: 1113–1119.
- 5 Criscione L, de Gasparo M, Buhlmayer P, Whitebread S, Ramjoue HP, Wood J. Pharmacological profile of valsartan: a potent, orally active, nonpeptide antagonist of the angiotensin II AT1-receptor subtype. *Br J Pharmacol* 1993; **110**: 761–771.

T Nagaoka, A Takahashi, E Sato, H Yokota, N Izumi and A Yoshida

Department of Ophthalmology, Asahikawa Medical College, Asahikawa, Japan
E-mail: nagaoka@asahikawa-med.ac.jp

Eye (2009) **23**, 1491–1492; doi:10.1038/eye.2008.274; published online 5 September 2008