manufacturer had to withdraw its retractable dog leashes after a recoil injury caused partial blindness in a 13-year-old girl.⁴ Subsequently manufacturers have advised securing the dog leash with a secondary security collar.⁵ Although this may appear to be an inconvenient second step, the benefits far outweigh the risk of permanent sight loss. It is also important to choose a type of leash that is adjusted to the size, weight, and force of the animal. When using leashes without the secondary security collar, eye protection is strongly recommended.

Conflict of interest

The authors declare no conflict of interest.

References

- 1 Jensen H. Expandable tie down straps for luggage racks dangerous for the eyes. *Nord Med* 1983; **98**: 296–297.
- 2 Gary RH, Menage MJ, Cook SD, Harcourt J. Eye injuries caused by elastic straps. *BMJ* 1988; **296**: 1097–1098.
- 3 Nichols CJ, Boldt HC, Mieler WF, Han DP, Olsen K. Ocular injuries caused by elastic cords. *Arch Ophthalmol* 1991; **109**: 371–372.
- 4 US Consumer Product Safety Commission. Worldwide Inc recalls retractable dog leashes; metal clasp can break and cause facial and body injuries to dog owners. 2008. <http://www.cpsc.gov/cpscpub/prerel/prhtml08/08387.html> (accessed 29 April 2011).
- 5 The Flexi Dog Leash Manufacturer. Retractable leash warnings. <http://www.flexiusa.com/operation/safety-advice.php> (accessed 21 August 2011).

S Waqar and P Simcock

West of England Eye Unit, Royal Devon and Exeter Hospital, Exeter, UK
E-mail: salmanwqr@gmail.com

Eye (2012) **26**, 617–618; doi:10.1038/eye.2011.352;
published online 6 January 2012

Sir,
Comment on ‘Phakic intraocular lenses outcomes and complications: Artisan vs Visian ICL’

We read with interest Hassaballa and Macky’s¹ article on phakic intraocular lenses (p-IOL) outcomes and complications. In their series significant complications resulting in elevation of intraocular pressure (IOP) requiring additional surgery occurred in two eyes (3%). As noted there are several possible mechanisms for increased IOP in patients implanted with a p-IOL. If IOP remains uncontrolled removal of a p-IOL may potentiate further pigment dispersion. There are case reports of intractable IOP elevation presumed secondary to pigment dispersion requiring surgery.^{2,3} We performed non-penetrating glaucoma surgery for a similar case of bilateral severe IOP elevation post p-IOL implantation in a patient with high myopia.³ The authors report pigment dispersion occurring in 28.6% (Artisan) and 15.38% (Visian) in their series. They did not document whether all patients underwent pre- and post-operative

gonioscopy to examine the anterior chamber angle. Stulting *et al*⁴ reported an incidence of iris pigment precipitates of 6.9%. A long-term incidence of pigment dispersion of 6.6% has also been described.⁵ This may be of significance as patients undergoing p-IOLs are generally from a younger demographic.

As the article stands, we agree with the authors on the need for long-term data on the efficacy and complications of p-IOLs. Evaluation of the possible effect on trabecular function from p-IOL-related pigment dispersion is warranted. Pigment dispersion is not always benign.

Conflict of interest

The authors declare no conflict of interest.

References

- 1 Hasaballa MA, Macky TA. Phakic intraocular lenses outcomes and complications: Artisan vs Visian ICL. *Eye* 2011; **25**: 1365–1370.
- 2 Sanchez-Galeana CA, Zadok D, Montes M, Cortés MA, Chayet AS. Refractory intraocular pressure increase after phakic posterior chamber intraocular lens implantation. *Am J Ophthalmol* 2002; **134**: 121–123.
- 3 Toh T, Nguyen DQ, Ruddle JB, Crowston JG, Coote MA. Bilateral acute severe intraocular pressure elevation requiring filtration surgery after implantation of toric Artisan phakic intraocular lenses. *Clin Exp Ophthalmol* 2011; e-pub ahead of print 30 June 2011; doi:10.1111/j.1442-9071.2011.02646.x.
- 4 Stulting RD, John ME, Maloney RK, Assil KK, Arrowsmith PN, Thompson VM. Three-year results of Artisan/Verisyse phakic intraocular lens implantation; results of the United States Food and Drug Administration Clinical Trial; the U.S. Verisyse Study Group. *Ophthalmology* 2008; **115**: 464–472.
- 5 Menezo JL, Menezo JL, Peris-Martínez C, Cisneros AL, Martínez-Costa R. Phakic intraocular lenses to correct high myopia: Adatomed, Staar, and Artisan. *J Cataract Refract Surg* 2004; **30**: 33–44.

DQ Nguyen^{1,2}, TY Toh^{3,4}, JB Ruddle^{1,2}, JG Crowston^{1,2} and MC Coote^{1,2}

¹Royal Victorian Eye and Ear Hospital, Melbourne, Australia

²Centre of Eye Research Australia, Melbourne, Australia

³Launceston Eye Institute, South Launceston, Tasmania

⁴The Eye Hospital, Launceston, Tasmania
E-mail: danqbnguyen@hotmail.com

Eye (2012) **26**, 618; doi:10.1038/eye.2011.355;
published online 6 January 2012

Sir,
Response to: Comment on ‘Phakic intraocular lenses outcomes and complications: Artisan vs Visian ICL’

We would like to thank Nguyen *et al*¹ for their valuable comments. Our study² is a retrospective one; in which, we reported in the ‘Material and methods’ section only the list of examinations and investigations done to all patients. In our study, not all patients had a

preoperative gonioscopy examination records in their files. However, anterior chamber (AC) depth measurements were done for all patients, and those with an AC depth less than 3 mm were excluded, and none of our patients had a narrow angle. We do agree that a pre-operative gonioscopy is essential in these cases, both for angle width assessment as well as exclusion of pigment deposits in the angle. The two glaucoma cases reported in our series were successfully treated with pars planavitreotomy, and a reduced-size pIOL exchange for the malignant glaucoma case with Artisan and the oversized Visian ICL case, respectively. Both cases developed in the first few days post-operative, and were closely followed up for 1 year with no further IOP rise. This excludes the possibility of pigment dispersion as an etiology. We do not have an explanation for the increased incidence of pigment dispersion in our series compared with others,^{3,4} but the possibility of a different ethnic and/or racial patient population should be considered.

Conflict of interest

The authors declare no conflict of interest.

References

- 1 Nguyen DQ, Toh TY, Ruddle JB, Crowston JG, Coote MC. Comment on 'Phakic intraocular lenses outcomes and complications: Artisan vs Visian ICL'. *Eye* 2012; **26**: 618.
- 2 Hasaballa MA, Macky TA. Phakic intraocular lenses outcomes and complications: Artisan vs Visian ICL. *Eye* 2011; **25**: 1365–1370.
- 3 Stulting RD, John ME, Maloney RK, Assil KK, Arrowsmith PN, Thompson VM. Three year results of Artisan/Verisyse phakic intraocular lens implantation; results of the United States Food and Drug Administration Clinical Trial; the U.S. Verisyse Study Group. *Ophthalmology* 2008; **115**: 464–472.
- 4 Menezo JL, Menezo JL, Peris-Martínez C, Cisneros AL, Martínez-Costa R. Phakic intraocular lenses to correct high myopia: Adatomed, Staar, and Artisan. *J Cataract Refract Surg* 2004; **30**: 33–44.

TA Macky and MA Hasaballa

Department of Ophthalmology, Cairo University
Hospitals, Cairo, Egypt
E-mail: tamer Macky@gmail.com

Eye (2012) **26**, 618–619; doi:10.1038/eye.2011.356;
published online 6 January 2012

Sir, Combined OCT and colour fundus photography in virtual clinic assessments of wet AMD patients

We would like to commend Hibbs *et al*¹ on their study emphasising the need for colour photography in screening patients for wet age-related macular degeneration (AMD). We agree with the authors that colour photography is essential, but would like to describe a service evaluation performed at Hull and East Yorkshire Eye Hospital. Data from this evaluation suggested that

even with colour photography combined with optical coherence tomography (OCT), wet AMD patients with retinal haemorrhage and no other features of disease activity can be under-treated without slit lamp assessment.

In our evaluation, we used spectral domain OCT with associated colour fundus photography (Topcon 3D OCT-1000) to examine 242 patients with wet AMD undergoing follow-up, to assess the need for Ranibizumab therapy. Clinicians were asked to record re-treatment decisions based on imaging and ETDRS visual acuity only as a 'virtual' assessment. Each patient was then immediately reviewed with slit lamp examination (SLE) and any change in management decision documented.

A total of 242 patients were evaluated and the results were as follows.

For 200 patient episodes, SLE did not provide any other useful clinical information. For 42 patients, the SLE provided additional information when compared with virtual assessment. We further examined these questionnaires.

In 21 patients ocular pathology unrelated to Ranibizumab therapy was detected, whereas in 6 patients OCT quality was too poor to allow virtual assessment. Of these 27 patients, SLE made no change in management of any ophthalmic disease for 11 patients. In our cohort, only 15 of the 242 (6.2%) patients had haemorrhage on SLE that was otherwise not detected on OCT/colour photography alone, but in only 6/15 (2.5% of the total) were there no other associated OCT features indicating disease activity.

We concluded that in our cohort 2.5% of patients would require SLE to detect isolated retinal haemorrhage undetected with colour imaging where there were no other OCT features of disease activity. However, in the context of the capacity issues associated with reviewing all patients on a monthly basis, it was felt that this was an acceptable percentage when set against the benefits of higher throughput in virtual clinics.

Conflict of interest

The authors declare no conflict of interest.

Reference

- 1 Hibbs SP, Smith A, Chow LP, Downes SM. Colour photographs for screening in neovascular age-related macular degeneration: are they necessary? *Eye* 2011; **25**(7): 918–921.

M Mookhtiar¹ and L Downey²

¹St James University Teaching Hospital,
Department of Ophthalmology, Leeds Teaching
Hospitals NHS Trust, Leeds, UK

²Hull & East Yorkshire Eye Hospital, Kingston upon
Hull, UK

E-mail: Murtuza@mookhtiar.co.uk

Some of the data has been presented as a poster presentation at Royal College of Ophthalmologists Annual Congress 2010.

Eye (2012) **26**, 619; doi:10.1038/eye.2011.345;
published online 13 January 2012