

A novel compound heterozygous mutation in the *BEST1* gene causes autosomal recessive Best vitelliform macular dystrophy

L Zhao^{1,2,3,5}, S Grob^{2,3,5}, R Corey⁴, M Krupa^{2,3}, J Luo^{2,3}, H Du^{2,3}, C Lee^{2,3}, G Hughes^{2,3}, J Lee^{2,3}, J Quach^{2,3}, J Zhu^{1,2,3}, PX Shaw^{2,3}, I Kozak² and K Zhang^{1,2,3}

¹Molecular Medicine Research Center and Department of Ophthalmology, West China Hospital, Sichuan University, Chengdu, China

²Department of Ophthalmology, Shiley Eye Center, University of California, San Diego, La Jolla, CA, USA

³Institute for Genomic Medicine, University of California, San Diego, La Jolla, CA, USA

⁴Vitreous and Retina Surgeons of Utah, Provo, UT, USA

Correspondence: K Zhang, Department of Ophthalmology, Shiley Eye Center, and Institute for Genomic Medicine, University of California, San Diego, 9415 Campus Point Drive Mail Code 0946 Room E214, La Jolla, CA 92093-0838, USA
 Tel: +1 858 246 0823;
 Fax: +1 858 246 0873.
 E-mail: kang.zhang@gmail.com

⁵These authors contributed equally to this work.

Received: 29 September 2011

Accepted in revised form: 29 January 2012

Published online: 16 March 2012

Abstract

Purpose To determine the genetic basis of early onset autosomal recessive Best vitelliform macular dystrophy (arBVMD) in a family with three affected children.

Design Clinical and family-based genetic study.
Methods Seven subjects making up a family with three children affected by Best vitelliform macular dystrophy were studied. Standard ophthalmic exam with dilated ophthalmoscopy and imaging were performed in each individual. The eleven exons of *BEST1* were directly sequenced.

Results All three affected children have the clinical characteristic features of Best vitelliform macular dystrophy: large macular vitelliform lesions, scattered vitelliform lesions along the arcades and in the peripheral retina, and an accumulation of serous retinal fluid. A novel compound heterozygous mutation in the *BEST1* gene was found in the three affected individuals (L41P and I201T). The unaffected parents and children only harbor one heterozygous mutation.

Conclusion arBVMD can be caused by the compound heterozygous mutation L41P and I201T in the *BEST1* gene.

Eye (2012) 26, 866–871; doi:10.1038/eye.2012.27; published online 16 March 2012

Keywords: autosomal recessive Best vitelliform macular dystrophy; *BEST1* gene; genetics

Introduction

Autosomal dominant Best vitelliform macular dystrophy (BVMD) is one of the most common

retinal dystrophies. The disease is characterized clinically by deposition of yellowish material in the retinal pigment epithelium (RPE) and subretinal space. This accumulated material forms the characteristic egg yolk-like appearance in the macula. While the autosomal dominant pattern of inheritance has been long observed, an autosomal recessive pattern has also been reported recently.¹ Homozygous or compound mutations in the *BEST1* gene can cause autosomal recessive BVMD (arBVMD).^{1–6}

BEST1 is located on chromosome 11q13 and encodes the bestrophin-1 protein that localizes to the RPE.^{1,7,8} The exact function of the bestrophin protein is not completely understood. However, it has been suggested that bestrophin-1 acts as either a chloride channel or a modulator of calcium channels.^{6,8–10}

To date, several mutations in arBVMD cases have been reported.^{1–3,5,11,12} These include homozygous and combinations of nonsense or missense mutations. In all of these cases, patients presented with central vision loss, characteristic retinopathy, absence of electro-oculogram light rise, and a reduced electroretinogram.¹ In this report, we present a novel compound heterozygous mutation in a family with three affected children.

Materials and methods

Participants and clinical examinations

The family being studied consisted of seven individuals, including two parents and five children. Three of the children complained of visual impairment. The parents and two of the children denied any visual symptoms. Both of the parents, the three affected children, and two

non-affected children had their blood drawn for genetic analysis.

Each family member received a complete ophthalmic examination. Exams included best-corrected visual acuity measurements and applanation tonometry for intraocular pressure measurements. Slit lamp exams included fundus biomicroscopy. In addition, fundus photography and spectral domain optical coherence tomography were completed.

DNA sequencing

Genomic DNA samples were extracted from peripheral blood leukocytes according to established protocols from seven of the family members. Direct DNA sequence analysis was completed for the *BEST1* gene. The eleven exons of the *BEST1* gene were amplified by PCR and sequenced using the Genetic Analyzer 3130 system (Applied Biosystems, Foster City, CA, USA). Primers used to amplify the exons in the *BEST1* gene are presented (Table 1).

Results

Clinical examinations

The family under study consisted of seven individuals, including two non-consanguineous Caucasian parents and five children. Three of the five children were diagnosed with arBVMD at age 4.

Child 1 (II-1 in Figure 1a) is a twelve-year-old female with best-corrected visual acuities of 20/400 OD and 20/30 OS. Dilated ophthalmic exam is significant for macular lesions and smaller scattered vitelliform lesions in the periphery of the right eye, intraretinal cysts, and elevation and separation of the neurosensory retina from the RPE in the left eye (Figure 2).

Child 3 (II-3 in Figure 1a) is a nine-year-old male with best-corrected visual acuities of 20/30 OD and 20/40 OS.

Dilated ophthalmic exam is significant for scattered vitelliform lesions, subretinal deposits, and subretinal fluid in the macula of the right eye (Figure 3).

Child 4 (II-4 in Figure 1a) is a five-year-old female with best-corrected visual acuities of 20/40 OD and 20/30 OS. Dilated ophthalmic exam is significant for bilateral yellowish, subretinal macular lesions with the lesion in the right eye extending outside the macular area down to the inferior arcade. There are some small deposits noted in the left macula and serous retinal fluid in both eyes.

The parents (I-1 and I-2) and the other two unaffected children (II-2 and II-5) have no clinical signs of disease (Figure 4).

Genetic analysis

The family history is consistent with an autosomal recessive inheritance pattern (Figure 1a). We found the unaffected father to have a T>C transition (c.122T>C) in exon 5, resulting in a L41P substitution (Figure 1b). The unaffected mother, however, was found to have a different variation, consisting of a T>C transition (c.602T>C) in exon 2, resulting in an I201T substitution (Figure 1c). The three affected children harbor both of these variations while the two unaffected children carry only one of the heterozygous mutations. Table 2 summarizes the genetic analysis results for each family member. These two changes were absent in 1000 normal control chromosomes.

Discussion

We report a family with an autosomal recessive BVMD that resulted from compound heterozygous *BEST1* mutations.¹ In these cases, the unaffected family members carry only one of the heterozygous mutations. While over 100 different *BEST1* mutations have been

Table 1 Sequencing primers list for *BEST1* gene

Exon	Forward (5'–3')	Reverse (5'–3')	PCR products (bp)
1	GAGGCTGAGAGAGGAGCTGA	CCATGGCCCCTCTAATTCT	363
2	GAGGTCCAGAGCAGGGAAGGGT	CAGCCCCAGCCACATCCTT	327
3	GAGGCAGTCCCACCTCTACC	GCAGCTCCTCGTGATCCTC	278
4	CTCCTGCCAGGCTTCTACGT	CCACCCATCTCCATTCT	326
5	GGTTCCTATAGGTCAGCAGGTG	GAAACCTTGTTTCCTGTGGAC	303
6	TGGTACCTGGAGAAGAGGTG	CCTTGGTCCTTCTAGCCTCA	219
7	CATCTGATTCAGGGTTCC	GACACTGCATCCTCGTCTCA	298
8	ATGGGGTGTGAAAATAGCAG	GAGGGGAAGGGTTGATCATT	290
9	CTCCAAGTCATCAGGCACAT	GCAGACCCCTGCACTAGGAG	284
10-1	GGTGTGGTCCTTTGTCCAC	TGACACTGTGAAGCTTTGACG	430
10-2	CTGGAAGCTTAAGGCTGTGG	TAGGCTCAGAGCAAGGGAAG	481
11	CTTTGCCCTCCTACTGCAAC	TCCTTAAGTGCCGTTGTCA	487




Figure 1 Pedigree and the *BEST1* sequencing results for a family affected by arBVMD. (a) Pedigree of the family affected by arBVMD. Father (I-1), mother (I-2), and the two children (II-2 and II-5) are normal; each one harbors only one mutation (L41P or I201T). The other three children (II-1, II-3, and II-4) are affected by arBVMD and have the compound mutation (L41P and I201T). (b) Sequencing data show that a T>C transition in exon 5 (arrow) causes the L41P mutation and (c) that a T>C transition in exon 2 (arrow) causes the I201T mutation.




Figure 2 (II-1) Fundus photographs (a, c) of both eyes significant for macular lesions surrounded by retinal elevation bilaterally. Multiple scattered vitelliform lesions are noted in the periphery outside the arcades. Optical coherence tomography images (b, d) are shown through the macular lesions. Significant distortion of the macular structure is noted with elevation and separation from the RPE and the presence of intraretinal cysts and subretinal fluid (c, d).

described in families affected by Best disease,^{13–17} only a few combinations of compound heterozygous mutations in arBVMD have been reported.^{1–5,11}

In our study, the affected children were found to have a combination of two heterozygous mutations: c.122T>C (L41P) and c. 602T>C (I201T). The L41P mutation has




Figure 3 (II-3) Fundus photographs (a, c) of both eyes significant for many small, scattered vitelliform lesions in the macula and larger vitelliform lesions along the arcades and in the periphery. Additionally, there are macular vitelliform lesions in both eyes with serous retinal fluid in the macula of the right eye. Optical coherence tomography images (b, d) through the macular and peripheral lesions are also shown. The peripheral lesion in the right eye (a, b) is located nasal to the macula (white arrow upper left panel), and the retina is also separated from the RPE. The scan through the macular lesion (d) shows elevation of the macular area with intraretinal fluid.




Figure 4 (I-2) Fundus photographs (a, c) and optical coherence tomography images (b, d) of both eyes reveal normal anatomy and no signs of retinal disease.

Table 2 Mutations in the *BEST1* gene and the associated clinical findings in patients with autosomal recessive Best vitelliform macular dystrophy

Family member	Age/gender	BCVA OD	BCVA OS	Mutation	Phenotype
Mother	35/F	20/20	20/20	I201T/+	Normal
Father	36/M	20/15	20/15	L41P/+	Normal
Child 1	12/F	20/400	20/30	L41P/I201T	arBVMD
Child 2	11/M	20/20	20/15	L41P/+	Normal
Child 3	9/M	20/30	20/40	L41P/I201T	arBVMD
Child 4	5/F	20/40	20/30	L41P/I201T	arBVMD
Child 5	4/M	20/25	20/25	I201T/+	Normal

been previously implicated in arBVMD as a compound heterozygous mutation with P152A in one case, and with R141H in a second case.¹ The I201T mutation has also been implicated in arBVMD. To date, three individual cases of a compound heterozygous mutation with I201T have been reported: I201T with P152A, I201T with IVS 7 + 4G > A, and I201T with the deletion Phe281del3CAGTTC.¹ We have shown for the first time that the combination of I201T with L41P can also lead to arBVMD.

In vitro studies using HEK293 cells have shown that co-transfection of the two mutations observed in the compound heterozygous state in arBVMD patients essentially abolished chloride channel activity. However, when six arBVMD mutations were coexpressed with wild-type bestrophin-1, none of the combinations suppressed the wild-type chloride channel activity.¹ Therefore, it is suggested that the autosomal recessive phenotype only manifests when the bestrophin-1 activity drops below a functional threshold.^{1,18} The mutations that make up the autosomal recessive phenotype do not cause enough structural change in the protein to bring its function below threshold alone in the carrier. They must be combined with other mutations to affect protein function.

The phenotype of BVMD varies with each patient and the stage of the disease. Initial stages are characterized by a circular or horizontally elliptical vitelliform lesion centered around the fovea and ranging in diameter from 0.5 disk diameter to 2 disk diameters, simulating the appearance of an egg yolk.¹⁹ Over the years, the material in the subretinal space may become less homogenous, giving rise to what is known as the 'scrambled-egg' stage, or it may gravitate inferiorly, forming the 'pseudohypopyon' stage.²⁰ Previous publications have discussed that the clinical presentation of cases of compound arBVMD are noticeably different from the autosomal dominant form.^{1,11,13} The most-reported distinguishing feature of arBVMD is extrafoveal and extramacular subretinal deposits.

Additionally, patients with BVMD are at increased risk for choroidal neovascularization. These membranes are often difficult to detect owing to the overlying vitelliform lesions.³ There have been several reports of favorable outcomes with the use of anti-VEGF therapy.^{21,22} This has turned out to be consistent with our study, where the patients' sub- or intraretinal fluid has resolved from injections of bevacizumab.

Our finding adds to the spectrum of clinical presentations caused by *BEST1* mutations. Understanding the genetics of the bestrophinopathies and the cause of different forms of the disease will allow for the development of potential gene therapy regimens and new diagnostic modalities.

Summary

What was known before

- Autosomal recessive BVMD can result from compound heterozygous *BEST1* mutations.

What this study adds

- A novel combination of two heterozygous *BEST1* gene mutations causes autosomal recessive BVMD.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

This study was approved by the Institutional Review Board of the University of California, San Diego. All subjects signed informed consent before participation in the study. We thank the family for their support. This study was supported by NEI/NIH grants, Research to Prevent Blindness, San Diego Clinical and Translational Research Institute 1TL1RR031979-01, and the VA Merit Award.

Access to data

The principal investigator, Dr Kang Zhang, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Author contributions

Design of the study (LZ, SG and KZ); conduct of the study (LZ, SG, MK, JL, HD and GH); data collection (SG, RC, JL, CL and IK); analysis and interpretation of the data (LZ, SG, MK, JQ, JZ, IK, PXS and KZ); and writing of the manuscript (LZ, SG, MK, PXS, IK and KZ).

References

- 1 Burgess R, Millar ID, Leroy BP, Urquhart JE, Fearon IM, De Baere E *et al*. Biallelic mutation of *BEST1* causes a distinct retinopathy in humans. *Am J Hum Genet* 2008; **1**: 19–31.
- 2 Kinnick TR, Mullins RF, Dev S, Leys M, Mackey DA, Kay CN *et al*. Autosomal recessive vitelliform macular dystrophy in a large cohort of vitelliform macular dystrophy patients. *Retina* 2011; **3**: 581–595.
- 3 Iannaccone A, Kerr NC, Kinnick TR, Calzada JI, Stone EM. Autosomal recessive best vitelliform macular dystrophy: report of a family and management of early-onset neovascular complications. *Arch Ophthalmol* 2011; **2**: 211–217.
- 4 Davidson AE, Sergouniotis PI, Burgess-Mullan R, Hart-Holden N, Low S, Foster PJ *et al*. A synonymous codon variant in two patients with autosomal recessive bestrophinopathy alters *in vitro* splicing of *BEST1*. *Mol Vis* 2010; **16**: 2916–2922.

- 5 Gerth C, Zawadzki RJ, Werner JS, Heon E. Detailed analysis of retinal function and morphology in a patient with autosomal recessive bestrophinopathy (ARB). *Doc Ophthalmol* 2009; **3**: 239–246.
- 6 Hartzell C, Qu Z, Putzier I, Artinian L, Chien LT, Cui Y. Looking chloride channels straight in the eye: bestrophins, lipofuscinosis, and retinal degeneration. *Physiology (Bethesda)* 2005; **20**: 292–302.
- 7 Petrukhin K, Koisti MJ, Bakall B, Li W, Xie G, Marknell T *et al*. Identification of the gene responsible for Best macular dystrophy. *Nat Genet* 1998; **3**: 241–247.
- 8 Sun H, Tsunenari T, Yau KW, Nathans J. The vitelliform macular dystrophy protein defines a new family of chloride channels. *Proc Natl Acad Sci USA* 2002; **6**: 4008–4013.
- 9 Tsunenari T, Sun H, Williams J, Cahill H, Smallwood P, Yau KW *et al*. Structure-function analysis of the bestrophin family of anion channels. *J Biol Chem* 2003; **42**: 41114–41125.
- 10 Rosenthal R, Bakall B, Kinnick T, Peachey N, Wimmers S, Wadelius C *et al*. Expression of bestrophin-1, the product of the VMD2 gene, modulates voltage-dependent Ca²⁺ channels in retinal pigment epithelial cells. *FASEB J* 2006; **1**: 178–180.
- 11 Schatz P, Klar J, Andreasson S, Ponjavic V, Dahl N. Variant phenotype of Best vitelliform macular dystrophy associated with compound heterozygous mutations in VMD2. *Ophthalmic Genet* 2006; **2**: 51–56.
- 12 Boon CJ, Klevering BJ, Leroy BP, Hoyng CB, Keunen JE, den Hollander AI. The spectrum of ocular phenotypes caused by mutations in the BEST1 gene. *Prog Retin Eye Res* 2009; **3**: 187–205.
- 13 Lotery AJ, Munier FL, Fishman GA, Weleber RG, Jacobson SG, Affatigato LM *et al*. Allelic variation in the VMD2 gene in best disease and age-related macular degeneration. *Invest Ophthalmol Vis Sci* 2000; **6**: 1291–1296.
- 14 Kramer F, Mohr N, Kellner U, Rudolph G, Weber BH. Ten novel mutations in VMD2 associated with Best macular dystrophy (BMD). *Hum Mutat* 2003; **5**: 418.
- 15 Caldwell GM, Kakuk LE, Griesinger IB, Simpson SA, Nowak NJ, Small KW *et al*. Bestrophin gene mutations in patients with Best vitelliform macular dystrophy. *Genomics* 1999; **1**: 98–101.
- 16 Downs K, Zacks DN, Caruso R, Karoukis AJ, Branham K, Yashar BM *et al*. Molecular testing for hereditary retinal disease as part of clinical care. *Arch Ophthalmol* 2007; **2**: 252–258.
- 17 Palomba G, Rozzo C, Angius A, Pierrottet CO, Orzalesi N, Pirastu M. A novel spontaneous missense mutation in VMD2 gene is a cause of a best macular dystrophy sporadic case. *Am J Ophthalmol* 2000; **2**: 260–262.
- 18 Davidson AE, Millar ID, Burgess-Mullan R, Maher GJ, Urquhart JE, Brown PD *et al*. Functional characterization of bestrophin-1 missense mutations associated with autosomal recessive bestrophinopathy. *Invest Ophthalmol Vis Sci* 2011; **6**: 3730–3736.
- 19 Blodi CF, Stone EM. Best's vitelliform dystrophy. *Ophthalmic Paediatr Genet* 1990; **1**: 49–59.
- 20 Yannuzzi LA. *The Retinal Atlas*. Saunders/Elsevier: Philadelphia, PA, 2010.
- 21 Querques G, Bocco MC, Soubrane G, Souied EH. Intravitreal ranibizumab (Lucentis) for choroidal neovascularization associated with vitelliform macular dystrophy. *Acta Ophthalmol* 2008; **6**: 694–695.
- 22 Leu J, Schrage NF, Degenring RF. Choroidal neovascularisation secondary to Best's disease in a 13-year-old boy treated by intravitreal bevacizumab. *Graefes Arch Clin Exp Ophthalmol* 2007; **11**: 1723–1725.