Acknowledgements

This work was supported by CONICET PIP 112 200801 01455; SECYT-UNC 2009 (Argentina).

References

- 1 Gray RH, Johnson GJ, Freedman A. Climatic droplet keratopathy. *Surv Ophthalmol* 1992; **36**: 241–253.
- 2 Urrets-Zavalía JA, Maccio JP, Knoll EG, Cafaro T, Urrets-Zavalía EA, Serra HM. Surface alterations, corneal hypoesthesia, and iris atrophy in patients with climatic droplet keratopathy. *Cornea* 2007; **26**: 800–804.
- 3 Muller LJ, Marfurt CF, Kruse F, Tervo TM. Corneal nerves: structure, contents and function. *Exp Eye Res* 2003; **76**: 521–542.
- 4 Niederer RL, McGhee CN. Clinical in vivo confocal microscopy of the human cornea in health and disease. *Prog Retin Eye Res* 2010; **29**: 30–58.
- 5 Patel DV, Tavakoli M, Craig JP, Efron N, McGhee CN. Corneal sensitivity and slit scanning iVCM of the subbasal nerve plexus of the normal central and peripheral human cornea. *Cornea* 2009; **28**: 735–740.

JA Urrets-Zavalía¹, JO Croxatto², JM Holopainen³,
TA Cafaro⁴, F Esposito¹, W Neira³ and HM Serra⁴

¹Department of Ophthalmology, University Clinic Reina Fabiola, Universidad Católica de Córdoba, Córdoba, Argentina

²Department of Ophthalmic Pathology, Fundación Oftalmológica Argentina Jorge Malbrán, Buenos Aires, Argentina

³Department of Ophthalmology, University of Helsinki, Helsinki, Finland

⁴CIBICI, Department of Clinical Biochemistry, Faculty of Chemical Sciences, Universidad Nacional de Córdoba, Córdoba, Argentina
E-mail: hserra@mail.fcq.unc.edu.ar

Eye (2012) **26**, 1021–1023; doi:10.1038/eye.2012.79;
published online 27 April 2012

Sir, Surgical repair of bilateral full thickness macular holes in a patient with blue sclera secondary to osteogenesis imperfecta

The management of full thickness macular holes (FTMH) in patients with osteogenesis imperfecta has not to our knowledge been reported. We describe a patient with blue sclera secondary to osteogenesis imperfecta, who underwent surgical repair of bilateral FTMH.

Figure 1 Photographs showing bilateral blue-grey scleral colouration OD (a) and OS (b). Pre-treatment OCT appearance of FTMH OD (c) and OS (d). Post-treatment OCT appearance showing closure of FTMH OD (e) and OS (f). Punched-out retinal defect on OCT in patient's son OD (g) and OS (h).

Case report

A 73-year-old lady was referred complaining of a 2-month history of reduced central vision in the right eye. Her left vision had been poor since childhood and she had not noticed a change. She was hypermetropic with spherical equivalents of +7.00 OD and +6.50 OS, and had blue sclera secondary to osteogenesis imperfecta type 1.

Examination revealed LogMAR acuities of 0.46 OD and 0.90 OS. Blue sclera was noted in both the eyes (Figures 1a and b) together with nuclear sclerotic cataract and bilateral FTMH (Figures 1c and d).

The amblyopic left eye had initial surgery in order to assess possible technical difficulties associated with the blue sclera and high hypermetropia. The left eye underwent 20 g combined phaco-vitreectomy, inner limiting membrane peel, 20% C₂F₆ gas tamponade and she postured face down for 5 days. Following successful closure of the left macular hole (Figure 1f) she underwent a similar procedure to the right eye but with 23 g instrumentation leading to successful closure of the right macula hole (Figure 1e). Postoperative corrected LogMAR acuities were 0.16 OD and 0.94 OS.

The patient's 46 year-old son, who had been registered blind since childhood was also examined. He did not have blue sclera but had a circular foveal abnormality bilaterally with punched-out retinal defects on OCT that have been described in cone dystrophy and Stargardt disease (Figures 1g and h).

Comment

Osteogenesis imperfecta is an inherited connective tissue disorder resulting from deficiency of type 1 collagen. The main ocular manifestation is blue sclera that has been shown to be associated with reduced scleral rigidity¹ and it has been proposed that this may lead to increased susceptibility to retinal detachment² and retinal haemorrhage.³ Our literature search did not reveal any reported association with FTMH.

Vitreectomy surgery in patients with blue sclera secondary to osteogenesis imperfecta has been described in the treatment of retinal detachment⁴ and vitreous opacities,⁵ the latter employing 25 g sutureless vitrectomy. We describe the treatment of FTMHs in such a patient, employing both 20 and 23 g vitrectomy. Ports were sutured in both the 20 and 23 g procedures as the surgeon (PRS) felt that the lack of scleral rigidity and thinness did not allow self-sealing of scleral tunnels. No operative complications were encountered.

Conflict of interest

The authors declare no conflict of interest.

References

- 1 Kaiser-Kupfer MI, Podgor MJ, McCain L, Kupfer C, Shapiro JR. Correlation of ocular rigidity and blue sclerae in osteogenesis imperfecta. *Trans Ophthalmol Soc UK* 1985; **104**: 191–195.

- 2 Madigan WP, Wertz D, Cockerham GC, Thach AB. Retinal detachment in osteogenesis imperfecta. *J Pediatr Ophthalmol Strabismus* 1994; **31**: 268–269.
- 3 Ganesh A, Jenny C, Geyer J, Shouldice M, Levin AV. Retinal hemorrhages in type 1 osteogenesis imperfecta after minor trauma. *Ophthalmol* 2004; **111**: 1428–1431.
- 4 Elliott D, Rezai KA, Bawa Dass A, Lewis J. Management of retinal detachment in osteogenesis imperfecta. *Arch Ophthalmol* 2003; **121**: 1062–1064.
- 5 Unver YB, Acar N, Altan T, Kapran Z, Ozturk G. 25-Gauge sutureless vitrectomy in osteogenesis imperfecta type 1. *Retin Cases Brief Rep* 2009; **3**: 286–287.

JD Benzmira, B Kapoor, N Edmeades, A Burger and PR Simcock

West of England Eye Unit, Royal Devon and Exeter Hospital, Exeter, UK
E-mail: james_benzmira@hotmail.com

Eye (2012) **26**, 1023–1024; doi:10.1038/eye.2012.63;
published online 13 April 2012

Sir,
Paediatric adjustable strabismus surgery

We commend Nihalani and Hunter¹ on an excellent review of the adjustable suture strabismus surgery literature, especially with regard to paediatric squints. As the authors mention, the benefits of adjustable sutures have been demonstrated in both adults and children, although there is a general reluctance in attempting this technique in children owing to the perception that it would not be tolerated by the patient.

Between 2007 and 2010, our experience of adjustable suture use in a paediatric population aged between 8–15 years (mean 12.3 years old) reflects the benefits of this technique. Our current practice involves a hangback technique with a 6/0 vicryl adjustable, slip-knot tie, followed by sub-Tenon's levobupivacaine (5 mg/ml). Within 2–3 h of recovery from general anaesthesia, topical tetracaine hydrochloride 1% is instilled and a prism cover test is performed. The suture is adjusted as required and the conjunctiva closed with 8-0 vicryl in the treatment room. Ninety-five percent (20/21) of our paediatric patients were able to tolerate this technique, with only one requiring a second general anaesthetic. The vast majority (90%) had a resultant angle <10 prism dioptres (Figure 1).

While we agree that a certain amount of experience is required with adjustable sutures for this population, we feel the most important factor is suitable patient selection and thorough preoperative counselling of both the patient and the parents. We find that with suitable scrutiny it is possible to identify those who will tolerate the adjustment, such that the surgeon should not feel intimidated by the prospect of adjustment in this group of patients.