

Predictive factors and outcomes of posterior segment intraocular foreign bodies

J Choovuthayakorn, L Hansapinyo, N Ittipunkul,
 D Patikulsila and P Kunavisarut

Eye (2012) 26, 484; doi:10.1038/eye.2012.8

Correction to: Eye (2011) 25, 1622–1626;
 doi:10.1038/eye.2011.229; published online
 16 September 2011

Since the publication of the above article,
 the authors have noticed that the sex ratios

in Tables 4 and 5 were labeled incorrectly.
 The correct tables are shown below.

The authors would like to apologize for this
 mistake.

Table 4 Univariate analysis: predictors for good visual outcome

Predictive factors	Final VA \geq 20/40, N (%)	Final VA < 20/40, N (%)	P-value
Mean age (years)	29.5	32.5	0.624 ^a
Sex (male:female)	6:1	46:3	0.287
Initial VA \geq 20/40	4 (66.67%)	2 (33.33%)	0.182
<i>Entry site</i>			
Cornea	22 (36.07%)	39 (63.93%)	1.000
Sclera	1 (10.00%)	9 (90.00%)	0.083
Corneosclera	5 (83.33%)	1 (16.67%)	0.022
RAPD	2 (11.11%)	16 (88.98%)	0.012
Lens injury	21 (31.82%)	45 (68.18%)	0.194
<i>Retinal injury</i>			
Break	7 (25.00%)	21 (75.00%)	0.223
RD	0 (0.00%)	5 (100.00%)	0.152
Hemorrhage	6 (28.57%)	15 (71.43%)	0.439
Endophthalmitis	0 (0.00%)	7 (100.00%)	0.044

Abbreviations: RAPD, relative afferent pupillary defect; RD, rhegmatogenous retinal detachment; VA, visual acuity.

^aMann–Whitney *U* test; in others Fisher exact test was used.

Table 5 Univariate analysis: predictors for poor visual outcome

Predictive factors	Final VA < 20/200, N (%)	Final VA \geq 20/200, N (%)	P-value
Mean age (years)	32.3	38.1	0.862 ^a
Sex (male:female)	36:1	17:3	0.084
Initial VA < 20/200	37 (56.92%)	28 (43.08%)	0.121
<i>Entry site</i>			
Cornea	26 (42.62%)	35 (57.38%)	1.000
Sclera	6 (60.00%)	4 (40.00%)	0.311
Corneosclera	1 (16.67%)	5 (83.33%)	0.230
RAPD	13 (72.22%)	5 (27.78%)	0.006
Lens injury	30 (45.45%)	36 (54.54%)	0.335
<i>Retinal injury</i>			
Break	13 (46.43%)	15 (53.57%)	0.485
RD	5 (100.00%)	0 (0.00%)	0.012
Hemorrhage	8 (38.10%)	13 (61.90%)	0.786
Endophthalmitis	7 (100.00%)	0 (0.00%)	0.002

Abbreviations: RAPD, relative afferent pupillary defect; RD, rhegmatogenous retinal detachment; VA, visual acuity.

^aMann–Whitney *U* test; in others Fisher exact test was used.