

# Relevance of TSH-receptor antibody levels in predicting disease course in Graves' orbitopathy: comparison of the third-generation TBII assay and Mc4-TSI bioassay

SY Jang<sup>1,2</sup>, DY Shin<sup>3</sup>, EJ Lee<sup>3</sup>, SY Lee<sup>4</sup>  
and JS Yoon<sup>4</sup>

*Eye* (2013) 27, 1231; doi:10.1038/eye.2013.193

**Correction to:** *Eye* (2013) 27, 964–971;  
doi: 10.1038/eye.2013.120; published online  
7 June 2013

Since the publication of the above article, the authors have noted that an affiliation is missing for the first author. The correct details are given above.

The authors would like to apologise for this error.

<sup>1</sup>Department of Ophthalmology, Soonchunhyang University Bucheon Hospital, Soonchunhyang University College of Medicine, Bucheon, Korea

<sup>2</sup>Yonsei University Graduate School of Medicine, Seoul, Korea

<sup>3</sup>Department of Endocrinology, Brain Korea 21 Project for Medical Science, Institute of Endocrine Research, and Severance Integrative Research Institute for Cerebral and Cardiovascular Disease, Yonsei University College of Medicine, Seoul, Korea

<sup>4</sup>Department of Ophthalmology, The Institute of Vision Research, Yonsei University College of Medicine, Seoul, Korea