- 4 Momota H, Narita Y, Miyakita Y, Shibui S. Secondary hematological malignancies associated with temozolomide in patients with glioma. *Neuro-oncology* 2013; **15**(10): 1445–1450.
- 5 Chow E, Merchant TE, Pappo A, Jenkins JJ, Shah AB, Kun LE. Cutaneous and subcutaneous Ewing's sarcoma: an indolent disease. *Int J Radiat Oncol Biol Phys* 2000; **46**(2): 433–438.

H-M Chen¹, W-Y Chuang^{2,3}, H-Y Chu¹ and H-Y Tan^{1,3}

¹Department of Ophthalmology, Chang Gung Memorial Hospital, Linkou, Taiwan

²Department of Pathology, Chang Gung Memorial Hospital, Linkou, Taiwan

³Chang Gung University College of Medicine, TaoYuan, Taiwan
E-mail: tanhsin@gmail.com

Eye (2014) **28**, 1143–1144; doi:10.1038/eye.2014.120;
published online 30 May 2014

Sir,
Re: Spontaneously resolved exudative retinal detachment caused by orbital cellulitis in an immune compromised adult

We thank the authors for presenting a very interesting case of orbital cellulitis and exudative serous retinal detachment with excellent MRI scans showing orbital pathology.¹ The authors suggest orbital cellulitis as a possible cause of the retinal detachment. The hematological tests suggest that the patient may have had a septicemia, with elevated ESR, C-reactive protein level, and a neutrophilic leukocytosis. Septicemia is well known to cause serous retinal detachments.

We wonder if the serous retinal detachment described in this case report was caused by the septicemia resulting from the primary infection, and if blood cultures were done to exclude this possibility.

Conflict of interest

The authors declare no conflict of interest.

Reference

- 1 Farah E, Kalantzis G, Papaefthimiou I, Koutsandrea C, Georgalas I. Spontaneously resolved exudative retinal detachment caused by orbital cellulitis in an immunocompromised adult. *Eye* 2014; **28**: 109–110.

S Perera and H Ali

Department of Ophthalmology, East Surrey Hospital, Redhill, UK
E-mail: Pereras@eyevision.fsnet.co.uk

Eye (2014) **28**, 1144; doi:10.1038/eye.2014.116; published online 30 May 2014

Sir,
Response to Perera and Ali

We would like to thank Perera and Ali¹ for their interest in our case report and their useful comments. It is true that serous retinal detachments can occur in patients suffering from septicemia or disseminated intravascular coagulation (DIC).^{2,3}

In our case,⁴ an 89-year-old immunocompromised patient presented with fever and elevated inflammatory markers (that is, raised ESR, C-reactive protein and neutrophilic leukocytosis), but the blood cultures that were performed were negative for any microorganism. Additionally, culture of nasal aspirates revealed methicillin-resistant *Staphylococcus aureus* and *Warneri-Staphylococcus*, which were consistent with ethmoid sinusitis leading to orbital cellulitis and subsequent exudative retinal detachment.

Conflict of interest

The authors declare no conflict of interest.

References

- 1 Perera S, Ali H. Re: Spontaneously resolved exudative retinal detachment caused by orbital cellulitis in an immune compromised adult. *Eye* 2014; **28**: 1144.
- 2 Lewis K, Herbert EN, Williamson TH. Severe ocular involvement in disseminated intravascular coagulation complicating meningococcaemia. *Graefes Arch Clin Exp Ophthalmol* 2005; **243**: 1069–1070.
- 3 Hoines J, Buettner H. Ocular complications of disseminated intravascular coagulation (DIC) in abruptio placentae. *Retina* 1989; **9**: 105–109.
- 4 Farah E, Kalantzis G, Papaefthimiou I, Koutsandrea C, Georgalas I. Spontaneously resolved exudative retinal detachment caused by orbital cellulitis in an immunocompromised adult. *Eye* 2014; **28**: 109–110.

I Georgalas, G Kalantzis, E Farah, I Papaefthimiou and C Koutsandrea

Department of Ophthalmology, 'G.Gennimatas' Hospital of Athens, University of Athens, Athens, Greece
E-mail: igeorgalas@yahoo.com

Eye (2014) **28**, 1144; doi:10.1038/eye.2014.117; published online 30 May 2014

Sir,
Maculopathy following extended usage of Clomiphene citrate

Clomiphene citrate (CC) is a selective estrogen receptor modulator mostly used for treatment of infertility associated with polycystic ovarian disease.¹ Unlike Tamoxifen, which is also a selective estrogen receptor