**Sir,
Intra-familial phenotype variability in patients with
Jalili syndrome**

Jalili syndrome¹ is a rare autosomal-recessive disorder, which is caused by mutations in the *CNNM4* gene.^{2,3} Two phenotypes are proposed: associated

with bull's eye maculopathy and peripheral retinal degeneration (type A), or with minor retinal dystrophy (type B).⁴

We report two new cases (siblings of 15 and 16 years of age) with different retinal findings, and their relationship with the existing phenotype classification. Fundus examination revealed severe bull's eye maculopathy in one sibling and diffuse retinal dystrophy in the other.

Figure 1 (a–e) Ocular and dental phenotype: ocular phenotype is illustrated by color fundus images and corresponding autofluorescence images, which show mild (III/4) to severe (III/5) maculopathy and retinopathy. Kinetic visual fields show a central scotoma and retained peripheral field sensitivity to large isopters (blue: V:4e; green: I:4e) in both patients, with the exception of the left eye of III/4 (lower panel). Horizontal line scans through the center of the macular are shown at the right. (a) Detailed retinal OCT analysis of the horizontal scans in this area of increased AF shows loss of the inner segment ellipsoid (illustrated by the absence of blue lines) and severe reduction of the outer retinal layers with merging of the outer limiting membrane (OLM) and the retinal pigment epithelium (RPE)/Bruch's membrane complex (absence of red lines). (b) Three stages of retinal layer changes are demonstrated on one horizontal line scan (yellow line, inset): absent inner segment ellipsoid (blue arrow), enhanced visibility of the inner plexiform layer (green arrow), and merging of the OLM and the RPE (red arrows). (c) Full-field ERG waveforms of the two patients and one age-matched control show severely reduced cone and rod function. (d) Color dental images and radiography show typical signs of AI (e).

Figure 1 (Continued)

Table 1 Review of published *CNNM4* mutations and associated phenotype

Mutation 1		Mutation 2	Author	N	M	F	Age (years)	VA	Nystagmus (Dpt)	Refraction (SE)	Fundus	ERG ^a	Origin
c.1312dupC; p.L438P/s*9	c.1312dupC; p.L438P/s*9	Present study		2	0	2	15, 16	20/200, 20/400 (1/2 progressive)	Fine pendular (1/2)	−0.5 to +2.0	Optic atrophy Macular atrophy of minor to severe extent	Scotopic: reduced, delayed Photopic: NR	Kosovo
c.1312dupC; p.L438P/s*9	c.1312dupC; p.L438P/s*9	Polok <i>et al</i> ³		2	1	1	7, 14	20/100 to 20/320	Pendular	Highly hypermetropic	Optic atrophy Macular atrophy w/ pigment mottling, periphery: white dots, 1/2: bone spiculae	Scotopic: b-wave reduced, slightly delayed Photopic: NR Repeat after 7 yrs: progressive deterioration	Kosovo
c.1312dupC; p.L438P/s*9	c.1312dupC; p.L438P/s*9	Parry <i>et al</i> ² Michaelides <i>et al</i> ⁶		2	2	0	8, 10	3/60	Fine pendular	Hypermetropic astigmatism	Macular atrophy and pigmentation	Rod response: abnormal Cone response: NR Repeat after 4 yrs: progressive deterioration	Kosovo
c.1312dupC; p.L438P/s*9	c.1312dupC; p.L438P/s*9	Zobor <i>et al</i> ⁷		1	0	1	9	0.05, 0.125		Slight myopia, astigmatism	Optic atrophy RPE atrophy macula	Scotopic: slightly delayed Cone response: NR	Kosovo
c.1312dupC; p.L438P/s*9	c.1312dupC; p.L438P/s*9	Luder <i>et al</i> ⁵		2	2	2	3, 4	20/200	Fine pendular to jerky	+8.0, +9.0	Optic atrophy Macular atrophy	Scotopic: reduced Photopic: NR Cone response: NR	Kosovo
c.599C>A; p.S200Y	c.599C>A; p.Ser200Y	Parry <i>et al</i> ² Jalili and Smith ¹ Jalili ⁴		31	17	14	0.25–50	6/36 to NLP	Fine pendular to jerky	average +3.0	RPE macular atrophy, normal optic disc (early) Chorioretinal atrophy, optic atrophy (late)	Scotopic (<i>n</i> = 3): slightly to severely reduced b-wave Flicker response: NR b-wave impaired, NR at age 10, cone: impaired to NR	Gaza A
c.1813C>T; p.R605*	c.1813C>T; p.R605*	Parry <i>et al</i> ² Jalili ⁴		3	1	2	5, 6, 10	2/60 to 6/60	Fine	+2.0 to 4.0	Normal macula, few RPE changes (1/3), optic disc unremarkable		Gaza B
c.586T>C; p.S196P	c.586T>C; p.S196P	Parry <i>et al</i> ²		2 ^b	2	2	5, 6						Turkey
c.1-?_1403+?del	c.1-?_1403+?del	Parry <i>et al</i> ²		4 ^b	3	1							Iran
c.2149C>T; p.Q717*	c.62_145 del: L21H/s*185	Parry <i>et al</i> ²		5 ^b	5								Guatemala
c.971T>C; p.L324P	c.971T>C; p.Q564*	Parry <i>et al</i> ²		1 ^b	1								Scotland
c.707G>A; p.R236Q	c.707G>A; p.R236Q	Polok <i>et al</i> ³		3	2	1	2, 6, 12	Low	Rapid			Scotopic: normal to reduced, photopic: severely attenuated to NR NR	Lebanon
		Polok <i>et al</i> ³		1	0	1	38					NR	

Table (Continued)

Mutation 1	Mutation 2	Author	N	M	F	Age (years)	VA	Nystagmus	Refraction (SE) (Dpt)	Fundus	ERG ^a	Origin
c.971T>C; p.L324P	c.971T>C; p.L324P	Doucette <i>et al</i> ⁸	4	1	3	16–28 (f.u.)	LP (10/200 at age 6) 20/200 to HM, progression	Yes	Slight myopia, astigmatism Myopia	Optic atrophy, macular atrophy, bone spiculae in midperiphery 1/4 Maculopathy, 1/4 bone spiculae, 2/4 ND	1/4 Rod normal, cone absent, 1/4 rod borderline, cone absent, 2/4 ND CRD (not sure if ERG was done)	Northern Europe
c.1484C>T p.T495I	c.1484C>T p.T495I	Abu-Safieh <i>et al</i> ⁹	1									
c.189del p.D63E/s*12	c.189del p.D63E/s*12	Coppieters <i>et al</i> ¹⁰	3	2	1					1/3 Reported: maculopathy, outer retinal atrophy w/ pigmentation		Algeria

Abbreviations: CRD, cone-rod dystrophy; Dpt, diopter; ERG, full-field electroretinography; F, female; f.u., follow-up; HM, hand motion detection; LP, light perception; M, male; ND, not done; NLP, no light perception; NR, non-recordable; RPE, retinal pigment epithelium; SE, spherical equivalent; VA, visual acuity; w/, with; yrs, years.

^aERG description as written in publication.

^bNumbers are based on the pedigree shown in publication.

Figure 2 Genetic testing results: Sanger DNA sequencing revealed a homozygous mutation in the two affected siblings (III:4 and III:5) as shown in the forward and reverse sequencing profiles from the DNA of the two patients. The mutation elongates the Oligo-C stretch in exon 1 of the *CNNM4* gene, which consists of six cytosine residues in the reference sequence, by 1 nucleotide (c.1312dup; p.Leu438Profs*9). The DNA from the mother (II:4) is heterozygous for this mutation.

ERG showed cone-rod dysfunction. OCT demonstrated thinning of the outer retinal layers, in particular the outer nuclear layer and the outer photoreceptor segments. Both patients showed amelogenesis imperfecta (Figure 1). Phenotype details in comparison with the reported patients with *CNNM4* mutations are listed in Table 1. The same homozygous mutation c.1312dup; p.Leu438Profs*9 was found in the affected patients (III:4, III:5). The mother (II:4) was heterozygous for this sequence alteration in *CNNM4* (Figure 2). The mutation within the cystathionine beta-synthase domain most likely results in a premature termination codon and nonsense-mediated mRNA decay of the mutant transcript. No mutation in the *ABCA4* gene was identified.

A similar dental phenotype with the characteristics of AI is described in all publications of patients with *CNNM4* mutations. In his phenotype dissection, Jalili described anterior open bite (AOB) in 2/30 and posterior open bite in 1/30 'type A' patients, whereas AOB was present in all three patients of the 'type B' phenotype. No open bite abnormality was seen on examination in our two patients and in one of the two patients reported by Luder *et al.*⁵

The intra-familial variability presented here is not consistent with a strict phenotype-genotype correlation, and may also argue against a rigid phenotype differentiation.⁴ As the patients with 'type B' phenotype were examined at a younger age than the patient with 'type A', it is possible that those patients with 'type B' may have shown minimal macular signs as most of them had a visual impairment and signs of a cone-rod dystrophy.

The proposed strict differentiation between type A and B may not be applicable to all affected patients and families.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

This study was funded in part by a grant from the Swiss National Science Foundation (grant number: 31003A 122359 to WB).

References

- 1 Jalili IK, Smith NJ. A progressive cone-rod dystrophy and amelogenesis imperfecta: a new syndrome. *J Med Genet* 1988; **25**: 738–740.
- 2 Parry DA, Mighell AJ, El-Sayed W, Shore RC, Jalili IK, Dollfus H *et al.* Mutations in *CNNM4* cause Jalili syndrome, consisting of autosomal-recessive cone-rod dystrophy and amelogenesis imperfecta. *Am J Hum Genet* 2009; **84**: 266–273.
- 3 Polok B, Escher P, Ambresin A, Chouery E, Bolay S, Meunier I *et al.* Mutations in *CNNM4* cause recessive cone-rod dystrophy with amelogenesis imperfecta. *Am J Hum Genet* 2009; **84**: 259–265.
- 4 Jalili IK. Cone-rod dystrophy and amelogenesis imperfecta (Jalili syndrome): phenotypes and environs. *Eye (Lond)* 2010; **24**: 1659–1668.
- 5 Luder HU, Gerth-Kahlert C, Ostertag-Benzinger S, Schorderet DF. Dental phenotype in Jalili syndrome due to a c.1312 dupC homozygous mutation in the *CNNM4* gene. *PLoS One* 2013; **8**: e78529.
- 6 Michaelides M, Bloch-Zupan A, Holder GE, Hunt DM, Moore AT. An autosomal recessive cone-rod dystrophy associated with amelogenesis imperfecta. *J Med Genet* 2004; **41**(6): 468–473.
- 7 Zobor D, Kaufmann DH, Weckerle P, Sauer A, Wissinger B, Wilhelm H *et al.* Cone-rod dystrophy associated with amelogenesis imperfecta in a child with neurofibromatosis type 1. *Ophthalmic Genet* 2012; **33**(1): 34–38.
- 8 Doucette L, Green J, Black C, Schwartzentruber J, Johnson GJ, Galutira D *et al.* Molecular Genetics of Achromatopsia in Newfoundland Reveal Genetic Heterogeneity, Founder Effects and the First Cases of Jalili Syndrome in North America. *Ophthalmic Genet* 2013; **34**(3): 119–129.
- 9 Abu-Safieh L, Alrashed M, Anazi S, Alkuraya H, Khan AO, Al-Owain M *et al.* Autozygome-guided exome sequencing in retinal dystrophy patients reveals pathogenetic mutations and novel candidate disease genes. *Genome Res* 2013; **23**(2): 236–247.
- 10 Coppieters F, Van Schil K, Bauwens M, Verdin H, De Jaegher A, Syx D *et al.* Identity-by-descent-guided mutation analysis and exome sequencing in consanguineous families reveals unusual clinical and molecular findings in retinal dystrophy. *Genet Med* 2014; **16**(9): 671–680.

C Gerth-Kahlert¹, B Seebauer^{2,3}, S Dold^{2,3}, JVM Hanson¹, H Wildberger¹, A Spörri⁴, H van Waes⁴ and W Berger^{2,3,5}

¹Department of Ophthalmology, University Hospital Zurich, Zurich, Switzerland

²Institute of Medical Molecular Genetics, Zurich, Switzerland

³Neuroscience Center Zurich, University and ETH Zurich, Zurich, Switzerland

⁴Clinic of Orthodontics and Paediatric Dentistry, Center of Dental Medicine, Zurich, Switzerland

⁵Zurich Center for Integrative Human Physiology, University of Zurich, Zurich, Switzerland
E-mail: christina.gerth-kahlert@usz.ch
Data were presented partially at the 2013 ARVO Annual Meeting.

Eye (2015) **29**, 712–716; doi:10.1038/eye.2014.314; published online 23 January 2015

Sir, Rupture of Descemet's membrane secondary to presumed non-accidental injury

We present the first case of multiple uniocular breaks in Descemet's membrane secondary to presumed non-accidental injury (NAI).

Case report

The first of twins, born at 35 weeks by spontaneous unassisted vaginal delivery, presented with a 2-week history of unexplained corneal haze in her left eye at the age of 4 months. Topical antibiotics and steroids were