

Determinants of visual acuity outcomes in eyes with neovascular AMD treated with anti-VEGF agents: an instrumental variable analysis of the AURA study

FG Holz, R Tadayoni, S Beatty, AR Berger, MG Cereda, P Hykin, G Staurenghi, K Witttrup-Jensen, J Nilsson, K Kim and S Sivaprasad

Eye (2017) 31, 166; doi:10.1038/eye.2016.276

Correction to: *Eye* (2016) 30, 1063–1071; doi:10.1038/eye.2016.90; published online 20 May 2016

Since the publication of this article, the authors would like to correct a mistake in the heading of Table 2. The lower 95% confidence interval (CI) and upper 95% CI should be replaced with the

Year 2 coefficient and *P*-value, respectively. Accordingly, the first column should be replaced with Year 1 coefficient and *P*-value (as shown). The coefficients for Year 1 are quoted in the text, but this does not change the interpretation of the findings.

The authors would like to apologise for this error.

Table 2 Correlation between number of ranibizumab injections and candidate variables (all countries; over 2 years)

Parameters	Year 1		Year 2	
	Coefficient ^a	<i>P</i> -value	Coefficient ^a	<i>P</i> -value
Age at start of therapy	-0.0289	0.1141	-0.0378	0.056
Sex	0.0171	0.384	-0.0126	0.5193
Health insurance (public)	-0.028	0.1544	-0.053	0.0071
Reimbursement type (national standards)	0.1297	< 0.001	0.1524	< 0.001
Medical history (prior disease)	-0.0268	0.1711	-0.0354	0.0703
Medical history (concomitant disease)	-0.0003	0.9866	0.018	0.3595
Medical history (ocular disease)	-0.0246	0.2083	-0.0331	0.0905
Baseline VAS	-0.0088	0.6758	0.0424	0.0444
Treatment duration	0.5645	< 0.001	0.7389	< 0.001
Switch to other treatments	0.0275	0.1599	0.0632	0.0012
Baseline presence of retinal breaks ^b	NA	NA	NA	NA
Baseline presence of pigment epithelial detachment	0.0821	0.0038	0.1163	< 0.001
Number of ophthalmoscopies	0.2499	< 0.001	0.3425	< 0.001
Optical coherence tomography, <i>n</i>	0.3777	< 0.001	0.4955	< 0.001
Fluorescein angiography, <i>n</i>	0.0509	0.0093	0.1154	< 0.001
ICGA, <i>n</i>	-0.0341	0.0814	0.0302	0.1228
Ophthalmoscopies/optical coherence tomography, <i>n</i>	0.3608	< 0.001	0.4673	< 0.001
Number of monitoring visits	-0.1349	< 0.001	0.0239	0.2224

Abbreviations: ICGA, indocyanine green angiography; NA, not available; VAS, visual acuity score. ^aPearson's correlation coefficient at 0.05 significance level. ^bNo observation with retinal break episode. The bold values highlights the candidate instrumental variables.