Continuing Medical Education:

Non-contact ultra-widefield retinal imaging of infants with suspected abusive head trauma

IH Yusuf, JK Barnes, THM Fung, JS Elston
and CK Patel

Release date: 24 February 2017; Expiration date: 24 February 2018

This activity has been planned and implemented through the joint providership of Medscape, LLC and Springer Nature. Medscape, LLC is accredited by the American Nurses Credentialing Center (ANCC), the Accreditation Council for Pharmacy Education (ACPE), and the Accreditation Council for Continuing Medical Education (ACCME), to provide continuing education for the healthcare team.

Medscape, LLC designates this Journal-based CME activity for a maximum of *1.00 AMA PRA Category 1 Credit(s)*[™]. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

All other clinicians completing this activity will be issued a certificate of participation. To participate in this journal CME activity: (1) review the learning objectives and author disclosures; (2) study the education content; (3) take the post-test with a 75% minimum passing score and complete the evaluation at www.medscape.org/journal/eye; (4) view/print certificate.

Learning Objectives

Upon completion of this activity, participants will be able to:

1. Interpret findings of non-contact ultra-widefield retinal imaging using the Optos P200MA scanning laser ophthalmoscope among infants being evaluated for abusive head trauma (AHT), based on a retrospective, observational case series
2. Identify advantages of the Optos P200MA scanning laser ophthalmoscope *vs* RetCam for the documentation of retinal haemorrhages in the context of AHT
3. Identify advantages of RetCam *vs* the Optos P200MA scanning laser ophthalmoscope for the documentation of retinal haemorrhages in the context of AHT

Authors/Editors disclosure information

Andrew J Lotery has disclosed the following relevant financial relationships: Served as an advisor or consultant for: Bayer HealthCare Pharmaceuticals; Roche. Served as a speaker or a member of a speakers bureau for: Bayer HealthCare Pharmaceuticals.

Imran Yusuf has disclosed no relevant financial relationships.

Joanna Kate Barnes has disclosed no relevant financial relationships.

Timothy HM Fung has disclosed no relevant financial relationships.

John S Elston has disclosed no relevant financial relationships.

Chetan K Patel has disclosed the following relevant financial relationships: Served as a speaker or a member of a speakers bureau for: Optos PLC.

Journal CME author disclosure information.

Laurie Barclay has disclosed the following relevant financial relationships: Owns stock, stock options, or bonds from: Pfizer.

Non-contact ultra-widefield retinal imaging of infants with suspected abusive head trauma

IH Yusuf, JK Barnes, THM Fung, JS Elston
and CK Patel

Abstract

Purpose The purpose of the study was to audit the use of non-contact ultra-widefield retinal imaging in infants with suspected abusive head trauma (AHT) using the Optos P200MA Scanning Laser Ophthalmoscope.

Patients and methods A retrospective, observational case series. Ten eyes of five consecutive infants (aged 1–15 months) with suspected (or in 1 case, known) AHT referred for an ophthalmological opinion were included. Each infant underwent non-contact ultra-widefield retinal imaging using the Optos P200MA scanning laser ophthalmoscope. Optos fundus fluorescein angiography (FFA) was performed in one infant with oral sedation. The other four infants did not require sedation. The main outcome measure was the acquisition of a single, definitive ultra-widefield retinal image in each eye. Safety was audited by determining adverse changes in heart rate and oxygen saturations that required cessation of imaging.

Results The Optos P200MA ultra-widefield scanning laser ophthalmoscope acquired good quality retinal images in all infants. Documentation of acute, widespread retinal haemorrhages contributed to a diagnosis of AHT in three infants. Chronic pre-macular haemorrhage and macular schisis were documented by FFA in a fourth infant. The absence of retinal haemorrhages was documented in a fifth infant contributing to the exclusion of a diagnosis of AHT. There were no adverse safety signals in any infant in this series.

Conclusion The Optos P200MA ultra-widefield scanning laser ophthalmoscope appears safe to use in infants with suspected AHT, providing high-quality retinal images in a single frame without ocular contact.

Optos P200MA may be used as alternative to RetCam to document retinal haemorrhages in stable infants with suspected AHT.

Eye (2017) **31**, 353–363; doi:10.1038/eye.2017.2; published online 24 February 2017

Introduction

Abusive head trauma (AHT)—also known as shaken baby syndrome—is the commonest cause of traumatic death in infancy.¹ It is associated with a 30% mortality, with a further 30% of infants suffering severe neurological sequelae.² AHT typically presents with an acute unexplained encephalopathy, intracranial haemorrhage (characteristically subdural), and retinal haemorrhages. Exclusion, by full investigation, of congenital or acquired coagulopathies and other masquerading organic disorders² is mandatory. The diagnosis of AHT is imperative, to ensure safeguarding of the infant, protection of any siblings and initiation of social services and police investigations.

The detection of retinal haemorrhages is central to the diagnosis of AHT.³ The appearance, distribution, laterality, and severity of haemorrhages will assist in objectively distinguishing between accidental and non-accidental causes of head injury in infancy.¹

Because of the child protection measures and legal proceedings initiated by a diagnosis of AHT, high-quality photo-documentation of the retinal appearance in cases of suspected AHT is critical. The diagnosis of AHT is likely to be challenged in Court and high-quality retinal images provide objective evidence of any abnormalities. Because some superficial retinal haemorrhages in AHT may resolve over a period of days, accurate documentation is required as soon as possible after admission. Photographs also enable the retinal complications of AHT such as pre-macular haemorrhage and macular

The Oxford Eye Hospital,
West Wing, John Radcliffe
Hospital, Oxford, UK

Correspondence:
CK Patel, The Oxford Eye
Hospital, John Radcliffe
Hospital, Headley Way,
Oxford OX3 9DU, UK
Tel/Fax: +44 (0)18 6523
4736.
E-mail: ckpatel@
btinternet.com

Received: 20 August 2016
Accepted in revised form:
16 November 2016
Published online:
24 February 2017

schisis to be documented to determine the necessity and timing of any potential intervention.

RetCam (Pleasanton, CA, USA) has become established for retinal imaging in infants with suspected AHT.⁴ The RetCam is a portable device that requires coupling fluid and ocular contact to acquire retinal images on a supine infant. Sedation may be required for RetCam imaging in some cases, especially if retinal angiography is required.⁵ Series evaluating the utility of the RetCam imaging system in AHT are often undertaken on paediatric intensive care units where infants are sedated or ventilated,⁴ although many series do not explicitly state whether sedation was used on RetCam imaging in this patient group.^{6,7} The RetCam is able to image a 130° field, typically requiring 10–12 montaged retinal images to record the combined appearance of both eyes. RetCam imaging is the current 'gold standard' method of documentation.

When our Retcam was not available in the ophthalmology outpatient facility, we started using the Optos P200MA Scanning Laser Ophthalmoscope (Optos, Dunfermline, UK), which has been demonstrated to attain high-quality retinal images in infants with retinopathy of prematurity (ROP),^{8,9} incontinentia pigmenti,¹⁰ and other paediatric retinal disorders¹¹ without the need for ocular contact. Optos captures a 200° retinal field enabling documentation of the retinal periphery in a single image, typically without the need for sedation.

We present a series of four infants presenting with suspected AHT, in whom ultra-widefield retinal photographs were acquired using the Optos P200MA Scanning Laser Ophthalmoscope. We describe the Optos imaging technique in infants, present ultra-widefield images, and illustrate their impact on patient management. We outline the relative advantages of Optos and RetCam in the context of retinal imaging of infants with suspected AHT. We demonstrate the utility of fundus fluorescein angiography (FFA) in the evaluation of retinal pathology in a fifth infant who had sustained AHT some weeks earlier. This audit was undertaken to evaluate the safety of this imaging modality in infants with suspected AHT.

Materials and methods

Patients

Four consecutive infants with suspected AHT were referred to the paediatric ophthalmology service at the Oxford Eye Hospital, UK between November 2013 and December 2014 for retinal examination and photography. Another infant, diagnosed with AHT elsewhere was referred for a surgical opinion. Infants were aged between 1 and 15 months, four were male and one female. The

audit protocol was approved by the clinical governance committee of the Oxford Eye Hospital (Registration number: 210416-NOTSS-YUSUF). All imaging was performed with the full informed consent of the parents, and in adherence with the tenets of the Declaration of Helsinki.

Retinal imaging protocol

Retinal photographs were acquired using the Optos P200MA Scanning Laser Ophthalmoscope in each infant as previously described.^{8–11} Mydriasis was achieved with topical phenylephrine 2.5% and cyclopentolate 0.5%. Topical proxymetacaine 0.5% was instilled into each eye followed by insertion of a paediatric Lieberman speculum to part the eyelids, secured to the temple with Micropore tape. A peripheral oxygen saturation probe monitored heart rate and peripheral capillary oxygen saturation with a paediatric neonatal nurse present, trained in neonatal resuscitation. The infant was held in the 'flying baby position' with the infants trunk supported by the forearm, chin supported by the hand, with the contralateral hand placed on the occiput for stability. A physiological saline solution was applied intermittently to the ocular surface to maintain a clear cornea. The infant was manoeuvred to the camera's aperture and fine adjustments were made by the imaging technician using an image capture control. Pseudo-colour fundal images were acquired from both eyes, taking <5 min in each case. Ultra-widefield fluorescein angiography was performed in one infant following peripheral venous cannulation. Ten percent sodium fluorescein was injected at a dose of 0.1 mg/kg, with images acquired from both eyes at regular time intervals with early phase images acquired on the right eye and late phases in both eyes. No infants required cessation of image acquisition due to reduced oxygen saturation or altered cardiovascular parameters. Optos fundus images are presented in pseudo colour, balanced from green and red laser images.

Results

Case 1

A 2-month-old male infant had been admitted to the paediatric high-dependency unit with unexplained facial soft tissue injuries, apnoea, and a nuchal rash. General examination revealed hypertonia with brisk reflexes. Neuroimaging revealed bilateral acute subdural haematomas with subarachnoid blood. Dilated retinal examination demonstrated a symmetrical distribution of over 100 retinal haemorrhages in each eye concentrated at the posterior pole, extending to the retinal periphery. The optic discs appeared unremarkable. Ultra-widefield

Figure 1 Ultra-widefield optomap colour images of right (a) and left eyes (b) demonstrating widespread, bilateral, multi-layered retinal haemorrhages affecting both the posterior pole and retinal periphery, with pre-macular haemorrhages (white arrowheads) identified on the right.

colour images were acquired without sedation (Figure 1). A diagnosis of AHT was made. Three weeks later, the eyes were convergent with reduced abduction and visual responsiveness. Dilated funduscopy revealed resolution of retinal haemorrhages.

Case 2

A 4-week-old male infant presented with a left parieto-occipital fluctuant swelling measuring 4×5.5 cm without discoloration. The child was born at 39 weeks gestation by caesarean section following abandoned forceps delivery. Scalp electrodes were used but vacuum-assisted delivery was not attempted. No clear explanation for the swelling was apparent, although the child had been vomiting. Computerised tomography (CT) imaging of the brain revealed a shallow left parieto-occipital sub-galeal fluid collection but no intracranial abnormalities. A paediatric ophthalmology consult was requested. Ultra-widefield colour images were acquired without sedation, documenting the absence of retinal haemorrhages to the retinal periphery (Figure 2). AHT was subsequently excluded; radiological skeletal survey and extended clotting screen were unremarkable and the sub-galeal collection resolved on repeat neuroimaging.

Case 3

A 4-month-old male infant had been admitted to the paediatric intensive care unit with unexplained loss of consciousness that required mechanical ventilation. The infant was previously fit and well, born through

normal vaginal delivery at 40 weeks gestation. Neuroimaging showed intracranial, subarachnoid, and subdural haemorrhages. Dilated retinal examination revealed bilateral pre-macular haemorrhages and extensive superficial, intra-retinal and pre-retinal haemorrhages in the posterior pole, extending to the periphery. The optic discs were normal. A diagnosis of AHT was made. RetCam images documented the retinal appearance in both eyes (Supplementary information 1a and 1b). Ultra-widefield retinal imaging without sedation 6 days later documented resolution of the majority of the peripheral and posterior pole retinal haemorrhages, with improvement of bilateral subretinal, pre-macular haemorrhages in both eyes (Figure 3). Three weeks later, fundal appearances had normalised.

Case 4

A 15-month-old male was admitted to the paediatric intensive care unit in poor condition after a seizure and collapse attributed to a fall from a chair. A CT scan performed on admission revealed a 6 mm acute subdural haematoma in the right fronto-parietal region with midline shift, although neurosurgical intervention was not required. A skeletal survey demonstrated a wrist fracture. Blood coagulation disorders were excluded on testing. On examination, multiple bruises were present on the forehead, chin, and cheek. Dilated fundal examination (using a speculum and indirect ophthalmoscope without scleral indentation) the day after admission in an awake, active child was difficult, with suspected retinal haemorrhages inferior to the left optic disc only. However, ultra-widefield retinal imaging 6 days later in

Figure 2 Ultra-widefield optomap colour images of right (a) and left eyes (b) documenting the absence of retinal haemorrhages in the posterior pole and retinal periphery of both right and left eyes. Optic disc appearance is unremarkable.

Figure 3 Ultra-widefield optomap colour images of right fundus (a) demonstrating haemorrhage at the posterior pole with widespread retinal haemorrhages extending into the retinal periphery. Macula exudative detachment (white arrowhead) is noted adjacent to the superotemporal arcade secondary to haemorrhage. The left (b) fundus demonstrates resolving haemorrhage at the posterior pole (white arrowhead) and resolving peripapillary intra-retinal haemorrhages extending into the retinal periphery, distributed adjacent to the retinal veins.

the outpatient clinic revealed multiple intra-retinal and pre-retinal haemorrhages in the left eye, and sparse retinal haemorrhages inferiorly in the right eye that were not identified clinically (Figure 4). The retinal appearances were consistent with inflicted trauma and a diagnosis of AHT was made.

Case 5

A 6-month-old female infant was referred to the paediatric vitreo-retinal service for surgical drainage of presumed pre-macular haemorrhage obscuring the fovea in both eyes. Two months earlier, the infant had presented

with an acute encephalopathy and been found to have bilateral subdural haematomas and extensive retinal haemorrhages including at the posterior poles and obscuring the macular. A diagnosis of AHT was made. Neurological recovery was swift, but visual function remained poor and acuities were fix and follow OD, and no interest OS. Ultra-widefield colour images and intravenous FFA (0.1 mg/kg) were undertaken under oral sedation (chloral hydrate 50 mg/kg). A right posterior pole haemorrhage was demonstrated with a fluid level and the fovea was not obscured (Figure 5a). FFA confirmed normal macular perfusion and fovea was not obscured. Retinal vessels anterior to the haemorrhage

Figure 4 Ultra-widefield optomap colour images of the right fundus (a) demonstrating several retinal haemorrhages inferiorly in the retinal periphery, which were not detected on clinical examination, and the left fundus (b) identifying multiple haemorrhages extending into the retinal periphery.

(Figure 5c), prove at least part of the collection was intra-retinal rather than in the retrohyaloid space. However, the inferotemporal arcade is masked, indicating that the haemorrhage is likely to be multi-layered. No vitreo-retinal intervention was therefore recommended. The left eye revealed macular schisis with posterior pole retinal detachment, partial posterior vitreous detachment with persistent retrohyaloid blood (Figures 5b and d).

Discussion

Non-contact retinal imaging in infants using the Optos P200MA scanning laser ophthalmoscope is capable of acquiring clinically useful, high-quality, ultra-widefield images in the following settings: (1) colour images of premature infants with ROP, demonstrating disease progression and regression following treatment⁹ (2) FFA images using oral fluorescein in infants with ROP to guide retinal photocoagulation treatment⁸ (3) FFA images using oral fluorescein in an infant with incontinentia pigmenti to guide surgical planning¹⁰ (4) Intravenous FFA in infants with a range of retinal vascular disorders including familial exudative vitreoretinopathy.¹¹

This series is the first to describe non-contact ultra-widefield retinal imaging of infants with suspected AHT. The Optos P200MA scanning laser ophthalmoscope effectively documented acute and chronic AHT-related retinal haemorrhages, accurately defined the anatomical location of haemorrhage in one infant who was referred for consideration of vitrectomy, and excluded retinal haemorrhages and consequently AHT in a fifth infant. A single, high-yield, definitive retinal image was attained in

each case to confirm the diagnosis and document clinical findings. In case 4, an experienced paediatric ophthalmologist had achieved a very limited retinal view, but Optos imaging documented much more widespread abnormalities than were suspected. There were no adverse changes in physiological parameters in any infant that necessitated cessation of imaging.

RetCam is a well-established retinal imaging modality for infants, both for colour photographs and FFA in the context of ROP^{8,9} and AHT.¹ Non-contact ultra-widefield retinal imaging using the Optos P200MA scanning laser ophthalmoscope offers many advantages over RetCam for the documentation of retinal haemorrhages in the context of AHT:

Non-contact imaging

Infants with AHT may suffer significant intracranial haemorrhage and are at increased risk of bradycardia and respiratory arrest from secondary brainstem dysfunction. The Optos P200MA imaging system does not require ocular contact, and may consequently reduce the risk of stimulating oculocardiac/vagal responses in infants with AHT—a risk with RetCam contact imaging.¹² However, the speculum that is necessary to undertake imaging is likely to cause the infant some distress.¹³ Ocular steering required on RetCam imaging to attain peripheral retinal images entails significant additional pressure on the globe, particularly without sedation when the infant's Bell's reflex may be preserved. The Optos P200MA may reduce the risk of inducing intra-retinal haemorrhages—which may be morphologically identical to those encountered in AHT—a

Figure 5 (a) Ultra-widefield optomap colour image of the right eye demonstrates the appearance of a pre-retinal haemorrhage at the posterior pole with a fluid level, partially obscuring the macula (white arrowhead). A resolving intra-retinal haemorrhage was noted adjacent to the optic disc (black arrowhead). The retinal periphery appears unaffected. (b) The left eye reveals macular schisis and an annulus of retrohyaloid blood (white arrowheads) at the limits of vitreo-retinal attachment. A fluid level of retrohyaloid blood is seen inferiorly, and the optic disc and fovea are obscured. (c) FFA confirms anterior displacement of retinal vessels over the posterior pole haemorrhage (white arrowhead) indicating that part of the haemorrhage is intra-retinal or subretinal, although the inferotemporal arcade is obscured, indicating that the haemorrhage is likely multi-layered. This information changed the clinical course: pars plana vitrectomy was not required. FFA further confirms normal macular perfusion. (d) FFA confirms the extent of pre-retinal haemorrhage and macular schisis. Perfusion of retinal arterioles is preserved. Fine perfused vessels (white arrowhead) are identified protruding anteriorly at the optic disc.

possibility with RetCam contact imaging, at least in the setting of ROP.¹⁴ The risk of inducing retinal haemorrhages with the RetCam in suspected AHT is theoretical and has never been reported. We have not encountered any adverse events in infants undergoing non-contact Optos P200MA imaging in our centre—across any indication—using the ‘flying baby position’ in more than 100 examinations to date.^{8–11} Larger numbers of Optos non-contact imaging studies are required before definitive conclusions can be made regarding safety.

Ocular contact required during RetCam imaging holds a theoretical risk of inducing retinal haemorrhages.¹⁴

However, in this report, the cause of the retinal haemorrhages was unclear. It must be noted that the retinal haemorrhages in this case (a premature infant) are likely to be different to the responses in a term infant with normal retinal development. The risk of cross infection between infants is also minimised as the P200MA imaging system does not contact the ocular surface.

Field of view

The Optos P200MA captures a 200° retinal field, superior to the 130° field possible with RetCam. Imaging of the

retinal periphery is important in the setting of AHT.¹ Systematic review of the distribution of retinal haemorrhages in abusive *vs* accidental head trauma identified 63% of children with AHT had peripheral retinal extension of haemorrhages (9% in non-AHT).¹ Ultra-widefield documentation of peripheral retinal haemorrhages increases the likelihood of AHT, and may increase diagnostic precision.¹ Accurate reporting of retinal haemorrhage location is paramount in AHT and a number of descriptive tools have been designed to aid documentation.³ To overcome challenges with photodocumentation of AHT in centres with no ophthalmic expertise, telemedicine-based imaging protocols are being evaluated.⁶ Ultra-widefield Optos P200MA may increase the sensitivity and specificity of retinal imaging of AHT in a telemedicine setting when compared to RetCam. However, a study directly comparing retinal imaging modalities is required in the setting of AHT before this conclusion can be made. The Optos P200MA imaging system may provide a greater field of imaging in the horizontal field (nasal and temporal) *vs* the vertical field (superior and inferior). In addition, the speculum and eyelashes may interfere with the inferior field of imaging (Figure 4).

General anaesthesia and sedation

Sedation is not required for the majority of infants imaged in the ophthalmic office setting with the Optos P200MA scanning laser ophthalmoscope owing to the speed of the imaging technique we have developed.^{8–11} Colour images of both eyes in this series took <5 min in each infant. Retinal haemorrhages are present bilaterally in 83% of infants with AHT¹; 2 images are required with Optos P200MA compared to 10–12 images with RetCam (typically 5–6 images are required for each eye to capture the entire retinal periphery which may be montaged). Sedation will be required more frequently to acquire high-quality images with RetCam than with Optos P200MA, due to a combination of ocular contact, eye steering, duration of imaging, and importance of ocular akinesia for image quality. Neurological observations are essential in infants with AHT due to intracranial haemorrhage; sedation may interfere with neurological parameters used to monitor infants, and presents risks related to the mechanism of action of the sedative drug (hypoventilation, hypotension etc.). Large infants and those over 6 months of age may require oral sedation for Optos P200MA retinal imaging.

Confocal optics

The Optos P200MA scanning laser ophthalmoscope is able to produce a uniform focus to multiple retinal

locations, simultaneously increasing clarity of documentation across the retinal field. This is not the case with RetCam; multiple images are required with manual focus of the lens system using a foot pedal, adding time to the process of image capture. AHT may be characterised by subretinal haemorrhages, retrohyaloid haemorrhage, schisis cavities, retinal or macular folds, retinal detachment, and papilloedema.¹ Confocal optics may produce clearer documentation of such AHT features, which are elevated from the retinal plane.

Speed of image acquisition and optics

Dual wavelength laser illumination of the Optos P200MA imaging system permits rapid image acquisition (0.25 s), overcoming the challenges to retinal imaging in mobile, non-sedated infants. The RetCam may suffer from image degradation in the setting of media opacity, such as vitreous haemorrhage in AHT.¹ Image acquisition time is prolonged by the need to replenish coupling media, and requirement for manual focusing.

Ultra-widefield fluorescein angiography

Ultra-widefield fluorescein angiography may increase diagnostic yield *vs* RetCam-assisted FFA as all retinal locations may be captured simultaneously at all time points in the study. Confocal FFA imaging ensures focus is maintained on all structures in the examination. This advantage changed the course of management in one of the presented infants (case 5). A fine retinal vessel overlying a pre-macular haemorrhage was identified by Optos-assisted FFA, indicating that the haematoma was located in the subretinal space. Vitrectomy to clear a presumed retrohyaloid haemorrhage was subsequently not advised and the haemorrhage cleared spontaneously. The plane of retinal haemorrhage carries distinct implications in the likelihood of AHT¹: the prevalence of multi-layered retinal haemorrhages is 77% in AHT and 73% of infants with AHT have evidence of retinal haemorrhages in all three layers (subretinal, intra-retinal, and subhyaloid).¹ We have demonstrated that confocal Optos-assisted FFA can accurately characterize the anatomical location of retinal haemorrhages in a single frame.

The RetCam has several advantages over the Optos P200MA imaging system in the setting infants with AHT:

- (1) *Portability.* Severe AHT may necessitate intubation to maintain the airway of infants secondary to impaired consciousness from intracranial bleeding. The RetCam is required to image the retina of a supine infant too unstable to transfer, or who is pharmacologically sedated or mechanically ventilated. The Optos

P200MA is not portable and is not currently capable of imaging an infant required to remain in a supine position. A dedicated P200MA imaging system on a neonatal high dependency unit—as has now become the case in our centre for the management of ROP—may permit ultra-widefield retinal imaging in unstable infants with AHT by eliminating the need for transfer to the ophthalmic office setting.

- (2) *Access to and familiarity with RetCam.* RetCam is widely used for paediatric retinal imaging. Many ophthalmic departments possess the RetCam, and paediatric ophthalmologists are familiar and comfortable with its use. Retinal imaging in infants with the Optos P200MA using the ‘flying baby position’ involves a learning curve for both ophthalmologist, imaging technician and paediatric nursing staff during the process of imaging. In our experience, imaging of 3–5 infants is typically required to become familiar and competent with the technique.

This observational series is the first to demonstrate that the Optos P200MA scanning laser ophthalmoscope can provide high-quality, ultra-widefield retinal images in infants with suspected AHT using a non-contact technique. We suggest that paediatricians and ophthalmologists should be aware that non-contact imaging without sedation in stable infants with suspected AHT may be the preferred modality to document and record any abnormalities.

Summary

What was known before

- Identification and documentation of retinal haemorrhages is critical to the diagnosis of abusive head trauma: the most common cause of traumatic death in infancy.
- The RetCam has become established for the documentation of retinal haemorrhages in infants, but requires ocular contact, pressure on the globe, and multiple, montaged images for documentation of the retinal periphery.
- Non-contact ultra-widefield imaging using the Optos P200MA has been used in infants with other paediatric retinal disorders such as retinopathy of prematurity and incontinentia pigmenti.

What this study adds

- The Optos P200MA scanning laser ophthalmoscope is able to acquire high-quality, ultra-widefield retinal images: documenting the appearance of the posterior pole and peripheral retina in a single image.
- The quality of images (both clarity and field) attained is superior to that acquired by RetCam.
- Optos P200MA scanning laser ophthalmoscope may be used to document the retinal appearance in stable infants with suspected abusive head trauma in the outpatient setting.

Conflict of interest

The authors declare no conflict of interest.

Author contributions

Design and conduct of the study: IHY, THMF, JSE, CKP. Collection, management, analysis, and interpretation of data: IHY, JKB, THMF, JSE, CKP. Preparation, review, or approval of the manuscript: IHY, JKB, THMF, JSE, CKP.

Acknowledgements

CK Patel had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

- 1 Maguire SA, Watts PO, Shaw AD, Holden S, Taylor RH, Watkins WJ *et al*. Retinal haemorrhages and related findings in abusive and non-abusive head trauma: a systematic review. *Eye* 2013; **27**(1): 28–36.
- 2 Carty H, Ratcliffe J. The shaken infant syndrome. *BMJ* 1995; **310**(6976): 344–345.
- 3 Tandon A, McIntyre S, Yu A, Stephens D, Leiby B, Croker S *et al*. Retinal haemorrhage description tool. *Br J Ophthalmol* 2011; **95**(12): 1719–1722.
- 4 Nakagawa TA, Skrinska R. Improved documentation of retinal hemorrhages using a wide-field digital ophthalmic camera in patients who experienced abusive head trauma. *Arch Pediatr Adolesc Med* 2001; **155**(10): 1149–1152.
- 5 Blair MP, Shapiro MJ, Hartnett ME. Fluorescein angiography to estimate normal peripheral retinal nonperfusion in children. *J AAPOS* 2012; **16**(3): 234–237.
- 6 Saleh M, Schoenlaub S, Desprez P, Bourcier T, Gaucher D, Astruc D *et al*. Use of digital camera imaging of eye fundus for telemedicine in children suspected of abusive head injury. *Br J Ophthalmol* 2009; **93**(4): 424–428.
- 7 Mulvihill AO, Jones P, Tandon A, Fleck BW, Minns RA. An inter-observer and intra-observer study of a classification of RetCam images of retinal haemorrhages in children. *Br J Ophthalmol* 2011; **95**(1): 99–104.
- 8 Fung TH, Muqit MM, Mordant DJ, Smith LM, Patel CK. Noncontact high-resolution ultra-wide-field oral fluorescein angiography in premature infants with retinopathy of prematurity. *JAMA Ophthalmol* 2014; **132**(1): 108–110.
- 9 Patel CK, Fung TH, Muqit MM, Mordant DJ, Brett J, Smith L *et al*. Non-contact ultra-widefield imaging of retinopathy of prematurity using the Optos dual wavelength scanning laser ophthalmoscope. *Eye* 2013; **27**(5): 589–596.
- 10 Patel CK, Fung TH, Muqit MM, Mordant DJ, Geh V. Non-contact ultra-widefield retinal imaging and fundus fluorescein angiography of an infant with incontinentia pigmenti without sedation in an ophthalmic office setting. *J AAPOS* 2013; **17**(3): 309–311.
- 11 Fung TH, Yusuf IH, Smith LM, Brett J, Weston L, Patel CK. Outpatient ultra wide-field intravenous fundus fluorescein angiography in infants using the Optos P200MA scanning laser ophthalmoscope. *Br J Ophthalmol* 2014; **98**(3): 302–304.

- 12 Kemper AR, Wallace DK, Quinn GE. Systematic review of digital imaging screening strategies for retinopathy of prematurity. *Pediatrics* 2008; **122**(4): 825–830.
- 13 Castellanos MA, Schwartz S, Leal R, Chan RV, Quiroz-Mercado H. Pain assessment in premature infants treated with intravitreal antiangiogenic therapy for retinopathy of prematurity under topical anesthesia. *Graefes Arch Clin Exp Ophthalmol* 2013; **251**(2): 491–494.
- 14 Adams GG, Clark BJ, Fang S, Hill M. Retinal haemorrhages in an infant following RetCam screening for retinopathy of prematurity. *Eye* 2004; **18**(6): 652–653.

Supplementary Information accompanies this paper on *Eye* website (<http://www.nature.com/eye>)

Non-contact ultra-widefield retinal imaging of infants with suspected abusive head trauma

To obtain credit, you should first read the journal article. After reading the article, you should be able to answer the following, related, multiple choice questions. To complete the questions (with a minimum 75% passing score) and earn continuing medical education (CME) credit, please go to www.medscape.org/journal/eye. Credit cannot be obtained for tests completed on paper, although you may use the worksheet below to keep a record of your answers.

You must be a registered user on Medscape.org. If you are not registered on Medscape.org, please click on the new users: Free Registration link on the left hand side of the website to register.

Only one answer is correct for each question. Once you successfully answer all post-test questions you will be able to view and/or print your certificate. For questions regarding the content of this activity, contact the accredited provider,

CME@medscape.net. For technical assistance, contact CME@webmd.net.

American Medical Association's Physician's Recognition Award (AMA PRA) credits are accepted in the US as evidence of participation in CME activities. For further information on this award, please refer to <http://www.ama-assn.org/ama/pub/about-ama/awards/ama-physicians-recognition-award.page>. The AMA has determined that physicians not licensed in the US who participate in this CME activity are eligible for *AMA PRA Category 1 Credits™*. Through agreements that the AMA has made with agencies in some countries, AMA PRA credit may be acceptable as evidence of participation in CME activities. If you are not licensed in the US, please complete the questions online, print the AMA PRA CME credit certificate and present it to your national medical association for review.

1. Your patient is a 5-month-old boy with suspected abusive head trauma (AHT). According to the retrospective, observational case series by Yusuf and colleagues, which of the following statements about findings of non-contact ultra-widefield retinal imaging using the Optos P200MA scanning laser ophthalmoscope (Optos; Dunfermline, UK) among infants being evaluated for AHT is *correct*?
 - A Optos P200MA acquired good quality, high-yield, definitive retinal images in three of five infants
 - B Acute, widespread retinal haemorrhages were documented in three infants, contributing to a diagnosis of AHT
 - C Optos fundus fluorescein angiography (FFA) did not reveal any significant findings
 - D Safety issues were a cause for concern in two infants, in whom imaging had to be stopped
2. According to the retrospective, observational case series by Yusuf and colleagues, which of the following statements about advantages of the Optos P200MA scanning laser ophthalmoscope *vs* RetCam (Clarity Medical Systems, Inc.; Pleasanton, CA, USA) for the documentation of retinal haemorrhages in the context of AHT is *correct*?
 - A The Optos P200MA captures a 130° retinal field
 - B Optos P200MA can acquire high-quality, posterior pole, and peripheral retina views in a single image
 - C With Optos P200MA, colour images of both eyes in this series took ~20 min in each infant
 - D Retinal haemorrhages induced by RetCam have been documented

3. According to the retrospective, observational case series by Yusuf and colleagues, which of the following statements about advantages of RetCam *vs* the Optos P200MA scanning laser ophthalmoscope for the documentation of retinal haemorrhages in the context of AHT is *correct*?
 - A RetCam is portable and can be used for a supine infant too unstable to transfer, or for a pharmacologically sedated or mechanically ventilated infant
 - B RetCam uses the 'flying baby position,' which makes it easier for staff to position the infant
 - C RetCam-assisted FFA may increase diagnostic yield *vs* the Optos P200MA
 - D RetCam produces a uniform focus to multiple retinal locations, simultaneously improving clarity of documentation across the retinal field

Activity evaluation

1. The activity supported the learning objectives.

Strongly disagree				Strongly agree
1	2	3	4	5

2. The material was organized clearly for learning to occur.

Strongly disagree				Strongly agree
1	2	3	4	5

3. The content learned from this activity will impact my practice.

Strongly disagree				Strongly agree
1	2	3	4	5

4. The activity was presented objectively and free of commercial bias.

Strongly disagree				Strongly agree
1	2	3	4	5