the transcervical approach with 16.1/1000 reported malformations in association with the transabdominal approach versus 11.2/1000 with the transcervical approach.

We also were surprised that we did not find an increased incidence of other defects thought to be due to vascular disruption. We offered the hypothesis that this may have been due to underreporting as the total rate of malformations reported was less than would be expected. The underreporting may have had less to do with how noticeable the defect was (i.e., clubfoot vs. cleft lip) than to the fact that some papers focused mainly on limb abnormalities, did not describe the defects completely, lumped defects together, or did not include data on terminations.

We did not use the term *disruptions* to describe the abnormalities reported, as that implies a known etiologic mechanism. At this time it is only theorized how some of these defects may arise. In addition, some of the defects reported, such as cleft lip and palate, may not be due to disruption of previously normally formed tissue as a disruption implies.

Joan M. Stoler, MD
Caroline McGuirk, MPH
Lewis B. Holmes, MD

Genetics and Teratology Unit
 Pediatric Service
 Massachusetts General Hospital and Harvard Medical School
 Boston, Massachusetts

Louise Ryan, PhD
 Division of Biostatistics
 Dana Farber Cancer Institute and Harvard School of Public Health
 Boston, Massachusetts

Ellice Lieberman, MD, DrPH
 Department of Obstetrics and Gynecology
 Brigham and Women's Hospital and Harvard Medical School
 Boston, Massachusetts

Reference

Brumback BA, Cook RJ, Ryan LM. A meta-analysis of case-control and cohort studies with interval-censored exposure data: application to chorionic villus sampling. *Biostatistics* 2000;1:203-217.

To the Editor:

Mao and colleagues¹ recently reported on two new cases with duplication 15q11.2q12, which includes the Prader-Willi syndrome (PWS)/Angelman syndrome (AS) critical region, and described the phenotype for these patients. Their patient with a paternal origin of duplication had PWS-like phenotype while the maternal origin of duplication had an AS-like phenotype. This report was of interest to us because we recently identified a 6-year, 10-month-old female with mild to moderate mental impairment and global developmental delays who had a dup(15)(q11.2q12), confirmed by SNRPN FISH probe.

Mao et al.¹ reviewed previously published cases with 15q proximal duplications, all which have been reported within the past 10 years. Most of these cases were evaluated for parental origin of the duplication. Including their cases, three paternal duplications have shown variable phenotype from normal to developmental delay and behavioral problems. A total of 16 reports of maternal duplication with phenotype ranging from developmental delay to autism and mental retardation have been published. This review of cases in the literature shows considerable variability of expression for both the maternal and paternal duplications. While the duplication in our case was not further characterized by methylation or linkage studies to determine parental origin, her clinical findings are such that it is at the mild end of the phenotypic spectrum that has been reported. We feel that it is important to report additional cases in which parental origin has been documented in order to better define the phenotype with the intent of providing prognostic information to families.

Sharon L. Wenger, PhD
 Department of Pathology
 West Virginia University
 Morgantown, West Virginia

Chet Johnson, MD
 Department of Pediatrics
 West Virginia University
 Morgantown, West Virginia

Reference

1. Mao R, Jalal SM, Snow K, Michels VV, Szabo SM, Babovic-Vuksanovic D. Characteristics of two cases with dup(15)(q11.2-q12): one of maternal and one of paternal origin. *Genet Med* 2000;2:131-135.