Statement on guidance for genetic counseling in advanced paternal age

Helga V. Toriello, PhD¹, and Jeanne M. Meck, PhD², for the Professional Practice and Guidelines Committee

Key Words: paternal age, genetic counseling, mutation, chromosome anomalies

Disclaimer: This guideline is designed primarily as an educational resource for healthcare providers to help them provide quality medical genetic services. Adherence to this guideline does not necessarily assure a successful medical outcome. This guideline should not be considered inclusive of all proper procedures and tests or exclusive of other procedures and tests that are reasonably directed to obtaining the same results. In determining the propriety of any specific procedure or test, the geneticist should apply his or her own professional judgment to the specific clinical circumstances presented by the individual patient or specimen. It may be prudent, however, to document in the patient's record the rationale for any significant deviation from this guideline.

In 1996, a practice guideline on genetic counseling for advanced paternal age was published. The current document updates the state of knowledge of advanced paternal age effects on single gene mutations, chromosome anomalies, and complex traits. **Genet Med 2008;10(6):457–460.**

There is no clearly accepted definition of advanced paternal age. A frequently used criterion is any man aged 40 years or older at the time of conception. The current population mean paternal age is 27 years.

Advanced paternal age is associated with an increased risk of new gene mutations. Because of the large number of cell divisions during spermatogenesis, the mutation rate for base substitutions is much higher in men than women, and increases with paternal age. The risk for genetic defects increases linearly for some conditions, and exponentially for others.^{1–3} The conditions most strongly associated with advanced paternal age are those caused by mutations in the form of single base substitutions in the *FGFR2*, *FGFR3*, and *RET* genes, and include Pfeiffer syndrome, Crouzon syndrome, Apert syndrome, achondroplasia, thanatophoric dysplasia, as well as *MEN2A* and *MEN2B*.⁴ Some dominant conditions that are caused by gene changes that include both point mutations and base pair deletions (e.g., neurofibromatosis) show a lesser association with paternal age. Other dominant conditions show no association with increased paternal age.⁵ Although Friedman⁶ had estimated that the risk for autosomal dominant disorders af-

fecting offspring of fathers aged 40 or more was 0.3–0.5%, it is now thought that the actual risk is lower.⁷ There is also a growing body of evidence that advanced paternal age is associated with an increased risk for complex disorders such as some congenital anomalies, schizophrenia, autism spectrum disorders, and some forms of cancer.^{8–12} For most conditions the relative risk is two or less. However, the mechanism for the increased risk is unknown, and in some cases, the observed paternal age effect may be an artifact of some other causative factor.

In general, for autosomes and sex chromosomes, there is no compelling evidence that chromosomal aberrations (aneuploidy or structural chromosome abnormalities) are significantly increased in newborns as paternal age increases. The low incidence of paternally derived extra chromosomes in trisomies combined with the relatively small number of children fathered by older men makes it difficult to demonstrate a paternal age effect. Two possible exceptions are trisomy 21 and Klinefelter syndrome. Recent data on Down syndrome suggest a paternal age effect, either acting alone or in combination with a maternal age effect.^{13,14} This observation is supported by reports of increased aneuploidy rates in sperm for some of the chromosomes, including 21 and the sex chromosomes.^{15–17} In summary, there is a wide range of genetic disorders that may be related to advanced paternal age (Table 1). Overall, it seems that the risk of birth defects and some chromosome disorders may be minimally increased, and the risk for later onset disorders may also show a small increase with advanced paternal age. There are currently no screening or diagnostic test panels which specifically target those conditions that increase with paternal age. If the older male's partner is currently pregnant, the pregnancy should be treated as any other according to pre-

From the ¹Genetics Services, Spectrum Health, Grand Rapids, Michigan; and ²Quest Diagnostics/Nichols Institute, Inc. Chantilly, Virginia.

Helga V. Toriello, PhD, Genetics Services, Spectrum Health, 21 Michigan St, Ste 465, Grand Rapids, MI 49503-2528. E-mail: helga.toriello@spectrum-health.org.

Approved by the Board of Directors July 28, 2007.

Go to www.geneticsinmedicine.org for a printable copy of this document.

American College of Medical Genetics, 9650 Rockville Pike, Bethesda, MD 20814

Submitted for publication March 4, 2008.

Accepted for publication March 5, 2008.

DOI: 10.1097/GIM.0b013e318176fab

Table 1
Paternal age risks

Type	Specific condition	Age (relative to reference age)	Relative risk (CI, if available)	Population risk (or reference risk)	Adjusted risk	References (first author's name only)
Autosomal dominant	Achondroplasia	>50 (25–29)	7.8	1/15,000	1/1923	Risch ¹
		30–34 (<20)	3.5		1/4285	Tiemann-Boege ²¹
		35–39 (<20)	4		1/3750	
		40–44 (<20)	8		1/1875	
		45–49 (<20)	9		1/1666	
		50–54 (<20)	12		1/1250	
	Apert	>50 (25–29)	9.5	1/50,000	1/5263	Risch ¹
	Pfeiffer	>50 (25–29)	6	1/100,000	1/16,666	Glaser ²²
	Crouzon	>50 (25–29)	8	1/50,000	1/6250	
	Progeria	Unknown	Effect seen	“Exceedingly rare”		
	MEN2A	Unknown	Effect seen	1/30,000		
	MEN2B	Unknown	Effect seen	1/30,000		
	Neurofibromatosis I	>50 (25–29)	3.7 ^a	1/3000–1/4000	1/810–1/1080	Risch ¹
		>40 (<30)	2.9		1/1034–1/1380	Bunin ²³
	Osteogenesis imperfecta	>35 (<25)	2.5	1/10,000	1/4000	Carothers ²⁴
		>35 (<35)	1.37 (0.73–6.89)		1/7300	Orioli ²⁵
	Thanatophoric dysplasia	>35 (<35)	3.18 (1.48–6.89)	1/20,000–1/50,000	1/6290–1/15,723	Orioli ²⁵
	Retinoblastoma	>45	3 ^a (0.21–41.7)	1/15,000–1/20,000	1/5000–1/6667	Dockerty, Yip ^{26,27}
		>35 (<35)	1.34 (1.04–1.74)		1/11,200–1/14,925	Moll ²⁸
		>50 (32.5)	5		1/3000–1/4000	DerKinderen ²⁹
Chromosomal	Down syndrome	40–44 (20–29)	1.37 (0.48–3.86)	1/1200 (mat. age 20–29)	1/876	Zhu ³⁰
		45–49 (20–29)	2.68 (0.76–9.51)		1/448	
		>49 (20–29)	4.5 (1.0–20.3)		1/267	
		40–44 (25–29)	1.45 (1.26–1.68)	Use maternal age as baseline for counseling purposes ^b		Yang ³¹
		45–49 (25–29)	1.28 (1.04–1.57)			
		>49 (25–29)	1.39 (1.04–1.83)			
		None given	“May be increased”			Kuhnert ¹⁶
		None given	“Paternal age effect in association with maternal age (>35) effect”			Fisch ¹⁴
	Klinefelter syndrome	>50 (20's)	1.6 ^c (0.69–3.0)	1/500 men	1/312 men	Lowe ³²
Congenital anomalies	VSD	>40 (<40)	1.69 ^a	1/200	1/118	Olshan ³³
	ASD	>35	1.95 ^a	1/400	1/205	Lian ¹¹
	Tracheoesophageal fistula	>50 (25–29)	2.55 (1.28–4.6)	1/3600	1/1412	Yang ³¹
Other complex disorders	Childhood leukemia	>35	1.5	1/25,000	1/16,667	Murray ³⁴
		>40 (<25)	1.14 (0.85–1.53)		1/21930	Yip ²⁷
	Childhood CNS tumor	30–34 (<25)	1.34 (1.04–1.72)	1/36,000	1/26,866	Yip ²⁷
		35–39 (<25)	1.4 (1.04–1.86)		1/25,714	
		>40 (<25)	1.69 (1.21–2.35)		1/21,302	

(Continued)

Table 1
(Continued)

Type	Specific condition	Age (relative to reference age)	Relative risk (CI, if available)	Population risk (or reference risk)	Adjusted risk	References (first author's name only)	
	Childhood type 1 diabetes	>34 (<25)	1.52 (1.1–2.09)	1/415	1/273	Cardwell ³⁵	
	Epilepsy	35–39	1.18 (1.02–1.26)	1/100	1/85	Vestergaard ³⁶	
		40–45	1.3 (1.08–1.55)		1/770		
	Schizophrenia	>50 (20–24)	4.62 (2.28–9.36)	1/100	1/22	Rasmussen ³⁷	
		35–44 (15–24)	1.6 (1.0–2.6)		1/62.5	Zammit ³⁸	
		45–54 (15–24)	1.6 (0.8–3.1)		1/62.5		
		>54 (15–24)	3.8 (1.3–11.8)		1/26		
		>49 (<25)	3		1/33	Malaspina ¹²	
		>32 (<28)	3 (1.49–6.04)		1/33	Tsuchiya ³⁹	
		Autism	>40 (<30)	5.75 (2.65–12.46)	1/1000	1/174	Reichenberg ⁴⁰
			Unknown	Effect seen			Cantor ⁹
	Autism spectrum disorders	35–39 (25–29)	1.38 (1.04–1.84)	1/200	1/145	Croen ⁴¹	
		>39 (25–29)	1.52 (1.1–2.1)		1/131		
	Breast cancer	>40 (<30)	1.6 (1.04–2.32)	1/8.5	1/5.3	Choi ⁴²	
	Prostate cancer	>38 (<27)	1.7 (1.0–2.8)	1/5.9	1/3.5	Zhang ⁴³	
	Multiple sclerosis	51–55 (21–25)	2.0 (1.35–2.96)			Montgomery ⁴⁴	
Other	Spontaneous miscarriages	>35 (<35)	1.26 (1.0–1.6)	1/7	1/5.3	Slama ⁴⁵	
		>39 (25–29)	1.6 (1.2–2.0)		1/4	Kleinhaus ⁴⁶	
	Relative infertility	>39 (<39)	2.3 (1.67–3.17)	1/14 couples	1/6.2	De la Rochebrochard ⁴⁷	
	Low birth weight	>34 (20–34)	1.7 (1.3–2.2)	1/40	1/23	Reichman ⁴⁸	
	Preeclampsia	35–44 (25–34)	1.24 (1.05–1.46)	1/62	1/50	Harlap ⁴⁹	
Total risk	For 86 examined congenital anomalies	>44 (25–34)	1.8 (1.04–1.51)	1/62	1/34		
		>40 (<20)	1.2	1/50	1/42	Lian ¹¹	
		>50 (<20)	1.3		1/38		

This table is meant to show the findings of various studies examining the effect of paternal age on the condition in question. It is not meant to be a comprehensive guide to counseling, but to merely indicate conditions which have been studied and results obtained from those studies.

^aIncreased risk not shown by other studies.

^bSuggestion for this adjustment made by the author of this document. There are no data regarding use of paternal age for counseling for serum screening results.

^cBased on frequency of XY sperm.

natal diagnosis guidelines established by the American College of Medical Genetics and American College of Obstetricians and Gynecologists,^{18–20} with the prenatal counseling session including a discussion about the potentially increased risk of Down syndrome attributable to increased paternal age. Because of this and the possibility of ultrasound detection of some of the features of the autosomal dominant conditions noted above (e.g., thanatophoric dysplasia), an ultrasound is recommended at 18–20-weeks gestation to evaluate fetal growth and development. However, it is unlikely to detect many of the conditions of interest. Prospective couples should receive individualized genetic counseling to address specific concerns.

RESOURCES

1. Archer NP, Langlois PH, Suarez L, Brender J, et al. Association of paternal age with prevalence of selected birth defects. *Birth Defects Res A Clin Mol Teratol* 2007; 79:27–34.
2. Cedergren MI, Selbing AJ, Kallen BA. Risk factors for cardiovascular malformation—a study based on prospectively collected data. *Scand J Work Environ Health* 2002;28:12–17.
3. Dzurova D, Pikhart H. Down syndrome, paternal age and education: comparison of California and the Czech Republic. *BMC Public Health* 2005;5:69.

4. MacDonald M, Hassold T, Harvey J, Wang LH, et al. The origin of 47, XXY and 47, XXX aneuploidy: heterogeneous mechanisms and the role of aberrant recombination. *Hum Mol Genet* 1994;3:1365–1371.
5. Sherman SL, Freeman SB, Allen EG, Lamb NE. Risk factors for nondisjunction of Trisomy 21. *Cytogenetic Genome Res* 2005;111:273–280.
6. Xue F, Colditz GA, Willett WC, Rosner BA, et al. Parental age at delivery and incidence of breast cancer: a prospective cohort study. *Breast Cancer Res Treat* 2007;104:331–340.

References

1. Risch N, Reigh EW, Wishnick MW, McCarthy JG. Spontaneous mutation and parental age in humans. *Am J Hum Genet* 1987;41:218–248.
2. Crow JF. The high spontaneous mutation rate: is it a health risk? *Proc Natl Acad Sci U S A* 1997;94:8380–8386.
3. Crow JF. Age and sex effects on human mutation rates: an old problem with new complexities. *J Radiat Res* 2006;47(suppl):B72–B82.
4. Jung A, Schuppe HC, Schill WB. Are children of older fathers at risk for genetic disorders? *Andrologia* 2003;35:191–199.
5. Splendore A, Jabs EW, Felix TM, Passos-Bueno MR. Parental origin of mutations in sporadic cases of Treacher Collins syndrome. *Eur J Hum Genet* 2003;11:718–722.
6. Friedman JM. Genetic disease in the offspring of older fathers. *Obstet Gynecol* 1981;57:745–749.
7. Hook EB. Paternal age and effects on chromosomal and specific locus mutations and on other genetic outcomes in offspring. In: Mastroianni L, Paulsen CA, editors. Aging, reproduction, and the climacteric. New York: Plenum Press, 1986:117–145.
8. Byrne M, Agerbo E, Ewald H, Eaton WW, et al. Parental age and risk of schizophrenia: a case-control study. *Arch Gen Psychiatry* 2003;60:673–678.
9. Cantor RM, Yoon JL, Furr J, Lajonchere CM. Paternal age and autism are associated in a family-based sample. *Mol Psychiatry* 2007;12:419–421.
10. Dalman C, Allebeck P. Paternal age and schizophrenia: further support for an association. *Am J Psychiatry* 2002;159:1591–1592.
11. Lian ZH, Zack MM, Erickson JD. Paternal age and the occurrence of birth defects. *Am J Hum Genet* 1986;39:648–660.
12. Malaspina D, Harlap S, Fennig S, Heiman D, et al. Advancing paternal age and the risk of schizophrenia. *Arch Gen Psychiatry* 2001;58:361–367.
13. Shi Q, Martin RH. Aneuploidy in human sperm: a review of the frequency and distribution of aneuploidy, effects of donor age and lifestyle factors. *Cytogenet Cell Genet* 2000;90:219–226.
14. Fisch H, Hyun G, Golden R, Hensle TW, et al. The influence of paternal age on Down syndrome. *J Urol* 2003;169:2275–2278.
15. Sartorelli EM, Mazzucatto LF, de Pina-Neto JM. Effect of paternal age on human sperm chromosomes. *Fertil Steril* 2001;76:1119–1123.
16. Kuhnert B, Nieschlag E. Reproductive functions of the ageing male. *Hum Reprod Update* 2004;10:327–339.
17. Buwe A, Guttenbach M, Schmid M. Effect of paternal age on the frequency of cytogenetic abnormalities in human spermatozoa. *Cytogenet Genome Res* 2005;111:213–228.
18. ACOG Practice Bulletin. Clinical management guidelines for obstetrician-gynecologists. Ultrasonography in pregnancy. No. 58. *Obstet Gynecol* 2004;104:1449–1458.
19. ACOG Practice Bulletin. Screening for fetal chromosomal abnormalities. No. 77. *Obstet Gynecol* 2007;109:217–227.
20. Driscoll DA, Gross SJ, ACMG Professional Practice and Guidelines Committee. First trimester diagnosis and screening for fetal aneuploidy. *Genet Med* 2008;10:73–75.
21. Tiemann-Boege I, Navidi W, Grewal R, Cohn D, et al. The observed human sperm mutation frequency cannot explain the achondroplasia paternal age effect. *Proc Natl Acad Sci U S A* 2002;99:14952–14957.
22. Glaser RL, Jabs EW. Dear old dad. *Sci Aging Knowledge Environ* 2004;21:2004:re1.
23. Bunin GR, Needle M, Riccardi VM. Paternal age and sporadic neurofibromatosis 1: a case-control study and consideration of the methodologic issues. *Genet Epidemiol* 1997;14:507–516.
24. Carothers AD, McAllion SJ, Paterson CR. Risk of dominant mutation in older fathers: evidence from osteogenesis imperfecta. *J Med Genet* 1986;23:227–230.
25. Orioli IM, Castilla EE, Scarano G, Mastroiaco P. Effect of paternal age in achondroplasia, thanatophoric dysplasia, and osteogenesis imperfecta. *Am J Med Genet* 1995;59:209–217.
26. Dockerty JD, Draper G, Vincent T, Rowan SD, et al. Case-control study of parental age, parity and socioeconomic level in relation to childhood cancers. *Int J Epidemiol* 2001;30:1428–1437.
27. Yip BH, Pawitan Y, Czene K. Parental age and risk of childhood cancers: a population-based cohort study from Sweden. *Int J Epidemiol* 2006;35:1495–1503.
28. Moll AC, Imhof SM, Kuik DJ, Bouter LM, et al. High parental age is associated with sporadic hereditary retinoblastoma: the Dutch retinoblastoma register 1862–1994. *Hum Genet* 1996;98:109–112.
29. DerKinderen DJ, Koter JW, Tan KE, Beemer FA, et al. Parental age in sporadic hereditary retinoblastoma. *Am J Ophthalmol* 1990;110:605–609.
30. Zhu JL, Madsen KM, Vestergaard M, Olesen AV, et al. Paternal age and congenital malformations. *Hum Reprod* 2005;20:3173–3177.
31. Yang Q, Wen SW, Leader A, Chen XK, et al. Paternal age and birth defects: how strong is the association? *Hum Reprod* 2007;22:696–701.
32. Lowe X, Eskenazi B, Nelson DO, Kidd S, et al. Frequency of XY sperm increases with age in fathers of boys with Klinefelter syndrome. *Am J Hum Genet* 2001;69:1046–1054.
33. Olshan AF, Schnitzer PG, Baird PA. Paternal age and the risk of congenital heart defects. *Teratol* 1994;50:80–84.
34. Murray L, McCarron P, Bailie K, Middleton R, et al. Association of early life factors and acute lymphoblastic leukaemia in childhood: historical cohort study. *Br J Cancer* 2002;86:356–361.
35. Cardwell CR, Carson DJ, Patterson CC. Parental age at delivery, birth order, birth weight and gestational age are associated with the risk of childhood Type 1 diabetes: a UK regional retrospective cohort study. *Diabet Med* 2005;22:200–206.
36. Vestergaard M, Mork A, Madsen KM, Olsen J. Paternal age and epilepsy in the offspring. *Eur J Epidemiol* 2005;20:1003–1005.
37. Rasmussen F. Paternal age, size at birth, and size in young adulthood—risk factors for schizophrenia. *Eur J Endocrin* 2006;155:S65–S69.
38. Zammit S, Allebeck P, Dalman C, Lundberg I, et al. Paternal age and risk for schizophrenia. *Br J Psychiatry* 2003;183:405–408.
39. Tsuchiya KJ, Takagai S, Kawai M, Matsomoto H, et al. Advanced paternal age associated with an elevated risk for schizophrenia in offspring in a Japanese population. *Schizophr Res* 2005;76:337–342.
40. Reichenberg A, Gross R, Weiser M, Bresnahan M, et al. Advancing paternal age and autism. *Arch Gen Psychiatry* 2006;63:1026–1032.
41. Croen LA, Najjar DV, Fireman B, Grether JK. Maternal and paternal age and risk of autism spectrum disorders. *Arch Pediatr Adolesc Med* 2007;161:334–340.
42. Choi JY, Lee KM, Park SK, Noh D-Y, et al. Association of paternal age at birth and the risk of breast cancer in offspring: a case control study. *BMC Cancer* 2005;5:143.
43. Zhang Y, Kreger BE, Dorgan JF, Cupples LA, et al. Parental age at child's birth and son's risk of prostate cancer. The Framingham study. *Am J Epidemiol* 1999;150:1208–1212.
44. Montgomery SM, Lambe M, Olsson T, Ekblom A. Paternal age, family size, and risk of multiple sclerosis. *Epidemiology* 2004;15:717–723.
45. Slama R, Bouyer J, Windham G, Fenster L, et al. Influence of paternal age on the risk of spontaneous abortion. *Am J Epidemiol* 2005;161:816–823.
46. Kleinhaus K, Perrin M, Friedlander Y, Paltiel O, et al. Paternal age and spontaneous abortion. *Obstet Gynecol* 2006;108:369–377.
47. De La Rochebrochard E, Thonneau P. Paternal age ≥ 40 years: an important risk factor for infertility. *Am J Obstet Gynecol* 2003;189:901–905.
48. Reichman NE, Teitler JO. Paternal age as a risk factor for low birthweight. *Am J Public Health* 2006;96:862–866.
49. Harlap S, Paltiel O, Deutsch L, Knaanie A, et al. Paternal age and preeclampsia. *Epidemiology* 2002;13:660–667.