REVIEWS

DRUG RESISTANCE IN MICRO-ORGANISMS. CIBA Foundation Symposium. Ed. by G. E. W. Wolstenholme and Cecilia M. O'Connor. London: J. & A. Churchill. 1957. Pp. 352. 50s.

This symposium arose out of the desire of Sir Charles Harington, its chairman, to promote fundamental knowledge of chemotherapy, which is at present largely empirical. The aspect considered is that of drug resistance, which makes the empirical successes much less favourable from a strictly utilitarian point of view. Thus in many hospitals over half of the staphylococci causing infections are resistant to penicillin, though perhaps not to other antibiotics. The nature of such resistance and its mode of origin is the core of the discussion. Naturally, the matter is of outstanding interest to geneticists, especially in respect of the controversy between the mutation-selection and "training" schools of thought. This is argued in detail and at length by protagonists who do not seem to convince one another.

The non-genetic viewpoint is stated by Dean and Hinshelwood who "have never doubted that the essential characters of a cell are inherent in the structure of certain fundamental units . . . not . . . easily susceptible to change". They "have never denied that structural mutations leading to increased drug resistance . . . can and do occur" but "have contested the assumption that random mutation and selection is the sole mechanism (or perhaps even the major mechanism) for adaptation to new media or for the development of drug resistance". Consideration of this essay suggests that some of the divergence depends upon a lack of appreciation of genetical principles. They "consider that the term 'heredity' is a rather ambiguous one to use of organisms which multiply by binary fission. . . . there is no question of parent and offspring". In attacking the "uncompromising mutation-selection theory," it is said that "so elaborate a polygenic system is frequently assumed that the results of recombination experiments become indistinguishable from those which would be given by quantitative cytoplasmic changes". Significant reports were given by Lederberg, Pollock and Eagle of mutation-selection situations in Hinshelwood's own material.

Every geneticist would agree that the behaviour of a cell in a given environment is dependent upon the interaction of its genotype and the environment, coloured by the previous physiological history of the cell, as shown so elegantly in the case of β -galactosidase in *E. coli*. They would also agree that there are permanent, or very persistent, hereditary changes not due to changes in structural elements like chromosomes. What is conspicuously lacking in so many of the examples, cited as non-genetic changes, is a genetic analysis of the difference between adapted and unadapted cells.

The list of participants is notable for the presence of two Russians from the Institute of Genetics, Moscow. Kossikov described "directed hereditary changes" of fermentative properties in yeast, but the changes have all the hall-marks of mutation-selection. A most interesting observation is that homozygous adapted cells have twice as much invertase as heterozygous cells. This paper is most significant for the way in which it avoids genetic interpretation.

D. G. CATCHESIDE.