REVIEWS

FARBLOSE ALGEN. Ein Beitrag zur Evolutionsforschung. By E. G. Pringsheim. Fischer Verlag, Stuttgart. 1963. Pp. 471. Price DM 64.

Scattered among the chlorophyll-pigmented algæ of every group—and also the flagellatæ and bacteria—are scores of colourless races which require organic sources of nutrition. From his vast knowledge Dr Pringsheim here describes these races exhaustively, their structure and classification, their requirements of growth and habitat.

Genetically there would appear to be little to say about the colourless algæ since only one form, in the genus *Polytoma*, is known to undergo sexual reproduction. Yet for the evolution of these plants—both their past and their future evolution—this itself is highly significant. Without sexual reproduction we should not expect the colourless algae to evolve new species. And, in fact, outside the special group to which *Polytoma* belongs, there is no evidence that they ever have done so. For each colourless race is paired with a coloured species from which by loss of chlorophyll it may be supposed to have been derived—the reverse process, it would seem, never occurring.

The experimental evidence that certain races of *Euglena gracilis* treated in various ways can lose their chloroplasts (or chromatophores) and become in form and function very like the natural colourless species *Astasia longa* agrees with this view.

A genetic problem of quite a different kind is raised by evolutionary change in the colourless algæ. It is that of the genetic continuity of organellæ. Not only the plastids but the determinants of fibres or flagella are probably irreplaceable when lost. This surprises Dr Pringsheim who thinks in terms of tissue-regeneration rather than of cytoplasmic inheritance. But it may well encourage others to begin with these problems where he has left off.

C. D. DARLINGTON.

NEW PATTERNS IN GENETICS AND DEVELOPMENT. By C. H. Waddington. Columbia University Press. 1962. Pp. xv+271. \$10.00.

The aim of this book, which is based on the Jesup lectures that the author gave at Columbia University in 1961, is to consider the origin of new patterns of structure in developing cells and tissues in the light of the new patterns of thought arising from recent advances in biology, particularly microbiological genetics and ultrastructural biology of tissues. The book is timely, for many biologists are now realising that advancing knowledge of gene structure and nature, and of gene-protein relationships has put us in a position to look in new ways at one of the oldest, least understood but most characteristically biological of the problems of biology: the problem of the organisation and development of form.

The author begins by discussing differentiation at the biochemical level, for “one can hardly discuss new patterns without devoting a good deal of attention to new substances”, in terms of “gene-action systems” which he divides into “gene-protein” and “protein-phenotype” phases. Jacob and Monod’s work on induction and repression of enzymes of *E. coli* provides “a good example of the type of conceptual scheme required”, and leads to discussion whether differentiation, initiated by gene-cytoplasm