REVIEWS

COMPUTER APPLICATIONS IN GENETICS. Edited by Newton E. Morton. University of Hawaii Press, Hawaii, 1969.

This book is accurately described by its title: it is a collection of papers on a variety of topics read at a meeting in Hawaii. Some of the papers merely describe the details of computer programmes written to estimate gene frequencies, genetic linkage, analyse human pedigrees and so on. One paper, indeed, is concerned only with the numerical analyses of the equations of maximum likelihood.

Other papers are of more general interest to human geneticists. Newton Morton describes his work on the problems of ascertainment in the analysis of genetic segregation. He gives the methods applicable to a very wide range of cases. Walter Bodmer discusses the use of association analysis in the study of multispecific sera and describes the results obtained for the HL-A leukocyte system. And J. R. Renwick discusses the evidence from the computation of likelihoods for a number of autosomal linkage groups in Man.

Three papers give results which should be of general interest to students of evolution. B. D. H. Latter discusses his attempts to formulate general models of the selection of quantitative characters. His models can be described by certain parameters giving the distribution of gene effects, linkage and so on. The goodness of fit of the different models can then be compared to the results of selection experiments. It is, of course, quite possible that no one model will ever be adequate to fit the results of many different selection experiments. But if one or two models can be found to give a generally good fit, Latter's models will be of very great value to the plant and animal breeder and for the study of natural selection: by fitting the comparatively few parameters of the models, the progress of selection will then be predictable.

Bruce Levin discusses the general "assets and liabilities" of genetic simulation, although he omits to mention its use for estimation and prediction, which, if Latter is successful, will perhaps become the most important asset. Levin is mainly concerned with the simulation of complex ecological and genetical situations as they are likely to occur in nature. As he points out, in a computer simulation all parameters and variables must be precisely defined whereas in a purely verbal argument they can be left undefined for intuition to get work on. Levin's own simulations of genetic variability in competing species show how unreliable purely verbal arguments can be. He found that there was no "genetic feed back" that could lead to the co-existence of competing species as suggested by Pimental *et al.* (*Am. Naturalist*, 993, 97-108, 1965) and that co-existence could only occur if the two species evolved to avoid competition—that is, if they diverged. Thus by his computer simulations, Levin appears to have beautifully confirmed Darwin's original theory of evolutionary divergence as a mechanism to avoid competition—a theory which does not seem to have been taken very seriously.

A. W. F. Edwards discusses the results of his attempts to grow evolutionary trees by computer. Apparently, the form of the tree cannot be determined using even the simplest model of evolution by genetic drift, and

Edwards is able to compute the form of a tree only by using the method that minimises the total amount of evolution. As he says, more complex models would allow all possible trees to be produced if the appropriate directional selection is postulated. The main conclusion is that non-adaptive characters will be the best to use in studying phylogenies because the amount of divergence will be proportional to the amount of evolution. Characters strongly subject to directional selection will therefore only confuse the matter. Taxonomists have mainly used characters like bone structures which are very constant in general form and probably therefore subject to stabilising selection. They will have undergone relatively little evolution. This may be the reason for the general agreement between evolutionary trees produced by the method of minimum evolution and generally accepted phylogenies.

PETER O'DONALD

Nuffield Unit of Medical Genetics, University of Liverpool

QUANTITATIVE GENETICS IN SHEEP BREEDING. H. Newton Turner and S. S. Y. Young. Macmillan, 1969. Pp. 332. \$20.

It has been one of the oddities of the application of quantitative genetics to animal improvement that, in spite of the fact that almost all of the theory referred to the concept of heritability and its value in comparing *a priori* different kinds of selection programmes, very few workers actually carried out selection on domestic animals, the notable exception being Lerner's poultry flock in California. In the early 1950's, a series of selection experiments for single characteristics in Merino sheep were started by Dr Turner in Australia as well as one line selected for maximum wool weight with restriction on other traits. The results of these experiments have only recently begun to appear in the literature, and the publication of a book by Dr Turner and one of her colleagues, Dr Young, is a major event for scientists concerned with animal improvement. This is a field in which authoritative up-to-date texts at an advanced level are almost non-existent. The experimental results and their application to practical problems of the improvement of wool production are discussed within the framework of a theoretical treatment involving a considerable amount of detailed algebra.

I suspect that I will use the book in the main to refer to the Australian experimental work. In that a better title would have been "The application of genetics in the improvement of the Australian Merino", it has some limitations as a textbook either of quantitative genetics or of sheep breeding. It became clear from early experimental results that with the main component of economic value, clean wool weight, having a high heritability and being measurable in both sexes, a programme of individual selection on a combination of several traits is best for the improvement of the Merino. In so far as the theoretical sections deal directly with this theme, they are good but away from it the standard is not so high. The sections on part-record selection (without any reference to Lerner's work on poultry and omitted from the index), progeny testing and the utilisation of heterosis fall into this category. The authors were clearly placed in some difficulty in writing a book on sheep improvement that an adequate theoretical text was not