REVIEWS

BASIC BACTERIOLOGY AND GENETICS. J. R. Sokatch and J. J. Ferretti. Yearbook Publishers Inc., Chicago, 1976. Pp. xii+264.

The growth of college and university courses in microbiology has provided the publishing industry with an opportunity to increase the number of texts that cover this subject. This has resulted in two distinct categories of books. The first, and most useful, are those general texts which can be used to complement a lecture course but are also considered to be works of reference upon the subject. The other group of books are those that are written with a specific audience in mind, usually a particular student intake. Books of this last category provide an introduction to a subject for a large number of students without intending to be a treatise. It is surprising therefore that Sokatch and Ferretti's book, which falls into this latter category, should deal only briefly with two fundamental areas of importance in bacteriology, pure culture technique and the identification and classification of bacteria. Perhaps this is information that is covered in practical classes; one hopes so. There is a trend, however, towards teaching those areas of a subject that are interesting and omitting others that are less stimulating. Unfortunately these two topics fall into this latter category and as they are essential to any training in bacteriology, their scant treatment is a serious failing of this book.

One useful feature of this book is the devotion of a separate chapter to antimicrobial agents. Usually this information is hidden in the chapters that deal with the physiology and biochemistry of bacteria. The other chapters deal with the structure of bacteria, their physiology, growth, and genetics and with gene expression. As is so common nowadays, models are described to the exclusion of the data that generated them. This saves space but does not lead to a full understanding of the systems. This book therefore requires certain supplementary information before it may be recommended as a course book.

PHILIP OLIVER

Department of Genetics, University of Cambridge

A DISCUSSION ON THE MEIOTIC PROCESS: R. Riley, M. D. Bennett and R. B. Flavell, (Organisers). *Philosophical Transactions of the Royal Society of London, B*, Vol. 277, 184-376. Royal Society, London, 1977. Price: £17.05.

This publication contains the contributions to a meeting held in December 1975. It was fitting that the meeting should be opened by Darlington with a characteristically thought-provoking and stimulating, if controversial, contribution. As Riley and Flavell said in their introduction the "meeting was organised to provide an opportunity for an assessment of the present state of our understanding and in the hope that at the same time it would provide indications of the most important directions for future research". The overall impression obtained from reading these papers is our lack of understanding on all the basic issues! Nevertheless, as H. Stern said in

his concluding remarks, we are now better able to formulate the problems to be tested than we were a decade ago, and the three main approaches of cytology, genetics, and biochemistry have more potential for discovery when combined than when pursued in isolation. Such combined approaches now appear feasible in attempting to answer some of the many outstanding questions about meiosis.

A few examples will illustrate the uncertainty that this publication reveals on fundamental issues. Riley and Flavell pointed out that although only a small fraction of the DNA can meet its homologue in the synaptonemal complex, yet recombination can occur at many sites. They suggested that the incorporation of DNA into the complex may be determined largely at random. Stern and von Wettstein each favour the idea that the regions of the DNA involved in recombination are selected in advance and trapped within the synaptonemal complex during its assembly. Maguire produced evidence supporting the notion that commitment of a particular site to crossing-over, or the event itself, must have occurred before pairing of the entire chromosome. Callan and Perry drew attention to the widespread occurrence of quite different patterns of localisation of chiasmata in the two sexes of the same organism, and yet these patterns are not associated with restriction on pairing. They also pointed out that chiasmata were never found in the transcribing regions of lampbrush chromosomes. Lindsley and Sandler showed how crossover frequency in *Drosophila* varies in a characteristic way with position in each chromosome, and they referred to Beadle's experiment of over 40 years ago that demonstrated that cross-over frequency was influenced by proximity to a centromere. But, as Riley and Flavell stated, there is reason to believe that recombination is initiated only at specific sites. Might not these be situated in the interchromomeric regions at leptotene, and all be incorporated in the synaptonemal complex in each meiotic cell? Which of them became recombination sites in a particular cell would then depend on controlling factors, some evidently tissue-specific to account for the differences between the sexes in hermaphrodite organisms. Bateson *et al.* (1923, *J. Genet.*, 13, 232) actually used the sex difference in cross-over frequency in *Primula sinensis* as an argument against the chromosome theory! Over 50 years later we are not much nearer understanding either the cause of this difference or its significance.

Several of the contributors referred to the perennial problem of interference between cross-overs, and Holliday discussed the puzzling fact that recombination events not involving flanking marker exchange do not show interference. He proposed that interference resulted from the presence within each synaptonemal complex of a limited amount of a DNA-binding protein that was able to stabilise cross-overs, perhaps forming the recombination nodules reported by Carpenter. Could it be that one of the functions of the synaptonemal complex is to restrict isomerisation (if that is the method by which flanking markers become recombinant) and that this enables controlling factors to determine the nature of each recombination event, that is, whether it is to be a cross-over or not? Unfortunately, the prospect of answering this question looks remote, since study will be required of flanking marker behaviour in allelic recombinants under conditions where chiasma distribution can be varied, for example, as between the sexes in organisms such as the grasshopper *Stethophyma grossum* or the flowering plant *Fritillaria meleagris* discussed by Callan and Perry. Alternatively, can a

technique be devised to detect the transfer or exchange of minute quantities of DNA between homologues at meiosis?

The firmest evidence of what goes on in meiosis was provided by Stern and Hotta's contribution. Their elegant biochemical studies with *Lilium* provide clear evidence for two important conclusions: (1) that each succeeding phase (prezygotene, zygotene, pachytene and postpachytene) has its own characteristic set of metabolic processes apparently concerned with commitment to meiosis, chromosome pairing, recombination, and chromosome segregation, respectively; and (2) that the DNA contains regions with specifically meiotic functions. At present *Lilium* stands alone, but it would seem to be a first priority to attempt to ascertain how far Stern and Hotta's conclusions apply generally. Lindsley and Sandler's genetic studies with meiotic mutants of *Drosophila* seem to be in agreement.

This publication is clearly essential reading for all who are interested in any aspect of meiosis. Errors are few though it was a surprise to read (p. 201 *et seq.*) that yeast and *Chlamydomonas* are prokaryotes! Thirty excellent plates beautifully illustrate the cytological contributions—and no doubt add considerably to the price of the work. But need this have been set quite so high?

H. L. K. WHITEHOUSE

Botany School, University of Cambridge, Downing Street, Cambridge

EVOLUTION. T. Dobzhansky, F. J. Ayala, G. L. Stebbins and J. W. Valentine. W. H. Freeman, San Francisco, 1977. Pp. 572. £11.60.

This book follows in the tradition of the distinguished synoptic treatises on evolution by Dobzhansky, Stebbins, Mayr and Simpson, which in the modern idiom owes its origin to Julian Huxley's pioneering review, "Evolution The Modern Synthesis", published 35 years ago. At that time Mendelian methods had uncovered sufficient of the relation between the gene and the phenotype, both at the level of the organism and of the population, to provide a coherent view of the processes of evolution. But the techniques used to study the molecular nature of hereditary materials and the chemistry of evolutionary responses are a recent introduction. It is the *forte* of the present volume that these exciting developments are at the centre of the synopsis. Thus, although much of the discussion on topics like, the genetic structure of populations, the origin of hereditary variation and patterns of speciation, has a very familiar ring, the supporting evidence has been so carefully up-dated, and subjected to a rigorously contemporary interpretation that readers will experience a stimulating degree of freshness throughout the presentation.

The first half of the book covers the cytogenetic basis of the response of individuals and populations to evolution: the most crucial details of micro-evolution including those uncovered by the authors, especially Dobzhansky and Stebbins, are concisely documented in this section. The discerning student will enjoy the introductory chapter on the nature of evolution and on its status as a core, cultural concept in the life sciences; a theme that is effectively developed in later chapters.

Chapter nine deals with phylogenetic relationships as revealed by analyses of the biologically important macromolecules. As befits the recent