REVIEWS

STOCHASTIC MODELS IN POPULATION GENETICS: W.-H. Li (Ed.). Dowden, Hutchinson and Ross. Stroudsburg, Pennsylvania, 1977. Pp. xiii+471. £22.50.

This reprinting of thirty-one papers in the stochastic theory of population genetics is as welcome an addition to the literature as it is, in some ways, disappointing. It is very expensive, so that the only scholars who will be able to afford it are those who are likely to have access to the original papers anyway; it is poorly printed, some of the editorial matter being in typescript and the quality of the facsimile reproductions varying from adequate to poor; and the selection of early work is open to criticism. Yet the introduction is a helpful guide to the literature, and there is no denying the benefit of having so many essential papers bound together.

The risk attached to such a collection is that it will be used by future generations as a definitive source-book, and that the editor's choice and comments will be accepted uncritically. Thus we read "For only two or three decades have mathematicians been aware of mathematical problems in population genetics". Karl Pearson was 3rd in the Cambridge Mathematical Tripos (3rd "Wrangler") in 1879; G. H. Hardy 4th Wrangler in 1898; H. T. J. Norton 7th Wrangler in 1907; R. A. Fisher took "Firsts" in both parts of the reorganised Mathematical Tripos in 1910 and 1912; J. B. S. Haldane took a First in Mathematics Moderations in his first year at Oxford (1912). With such antecedents, even as distinguished a modern contributor as P. A. P. Moran will blush to find himself described as "one of the first few mathematicians who became aware of the mathematical problems in genetics".

The collection starts with R. A. Fisher's 1922 paper, following four type-written pages of S. Wright's reminiscences (Fisher and J. B. S. Haldane are given the briefest biographical sketches in the book, the reader being referred elsewhere). There is a growing misunderstanding of the place of this paper in the history of the subject, to which this volume contributes. "Paper 1 by Fisher is a history-making paper for several reasons. . . . This was the first study of the maintenance of genetic variability in terms of stochastic models. Considering a population not so much as an aggregate of living individuals but as an aggregate of genes, Fisher introduced two very important models. The first was the branching process (or the so-called Galton-Watson process), in which he used the method of generating function to study the survival of a mutant gene in a large population. The second model was the simple heat diffusion equation. . . ." The quotation is given at length because it exemplifies the modern American use of the word "model" which is the source of some of the misunderstanding. In England the word "model" refers to the mathematical construct describing a process, whilst the word "method" refers to the mathematical technique used to analyse the model. Thus the branching process is a model, but the use of generating functions to study it is a method, as in the quotation. The editor goes on, however, to call the diffusion equation a "model" in accordance with modern American usage which, for example, refers to the method of maximum likelihood as a "model for estimation". This usage

has obscured the early history of the subject, with the result that many of the later papers in this volume refer to "Wright's model" when describing Fisher's 1922 model. Using the words in their British sense, the early contributions can be assigned as follows.

In 1922 Fisher introduced (1) the branching-process model, using functional iteration as a method of analysis; (2) the chain-binomial model, erroneously ascribed to Wright by many authors, including, in the present volume, Ewens (paper 15), Moran (paper 27), and Karlin and McGregor (papers 28 and 29), together with the associated concept of the distribution of gene frequencies, and in particular the stationary distribution; and (3) the method of analysis for this model involving the diffusion approximation and its associated partial-differential equation. By contrast, in 1931 Wright introduced (1) the concept of effective population size and the treatment of the rate of decrease of heterozygosity, using his inbreeding coefficient method, and (2) the method of finding the stationary gene-frequency distributions on Fisher's model by solving integral equations.

The second paper in the volume is by Haldane, and is the only one of his included; but then he wrote, in *The Causes of Evolution*, "The investigation of the case where the total population is finite has been wholly due to Fisher and Wright".

The next three papers (and seventy-nine pages) are by Wright, and are mostly concerned with finding the stationary gene-frequency distributions under various extensions of Fisher's model. The longest is the 1931 paper: "a monumental work. It still has not found its peer". The student who relies on this part of the volume (Part I: "Pioneering Works") for his knowledge of the origins of the subject may wonder what Fisher was doing whilst Wright was producing this monumental work without peer, and he should satisfy his curiosity by reading *The Genetical Theory of Natural Selection*, pp. xiv+272, Oxford, Clarendon Press 1930 (second revised edition: Dover Publications, 1958), a book which Haldane described as "brilliant" and Wright considered "certain to rank as one of the major contributions to the theory of evolution". No part of this book is reproduced in the present volume, nor does the editor mention it explicitly in his introductory review. Moreover, Fisher's 1930 paper "The distribution of gene ratios for rare mutations" is not reproduced on the grounds that it will in due course be reproduced elsewhere (in another volume in the same series as the present one). This is most unsatisfactory, for the whole point of a volume of reprints is balance and completeness—all the papers are available elsewhere.

Fisher had seen a draft of Wright's 1931 paper in 1929, and thus in 1930 was able to correct the form of the diffusion equation which he had given erroneously in 1922. He then obtained "the solution for steady decay" followed by the solution for "neutral mutations". Noting that the integral of the gene-frequency distributions thus obtained fails to converge, Fisher then developed a method of functional equations to investigate these two discrete distributions more accurately. In both cases he established the exact forms for the terminal classes and gave the corresponding diagrams in *The Genetical Theory* (pp. 87 and 91). These two diagrams compare with figs. 3 and 5 of Wright (1931). Moreover, Wright, for his part, specifically states that the selections on the distributions of gene frequencies headed "Selection" and "General formula" were rewritten "in order to take

account of the correction of my formula, suggested by Fisher's results in *The Genetical Theory*". By omitting Fisher's 1930 contributions the present volume can only contribute to the misunderstanding which surrounds the early history of the subject.

Part II, "Classical single-locus problems", reprints papers by M. Kimura, A. Robertson and W. J. Ewens from the period 1955–1963, and Part III "Molecular Population Genetics" contains papers by Kimura (sometimes jointly), Ewens and M. Nei, from 1964 to 1972, together with a typewritten copy of an extract from an article by Wright in the 1949 *Encyclopaedia Britannica*: "It is remarkable that an article dated 1949 is the first paper for this section. Four years before Watson and Crick (1953) proposed their model of DNA structure, Wright had already used the model of infinite alleles to study the number of alleles that could be carried by a population". However, a year earlier still, G. Malécot (1948) had introduced precisely this model in *Les mathématiques de l'hérédité*, concluding that the equilibrium value of the inbreeding coefficient is $1/(4Nu + 1)$ where N is the effective population size and u the mutation rate, each new mutation being distinct. This finding, often attributed to Kimura and Crow (1964), would have been a worthy addition to the volume. The editor writes that he had hoped to include another part of the English translation of Malécot's book, but could not obtain permission to reprint it. Perhaps the original French publisher, Masson et Cie, would have been more co-operative.

The last paper in Part III, Nei on genetic distance, prompts a rather strange editorial comment: "In the last several decades, a number of measures of genetic distance have been proposed, but most of them are constructed from the statistical point of view, and it is not clear what biological unit they are going to measure". Nei's method is to be preferred. But at least one of the measures referred to was created specifically to account for gene-frequency divergence due to random genetic drift, and it seems strange to dismiss it in a book called *Stochastic Models in Population Genetics*.

Part IV, "Multilocus Problems", contains four papers and is principally concerned with two linked loci, whilst Part V, "Structured Populations", contains five papers, starting with Wright's "Isolation by distance".

Part VI is rather curiously called "Mathematical Problems" and contains well-known papers by W. Feller, P. A. P. Moran, S. Karlin and J. McGregor, and G. A. Watterson, and a recent (1975) paper by M. F. Norman. These mostly relate to the Markov-chain approach to Fisher's model or to Moran's model (here introduced for the first time). It is a pity that the paper first applying Markov methods to Fisher's model is not included: Malécot (1944). The modern reader will too easily conclude from the papers in this section that the central model of stochastic population genetics was introduced by Wright and the method of analysis by Feller, rather than by Fisher and Malécot respectively.

The major lacuna in the volume is the absence of any papers on the stochastic properties of populations of known structure. Neither the work on regular mating systems (from H. S. Jennings, 1916 and Wright, 1921, to Fisher's *Theory of Inbreeding*, 1949) nor the work on arbitrary genealogies (from C. W. Cotterman, 1940, to the modern developments by M. Gillois, A. Jacquard, and others) is included.

The detailed consideration of a volume such as this raises the question of its real purpose. If it is to introduce students to stochastic theory in

genetics, it is likely to confuse all but the best, but if it is to introduce the new generation to the history of the subject, it is likely to create a distorted impression. Perhaps its greatest value will be to those who, with discrimination, write the text and tell the history.

A. W. F. EDWARDS

Gonville and Caius College, Cambridge

DNA INSERTION SEQUENCES, PLASMIDS AND EPISOMES. Edited by A. I. Bukhari, J. A. Shapiro and S. L. Adhya. Cold Spring Harbor Laboratory, Nov. 1977. Pp. xii + 782. \$36 postpaid.

This admirable book documents one of the recent important surprises in genetics—the discovery of the lability of the genome, especially in prokaryotes but to an unknown but significant extent in eukaryotes also. A potent combination of genetic methods and the newer techniques of hetero-duplexing and endonuclease fragmentation reveals not only that bacteria can acquire ready-assembled kits of useful accessory genes in the form of plasmids, but also that the plasmids themselves are in a state of flux through transpositions from one to another of blocks of genes. The whole system resembles a series of Chinese boxes; the large transposable blocks, which may be tens of kilobases in terms of DNA, often contain smaller transposable units (*transposons*) of the order of one or a few kilobases. Furthermore, the property of transposability itself depends on small (of the order of hundreds of bases) repeated sequences flanking the transposable segments; under the name of IS, or inserted sequences, these repeats had earlier been identified as individually transposable elements which, though apparently near-random insertion in the chromosome, account for a significant proportion of spontaneous mutation in *E. coli*.

The key to transposability in general appears to be the presence of repeats (usually inverted, *i.e.* palindromic) at the ends of the sequences transposed. This applies both to the transposons and large gene blocks, where the repeats are provided by whole IS or similar elements, and to the IS elements themselves, which, in two cases so far analysed, have palindromic or near-palindromic sequences at their ends. It looks very much as if the significance of the palindromes must be in providing recognition sequences for a series of site-specific recombination enzymes. It is envisaged that such an enzyme or enzyme system can “see” and cut the two identical palindrome ends simultaneously, lift out the intervening sequence and reinsert it in a new location. How this model relates to the cases where the flanking repeats are in direct, rather than inverted, orientation with respect to one another is not altogether clear, but it may be that the general (*rec A*) recombination system, operating through sequence homology rather than site specificity, may have a role in these cases.

These phenomena, or something like them, find a parallel in eukaryotes in the classical work of McClintock on maize, to which has been more recently added a number of examples of transposable elements in *Drosophila*, due to the work of Green and others. What is so far known of the labile mating type systems in both of the popular yeasts (*Saccharomyces* and *Schizosaccharomyces*) are strongly suggestive of controlled DNA rearrangements, and we know that DNA rearrangements are (though omitted from this book) involved in at least one case of cellular differentiation—