

THE GENETICAL RESPONSE TO NATURAL SELECTION BY VARIED ENVIRONMENTS II. OBSERVATIONS ON REPLICATE POPULATIONS IN SPATIALLY VARIED LABORATORY ENVIRONMENTS

C. S. HALEY AND A. J. BIRLEY

Department of Genetics, University of Birmingham, P.O. Box 363, Birmingham B15 2TT,
England

Received 18.iv.83

SUMMARY

From each of two populations of *Drosophila melanogaster*, collected two months previously, from Chateau Tahbilk, S. Australia and Groningen, The Netherlands, duplicate populations were initiated in each of four environments which differed in their degree of environmental heterogeneity. Differing combinations of three food media based on oatmeal/treacle, potato or fig were used to simulate levels of environmental heterogeneity within the populations. The polymorphic loci, *Adh*, *Est-6*, *G-6pdh*, α -*Gpdh*, *Pgm*, *Lap-D* and *Aph* in both the Chateau Tahbilk and Groningen derived populations and *6Pgdh*, which was only polymorphic in the populations which came from Chateau Tahbilk, were monitored in the experiment. The populations maintained a size of about 2500 adults and were sampled after 16 and 32 generations.

Large changes of phenotype frequency were shown by all loci. Despite a frequent divergence of phenotype frequencies between duplicate cages, systematic effects of occasion and environment were present and allele frequencies at many loci were shown to be changing at a faster rate than could be due to random genetic drift.

Genetic heterozygosity differed between environments but was not positively correlated with degree of environmental heterogeneity.

1. INTRODUCTION

An approach to the assessment of the effects of selection by different or varied environments upon enzyme and other protein polymorphisms is to study changes in genotype frequency in laboratory populations. This approach has been used to study the effects of different types of environment and also to obtain an estimate of the fraction of loci showing a selective response to the imposed environmental regimes. Examples of this method are provided by Powell (1971), McDonald and Ayala, (1974) and Minawa and Birley (1975, 1978). Although such studies do not conclusively demonstrate that the loci under observation are themselves the object of selection, they still exemplify the short-term response to selection and may be indicators of the extent of selection by variation in the natural environment.

An additional advantage of the study of laboratory populations in artificial or modelled environments is that it permits the examination of the maintenance of polymorphisms by, for example, either spatial or temporal variation of the environment. Theoretical models of spatial variation in which individuals mate at random and traverse the whole range of environments show that conditions for gene frequency equilibrium are very limited (Gillespie, 1974, Templeton and Rothman, 1974; Strobeck, 1975).

The conditions become slightly less restrictive if individuals remain for a significant part of their lifespan in one environmental niche but mating is at random over all niches (Levene, 1953; Levins and MacArthur, 1966; Maynard Smith, 1966). If individuals both mate and produce progeny in a particular niche, the conditions under which genetic variation can be maintained become much less stringent (Levins and MacArthur, 1966, Levins, 1968).

Cyclical variation in the environment is also capable of maintaining genetical variation (Dempster, 1955; Haldane and Jayakar, 1963). Recently, studies of the effects of random fluctuations in the environment over time with particular reference to enzyme polymorphisms have shown that these may maintain genetical variation (Gillespie and Langley, 1974; Gillespie, 1976).

Previous studies with laboratory populations subject to environmental variation have provided evidence for a positive association between environmental heterogeneity and genetic heterozygosity. Powell (1971) has shown that the rate of loss of heterozygosity at a number of allozyme loci in *Drosophila willistoni* is retarded in more variable environments. A similar study was reported by McDonald and Ayala (1974) with *Drosophila pseudoobscura*.

In the present study a set of environments was created with a gradient of spatial environmental heterogeneity such that the more heterogeneous environments contained discrete environmental niches. Spatial environmental variation was produced by way of differing food media, the only temporal variation present in the design were aging effects of the food media, population density variation and random environmental effects. Thus the design provided a test of the effect of environmental heterogeneity upon genic heterozygosity and additionally a test for effects of environmental niches upon the enzyme polymorphism. Furthermore, two different recently collected natural populations were studied in the same set of environments in order to test for any generality of response, and the replication of large populations in all major environments allowed a comprehensive assessment of results.

2. MATERIALS AND METHODS

(i) *The sampled populations*

Two populations of *Drosophila melanogaster* were used in this experiment. They were a) the "Groningen" population, collected in a fruit market in The Netherlands by Dr W. Van Delden and b) the "Chateau Tahbilk" population, collected from a vineyard in Southern Australia by Dr A. MacKenzie. Both populations were collected as circa 100 isofemale lines and were maintained for three generations in the laboratory before the experiment commenced.

(ii) *The experimental populations*

For both the "Groningen" and "Chateau Tahbilk" populations four hundred virgin flies (equal numbers of both sexes) were collected from each isofemale line and placed into a population cage. The flies were allowed to intermingle for 4 hours and then the populations were each divided to

derive to sub-populations from each of the "Groningen" and "Chateau Tahbilk" populations. Twenty vials each containing a standard oatmeal/molasses medium (see below) were placed in every cage and the populations left for 18 hours at 25°C. After this period all of the vials were removed, discarded and replaced with fresh vials containing the same food medium. By this time most of the female flies are gravid and hence potential contributors to the next generation. After a further 24 hours at 25°C the vials were removed from the cages and the food medium in the vials was removed and quartered. Each quarter was then placed into a $\frac{1}{3}$ pint milk bottle again containing the standard oatmeal/molasses (see later) food medium. Each bottle received about 300 eggs and the food medium was sufficient to support at least 500 larvae, without intense competition, at 25°C. The emergent adults were collected from all of the bottles and placed in a population cage. After intermingling for 4 hours the flies were removed and 2000 flies, equal numbers of both sexes, were placed into the experimental cages. All cages were maintained at 25°C and originally contained 18 tubes of standard oatmeal/molasses food medium.

(iii) *The experimental environments*

Three types of food medium were used in different combinations with each other. They were standard oatmeal/molasses medium (type A), fig based (type B) and potato based (type C). Type A medium consisted of 72 g oatmeal (Mornflake, fully stabilized); 35 g molasses (Fowler's); 6 ml Nipagin-M as a 1 per cent solution in 95 per cent ethanol; 6 gms agar (Köbe) and 400 mls water. Type B was 100 g dried figs subsequently rehydrated and homogenised in 500 ml water; 15 g dried flaked, killed baker's yeast; 5 g agar (Köbe) and 5 mls Nipagin solution. Type C comprised 50 g dried potato; 500 ml water; 15 g dried flaked yeast; 5 g agar (Köbe) and 5 ml Nipagin solution. Each vial of food medium contained 8 ml of one of these media, medium type A was additionally topped with 1 ml of live baker's yeast suspension. Four types of environment were established from both the "Groningen" and "Chateau Tahbilk" derived populations. The environments can be represented by three letters each representing one of the three vials of food medium, with which the populations were supplied for their maintenance. The four environments were: (1) AAA; (2) AAB; (3) AAC; (4) ABC. Hence the ABC environments were supplied with one vial of each of the three types of medium on each feeding occasion. Duplicate cages of all environments were established there being 16 cage populations in all. In addition, a series of populations was also established with the environmental compositions, ABB, ACC, BBB, BBC, BCC and CCC. For reasons of space, this series was unreplicated and only set up with flies from the "Groningen" population.

The cages were maintained at 25°C. The cycle of media vial changes was over two weeks, three vials being changed on Monday, Wednesday and Friday of the first week and Tuesday and Thursday of the second week. Each cage contained 18 vials of medium thus the mean time a vial spent in a cage was 16.8 days. By this time all flies had emerged and the food medium had been fully utilised.

The experimental size necessitated the use of two identical incubators. In an attempt to compensate for the effect of any slight difference between

the incubators, the population cages were randomly re-allocated to the various positions within the incubators every time new food was supplied. This procedure has the effect of distributing over cages any environmental variance between different incubators and shelves within incubators.

(iv) *Sampling the environments*

All populations were sampled (see below) 11 and 22 months after their establishment. Allowing a generation interval of 3 weeks for *Drosophila melanogaster* populations with overlapping generations (Barker, 1962), these periods represent generations 16 and 32 respectively and will be referred to as sampling occasions one and two. Additionally, all environments containing only one type of medium or all three types of media were sampled two months, *i.e.*, three generations, after the initiation of the experiment, this will be referred to as sampling occasion zero.

Three sub-samples of eggs were taken from every cage when it was sampled, hence all the component types of medium were sampled from each cage. The minimum number of individuals typed from each locus from each of the three sub-samples within a population cage on each occasion was approximately 30. There was no more variation between sub-samples within cages than would be expected due to chance, thus the sub-samples within a population cage were pooled for the purposes of analysis. The egg samples were raised at low density at 25°C in culture bottles containing medium of the same type as that upon which the eggs were sampled. Third instar larvae were collected for typing with respect to the *Adh* locus and late pupae for typing at the *Lap-D* locus. Emergent flies were aged for 4 days after collection and then typed with respect to the *Adh*, *Est-6*, *G-6pdh*, *Pgm* and α -*Gpdh* loci in populations. *6-Pgdh* was only polymorphic and hence typed in the series of populations derived from "Chateau Tahbilk". After collection all samples were stored at -20°C prior to gel electrophoresis.

(v) *Gel electrophoresis*

Gel-electrophoresis of individual fly homogenates was carried out in 12 per cent starch gels (Connaught Medical Laboratories, Toronto). Poulik's (1957) discontinuous buffer system was used to type *Lap-D*, *Adh*, *Est-6* and α -*Gpdh*. A Tris-citrate continuous buffer system pH 7.0 (Ayala, 1972), was used to type *G6Pdh*, *Pgm* and *6Pgdh* and a tris-borate-EDTA continuous buffer was used for *Adh* (Wallis and Fox, 1968).

(vi) *Data*

All polymorphic loci studied had two electromorphs apart from the *Est-6* and *Pgm* loci in the "Chateau Tahbilk" population which both had two additional rare electromorphs. All electromorphs present in the "Groningen" population were also present in the "Chateau Tahbilk" population. The amount of data collected precludes its full inclusion here. However, a summary of the data, in the form of the frequency of one allele class from each locus in each population cage on each sampling occasion, is given in table 1 for the "Chateau Tahbilk" populations and in table 2

for the "Groningen" populations. The allele classes shown in Tables 1 and 2 are those which were commonest in the "Chateau Tahbilk" populations.

3. RESULTS

(a) *Phenotypic frequencies*

The variation in phenotypic frequencies at a locus was analysed in respect of environments, sampling occasions and replicate populations using the log-likelihood ratio statistic G^2 or $2I$ (Kullback, 1959). The analysis permits the total homogeneity of phenotypic frequency at a locus to be partitioned in accordance with table 3. The partitions of the total homogeneity of phenotype frequencies of any one locus represent:

(1) homogeneity of phenotypes over environments, irrespective of sampling occasion and replicate population (environments).

(2) homogeneity of phenotypes over occasions, irrespective of environments (occasions).

(3) the homogeneity of phenotypes over environments conditional upon occasions (environments \times occasions).

(4) the homogeneity of phenotypes over replicates within environments (replicates within environments).

(5) the homogeneity of replicates within environments conditional upon occasions (replicates within environments \times occasions).

The appropriate analyses for "Chateau Tahbilk" populations are shown in table 4 and for "Groningen" populations, in Table 5. All loci, except *Pgm* and *Est-6* in "Chateau Tahbilk" were di-allelic (i.e., had two allele mobility classes) and thus had three phenotypic classes. However, at the *Adh*, *G-6Pdh* and *Pgm* loci in the "Groningen" population one homozygote was quite rare and was pooled with the heterozygote for the purposes of analysis in order to avoid having classes with expected numbers of less than 4.

Pgm and *Est-6* are tetra-allelic in the "Chateau Tahbilk" population with allele frequencies such as to give expected numbers some classes of phenotype less than 4. Consequently the rarest phenotypic classes corresponding to these loci were pooled to give classes with expected values greater than 4. This pooling resulted in three phenotypic classes for *Pgm* and four phenotypic classes for *Est-6*.

The G^2 or $2I$ values from each of the five items in this analysis are tested for significance by comparison with the χ^2 -distribution for the appropriate degrees of freedom. The analyses show a substantial change in phenotype frequency due to divergence between replicate populations. This is shown both in the item testing homogeneity between replicate populations within environments (item 4), and the item testing the homogeneity of replicate populations within environments and occasions (item 5), which are often significant at least the 5 per cent level. A significant G^2 value for one of the main effects, that is the environments item (item 1), the occasions item (item 2) on the environments \times occasions item (item 3), can only be interpreted as demonstrating an effect in the absence of a significant G^2 value for the appropriate error item. Thus the effect of environments (item 1) was assessed relative to the variation between replicate cages within environments (item 4) and the effects of occasions (item 2) and

TABLE I
 "Chateau Tahbilk" population-allele class frequencies

Adh	Sampling Occasion			Est-6	Environment	Sampling Occasion		
	0	1	2			0	1	2
	AAA ₁	0.736	0.725	0.733	AAA ₁	0.633	0.699	0.477
	AAA ₂	0.744	0.696	0.634	AAA ₂	0.637	0.751	0.672
	AAB ₁	—	0.735	0.316	AAB ₁	—	0.756	0.644
	AAB ₂	—	0.756	0.439	AAB ₂	—	0.511	0.5
	AAC ₁	—	0.685	0.767	AAC ₁	—	0.781	0.667
	AAC ₂	—	0.770	0.801	AAC ₂	—	0.825	0.815
	ABC ₁	0.779	0.741	0.684	ABC ₁	0.711	0.707	0.689
	ABC ₂	0.702	0.654	0.528	ABC ₂	0.623	0.644	0.673

G-6Pdh	Sampling Occasion			Pgm	Environment	Sampling Occasion		
	0	1	2			0	1	2
	AAA ₁	0.812	0.684	0.661	AAA ₁	0.831	0.791	0.689
	AAA ₂	0.864	0.878	0.861	AAA ₂	0.822	0.689	0.622
	AAB ₁	—	0.822	0.639	AAB ₁	—	0.955	0.917
	AAB ₂	—	0.756	0.511	AAB ₂	—	0.928	0.928
	AAC ₁	—	0.787	0.7	AAC ₁	—	0.888	0.872
	AAC ₂	—	0.867	0.811	AAC ₂	—	0.794	0.744
	ABC ₁	0.811	0.822	0.922	ABC ₁	0.858	0.872	0.867
	ABC ₂	0.814	0.708	0.477	ABC ₂	0.844	0.773	0.911

RESPONSE TO VARIED ENVIRONMENTS

α -GPdh	Environment	Sampling Occasion			6-PGdh	Environment	Sampling Occasion		
		0	1	2			0	1	2
	AAA ₁	0.771	0.789	0.922	AAA ₁	0.706	0.444	0.281	
	AAA ₂	0.781	0.817	0.85	AAA ₂	0.910	0.934	0.939	
	AAB ₁	—	0.741	0.833	AAB ₁	—	0.756	0.628	
	AAB ₂	—	0.600	0.828	AAB ₂	—	0.824	0.783	
	AAC ₁	—	0.756	0.844	AAC ₁	—	0.808	0.789	
	AAC ₂	—	0.806	0.811	AAC ₂	—	0.784	0.817	
	ABC ₁	0.695	0.789	0.889	ABC ₁	0.866	0.776	0.783	
	ABC ₂	0.686	0.706	0.764	ABC ₂	0.832	0.693	0.509	

Lap-D	Environment	Sampling Occasion			Aph	Environment	Sampling Occasion		
		0	1	2			0	1	2
	AAA ₁	0.814	0.861	0.933	AAA ₁	0.5	0.471	0.404	
	AAA ₂	0.795	0.761	0.811	AAA ₂	0.528	0.424	0.289	
	AAB ₁	—	0.639	0.528	AAB ₁	—	0.611	0.494	
	AAB ₂	—	0.822	0.783	AAB ₂	—	0.506	0.278	
	AAC ₁	—	0.850	0.872	AAC ₁	—	0.610	0.367	
	AAC ₂	—	0.889	0.868	AAC ₂	—	0.459	0.331	
	ABC ₁	0.783	0.711	0.728	ABC ₁	0.452	0.382	0.4	
	ABC ₂	0.805	0.653	0.628	ABC ₂	0.531	0.461	0.354	

TABLE 2
 "Groningen" population allele class frequencies

Adh	Environment	Sampling Occasion			Est-6	Environment	Sampling Occasion		
		0	1	2			0	1	2
	AAA ₁	0.888	0.920	0.806		AAA ₁	0.748	0.655	0.811
	AAA ₂	0.868	0.906	0.956		AAA ₂	0.788	0.670	0.744
	AAB ₁	—	0.875	0.917		AAB ₁	—	0.659	0.806
	AAB ₂	—	0.856	0.893		AAB ₂	—	0.685	0.706
	AAC ₁	—	0.912	0.822		AAC ₁	—	0.620	0.622
	AAC ₂	—	0.929	0.878		AAC ₂	—	0.803	0.753
	ABC ₁	0.886	0.933	0.933		ABC ₁	0.712	0.713	0.635
	ABC ₂	0.887	0.867	0.882		ABC ₂	0.732	0.655	0.644
	ABB	—	0.917	0.875		ABB	—	0.600	0.617
	ACC	—	0.944	0.917		ACC	—	0.710	0.656
	BBB	0.883	0.899	0.867		BBB	0.711	0.584	0.489
	BBC	—	0.916	0.869		BBC	—	0.706	0.667
	BCC	—	0.933	0.989		BCC	—	0.747	0.539
	CCC	0.858	0.854	0.806		CCC	0.708	0.591	0.680

G-6Pdh	Environment	Sampling Occasion			Pgm	Environment	Sampling Occasion		
		0	1	2			0	1	2
	AAA ₁	0.954	0.928	0.944		AAA ₁	0.975	0.961	0.787
	AAA ₂	0.927	0.861	0.944		AAA ₂	0.949	0.894	0.830
	AAB ₁	—	0.956	0.917		AAB ₁	—	0.962	0.783
	AAB ₂	—	0.914	0.989		AAB ₂	—	0.883	0.878
	AAC ₁	—	0.941	0.856		AAC ₁	—	0.949	0.928
	AAC ₂	—	0.978	0.961		AAC ₂	—	0.922	0.9
	ABC ₁	0.924	0.922	0.939		ABC ₁	0.973	0.983	0.989
	ABC ₂	0.942	0.945	0.9		ABC ₂	0.946	0.967	0.939
	ABB	—	0.956	0.967		ABB	—	0.962	0.939
	ACC	—	0.967	0.944		ACC	—	0.967	0.85
	BBB	0.944	0.890	0.933		BBB	0.948	0.950	0.95
	BBC	—	0.928	0.961		BBC	—	0.978	0.966
	BCC	—	0.939	0.933		BCC	—	0.959	0.961
	CCC	0.935	0.956	0.967		CCC	0.954	0.950	0.994

α -GPDh	Environment	Sampling Occasion		
		0	1	2
AAA ₁	AAA ₁	0.513	0.450	0.478
AAA ₂	AAA ₂	0.481	0.444	0.317
AAB ₁	AAB ₁	—	0.528	0.594
AAB ₂	AAB ₂	—	0.483	0.428
AAC ₁	AAC ₁	—	0.575	0.706
AAC ₂	AAC ₂	—	0.450	0.467
ABC ₁	ABC ₁	0.498	0.506	0.567
ABC ₂	ABC ₂	0.448	0.461	0.528
ABB	ABB	—	0.528	0.561
ACC	ACC	—	0.639	0.717
BBB	BBB	0.290	0.471	0.611
BBC	BBC	—	0.578	0.606
BCC	BCC	—	0.539	0.472
CCC	CCC	0.466	0.450	0.439

Lap-D	Environment	Sampling Occasion			Aph	Environment	Sampling Occasion		
		0	1	2			0	1	2
AAA ₁	AAA ₁	0.774	0.900	0.719	AAA ₁	0.523	0.715	0.651	
AAA ₂	AAA ₂	0.831	0.844	0.806	AAA ₂	0.595	0.667	0.556	
AAB ₁	AAB ₁	—	0.861	0.831	AAB ₁	—	0.657	0.494	
AAB ₂	AAB ₂	—	0.891	0.822	AAB ₂	—	0.681	0.722	
AAC ₁	AAC ₁	—	0.876	0.805	AAC ₁	—	0.708	0.738	
AAC ₂	AAC ₂	—	0.814	0.739	AAC ₂	—	0.608	0.661	
ABC ₁	ABC ₁	0.816	0.826	0.789	ABC ₁	0.517	0.551	0.578	
ABC ₂	ABC ₂	0.782	0.772	0.742	ABC ₂	0.573	0.651	0.640	
ABB	ABB	—	0.806	0.761	ABB	—	0.625	0.517	
ACC	ACC	—	0.778	0.883	ACC	—	0.703	0.590	
BBB	BBB	0.763	0.852	0.742	BBB	0.549	0.602	0.744	
BBC	BBC	—	0.850	0.9	BBC	—	0.591	0.683	
BCC	BCC	—	0.761	0.606	BCC	—	0.676	0.670	
CCC	CCC	0.75	0.847	0.683	CCC	0.517	0.840	0.644	

TABLE 3
The log-likelihood analysis of phenotype frequency

Item	degrees of freedom	2I or G
1. Phenotypes × Environments	$(q-1)(s-1)$	$2 \sum_{i,k} \sum_{i,k} n_{i,k} \ln \frac{n_{i..k} N}{n_{i..} n_{.k}}$
2. Phenotypes × Occasions	$(q-1)(t-1)$	$2 \sum_{i,l} \sum_{i,l} n_{i..l} \ln \frac{n_{i..l} N}{n_{i..} n_{..l}}$
3. Phenotypes × Environments × Occasions	$(q-1)(s-1)(t-1)$	$2 \sum_{i,k,l} \sum_{i,k,l} n_{i,k,l} \ln \frac{n_{i..k..l} N}{n_{i..} n_{.k..} n_{..l} N}$
4. Phenotypes × Replicates within environments	$s(q-1)(r-1)$	$2 \sum_{i,j,k} \sum_{i,j,k} n_{ij,k} \ln \frac{n_{ij,k} N}{n_{.j,k} n_{i..}}$
5. Phenotypes × Replicates within environments and occasions	$s(q-1)(r-1)(t-1)$	$2 \sum_{i,j,k,l} \sum_{i,j,k,l} n_{ijkl} \ln \frac{n_{ijkl} N}{n_{.jkl} n_{i..} n_{.jk..} n_{..l}}$
where Phenotypes (i) = 1, q	$N = n_{....} = \sum_{i,j,k,l} n_{ijkl}$	
Replicates (j) = 1, r	and $n_{..} = \sum_i n_{ij,k,l}$	
Environments (k) = 1, s		
Occasions (l) = 1, t	$n_{i..l} = \sum_{i,j,k} n_{ijkl}$ etc.	

environments \times occasions (item 3) were assessed relative to item 5, replicate cages within environments \times occasions.

The only cases in which the main effects are significant and the appropriate error terms are not significant are: in the "Chateau Tahbilk" populations; the occasions and environments \times occasions items for the *Adh* locus and the occasions items for the *Aph* locus; and, in the "Groningen" populations; the occasions and environments \times occasions items for the *Est-6* locus, the environments item for the *Pgm* locus and the occasions item for the *Lap-D* locus. As an approximate, and probably conservative, test for those main effects which are significant as G^2 values but which had significant error terms, an F -ratio test following that described by Minawa and Birley (1978) was performed. The F -ratio is constructed from the G^2 value for the item to be tested over its degrees of freedom divided by the appropriate error term, also over its degrees of freedom. This test reveals a further two cases in which the effect of occasions is significantly greater than its appropriate error; for the *G-6Pdh* and α -*Gpdh* loci in the "Chateau Tahbilk" population, and a further one case in which the effect of environments is significantly greater than its appropriate error; for the *Adh* locus in the "Chateau Tahbilk" population. Thus for every locus studied, other than *6-Pgdh*, in one or other of the two populations studied, there is evidence for systematic changes in phenotype frequency indicative of the action of selection. For all loci other than *6-Pgdh* and *Pgm* in one or other of the populations there is evidence for an effect of occasions. However, for only three loci (*Adh* in "Chateau Tahbilk" and *Est-6* and *Pgm* in "Groningen") does the presence of a significant environments or environments \times occasions item demonstrate any differential effect of the experimental environments.

Inspection of tables 4 and 5 reveals little similarity between the "Groningen" and "Chateau Tahbilk" populations in the systematic responses of individual loci. There are no instances where a locus produces a significant occasions item in both populations, thus there is little evidence that pure directional selection acting solely on the loci under observation. Additionally, it does not seem as though the two populations are converging on common allele frequencies; inspection of tables 1 and 2 reveals that the general trend of allele class frequencies for a locus in a population for which there is a significant occasions item, is away from the prevailing allele class frequency in the other population for all six loci with a significant occasions item apart from *Lap-D*. Similarly, there is little concordance between the responses to the experimental environments in the two populations. The two loci which produced significant environments items were *Adh* (in "Chateau Tahbilk") and *Pgm* (in "Groningen", although the F ratio in "Chateau Tahbilk" is approaching significance: $p = 0.056$) the mean common allele class frequency for each of the four replicated environments in each population on sampling occasion 2 for these loci are shown below:

Environment	<i>Adh</i>		Environment	<i>Pgm</i>	
	Chateau Tahbilk	Groningen		Chateau Tahbilk	Groningen
AAA	0.684	0.881	AAA	0.656	0.809
AAB	0.378	0.905	AAB	0.923	0.831
AAC	0.784	0.850	AAC	0.808	0.914
ABC	0.606	0.908	ABC	0.889	0.964

TABLE 4
 "Chateau Tahbilk"—phenotype frequency analysis

Item	Phenotypes	Environments (E) Occurrences (O)	D.F.	G	P	F	P
<i>Adh</i>	xEnvironments	(E)	6	107.84	<0.001***	4.49	0.028*
	xOccurrences	(O)	2	82.17	<0.001***	58.80	<0.001***
	xExO		6	44.92	<0.001***	10.71	0.002**
	xReplicates within xReplicates within	E ExO	8 8	31.76 5.59	<0.001*** 0.69		
<i>Est-6</i>	xE		9	77.95	<0.001***	1.22	0.366
	xO		3	17.33	<0.001***	2.35	0.124
	xExO		9	27.36	0.001***	1.23	0.361
	xR xR	E ExO	12 12	85.12 29.54	<0.001*** 0.003**		
<i>G-6Pdh</i>	xE		6	27.50	<0.001***	0.25	0.921
	xO		2	33.12	<0.001***	5.03	0.039*
	xExO		6	20.77	0.002**	1.05	0.461
	xR xR	E ExO	8 8	144.64 26.36	<0.001*** <0.001***		
<i>Pgm</i>	xE		6	131.32	<0.001***	3.42	0.056
	xO		2	5.27	0.072	9.99	0.007***
	xExO		6	8.11	0.230	5.12	0.019*

<i>α-GPd_h</i>	P	XR	8	51.18	<0.001***			
		XO	8	2.11	0.977			
		XE	6	19.71	0.003**	1.44	0.308	
<i>6-PGd_h</i>	P	XO	2	35.55	<0.001***	5.60	0.030*	
		XE	6	14.23	0.027*	0.75	0.627	
		XR	8	18.21	0.020*			
		XO	8	25.41	0.001**			
		XE	6	92.79	<0.001***	0.42	0.847	
		XO	2	3.30	0.192	0.59	0.577	
<i>Lap-D</i>	P	XE	6	20.04	0.003**	1.20	0.394	
		XO	6	297.23	<0.001***			
		XR	8	22.32	0.004**			
		XE	6	115.96	<0.001***	2.52	0.113	
		XO	2	0.22	0.896	0.13	0.880	
		XR	8	11.87	0.065	2.43	0.122	
<i>Aph</i>	P	XE	8	61.44	<0.001***			
		XO	8	6.52	0.589			
		XE	6	14.86	0.021*	0.64	0.698	
		XO	2	40.96	<0.001***	12.56	0.003**	
		XR	6	10.80	0.095	1.10	0.437	
		XR	8	30.82	<0.001***			
		XO	8	13.04	0.110			

TABLE 5
Groningen—phenotype frequency analysis

	Phenotypes	Item	(E) (O)	D.F.	G	P	F	P
Adh		xEnvironments	(E)	3	2.36	0.501	0.15	0.924
		xOccasions	(O)	1	1.17	0.279	0.35	0.586
		xExO		3	14.37	0.002**	1.43	0.298
		xReplicates within	E	4	21.08	<0.001***		
		xReplicates within	ExO	4	13.41	0.009**		
Est-6	P	xE		6	8.31	0.216	0.41	0.853
		xO		2	8.52	0.014*	3.93	0.065
		xExO		6	14.58	0.024*	2.24	0.144
		xR	E	8	27.14	<0.001***		
		xR	ExO	8	8.68	0.370		
G-6Pdh	P	xE		3	3.31	0.346	0.30	0.825
		xO		1	0.03	0.862	0.01	0.925
		xExO		3	14.85	0.002**	1.04	0.465
		xR	E	4	15.03	0.005**		
		xR	ExO	4	18.99	<0.001***		

<i>Pgm</i>	P	xE	3	63.18	<0.001***	9.30	0.028*
		XO	1	33.95	<0.001***	5.37	0.103
		xExO	3	8.33	0.040*	0.59	0.653
		XR	4	9.06	0.060		
<i>α-GPdh</i>		E	4	18.96	<0.001***		
		ExO	4	18.96	<0.001***		
	P	xE	6	35.08	<0.001***	1.09	0.442
		XO	2	4.19	0.100	0.94	0.430
<i>Lap-D</i>		xExO	6	8.77	0.187	0.66	0.685
		XR	8	42.96	<0.001***		
		XR	8	17.75	0.023*		
	P	xE	6	12.37	0.054	1.79	0.218
<i>Aph</i>		XO	2	20.15	<0.001***	7.53	0.014*
		xExO	6	7.39	0.286	0.92	0.527
		XR	8	9.19	0.327		
		XR	8	10.71	0.219		
<i>Aph</i>	P	xE	6	11.42	0.076	0.44	0.834
		XO	2	3.01	0.222	1.26	0.334
		xExO	6	8.84	0.183	1.23	0.382
		XR	8	34.40	<0.001***		
	ExO	8	9.55	0.298			

As inspection of these figures reveals, there is no similarity between the two populations for the *Adh* locus; the environments producing the two lowest frequencies in "Chateau Tahbilk" produce the highest ones in "Groningen". At the *Pgm* locus the AAA environment produces the lowest frequency in both populations but there is little similarity between the other environments.

In order to compare the overall degree of change in the two populations the average value of G^2 (i.e., the total value of G^2 over the total degrees of freedom) over all loci for each of the five items in the analysis were calculated and are shown below:

	Chateau Tahbilk		Groningen	
	d.f.	Average G^2	d.f.	Average G^2
Phenotypes \times Environments (E)	51	11.53	33	4.12
\times Occasions (O)	17	12.82	11	6.46
\times E \times O	51	3.10	33	2.34
\times Replicates within E	68	10.59	44	3.61
\times Replicates within E \times O	68	1.92	44	2.23

It can be seen from these figures that the first three averages are all larger than their errors and that this is especially so for the two occasions items. Also most items in the "Chateau Tahbilk" population are appreciably larger than in the "Groningen" population. This is despite the fact that the item measuring replicate divergence between sampling occasions 1 and 2 (phenotypes \times replicates within E \times O) is approximately the same in both populations. The large averages for phenotypes \times replicates within environments compared with those for phenotypes \times replicates within environments \times occasions indicates that most replicate divergence occurred prior to sampling occasion 1.

(b) *Allele frequencies*

Analysis of phenotypic frequencies have demonstrated a good deal of genetic change has occurred in the set of populations. A small proportion of the response to selection can be attributed or related to the effect of environments. A substantial response to selection has, however, occurred between the two times of sampling, i.e., over 16 generations. It also appears that some of the populations may be evolving uniquely and largely independently of their environments. Therefore an analysis of the data is required which examines changes of allele frequency in replicate populations making allowance for the effects of random genetic drift. That is, we require to separate the effects of natural selection and random genetic drift as causes of the non-systematic changes seen in replicate populations. The experiment spanned approximately 32 generations and the observed population sizes (N) maintained in the cages were about 2500 to 3000 adults. In this case allele frequency changes observed between sampling occasions must be largely due to natural selection since the number of generations is much less than the effective population size (N_e) provided that N_e/N for *Drosophila melanogaster* is taken to be about 0.8 (Crow and Morton, 1955).

A test for selection taking into account the effect of random genetic drift is available from the method of Fisher and Ford (1947). These authors define the sampling variance of the observed allele frequencies around the population allele frequency as

$$V(p_i) = \frac{\hat{p}_i \hat{q}_i}{2n_i} - \frac{\hat{p}_i \hat{q}_i}{2N_i}$$

where N_i is the total population size on sampling occasion i , n_i is the sample taken on occasion i and p_i and q_i are the allele frequencies on occasion i .

This relationship is for sampling from a finite population without replacement. In the present experiment the population samples were taken as samples from continuously produced egg-population rather than a population of adult flies. Hence we can omit the quantity $p_i q_i / 2N_i$. Fisher and Ford (1947) demonstrated that following the angular transformation $\theta_i = \sin^{-1} \sqrt{p_i}$, the sampling variance of θ was independent of its magnitude and

$$\begin{aligned} V(\theta_i) &= \frac{180^2}{4\pi^2} \frac{\text{Var}(p_i)}{p_i q_i} \\ &= \frac{820 \cdot 7}{2n_i} \end{aligned}$$

For a chosen reference generation (r) the variance of the observed allele frequency about its actual value in the reference generation is $820 \cdot 7 / 2n_r$. The variance of the transformed observed allele frequency in any generation (i) about its actual value in the referenced generation due to sampling and random genetic drift then becomes $820 \cdot 7 / 2n_i + x \cdot 820 \cdot 7 / 2N$ where x is the number of generations separating the two samplings. A chi-square test for selection is obtained as:

$$\chi^2(j-1) = \sum_i T_i \theta_i - \hat{\theta} \sum_i T_i$$

where j is the number of samples taken, T is a matrix obtained as the product of θ , a column vector of the observed values of θ_i , and I the inverse matrix of the expected variances of the estimates of θ_i around its population value in the reference year; these values are arrayed along the leading diagonal of the matrix. The maximum likelihood estimate $\hat{\theta}$ of θ in the reference year is

$$\hat{\theta} = \frac{\sum_i T_i}{\sum_{ij} I_{ij}}$$

we of course assume random union of gametes.

Data was available from all population cages on sampling occasions 1 and 2 (approximately generations 16 and 32 respectively) and this was supplemented with data from occasion 0 (generation 3) where available. The *Est-6* and *Pgm* loci in the "Chateau Tahbilk" population have four alleles but as for all other loci, only changes in the commonest allele class were included on the analysis. The results are presented in table 6 for the "Chateau Tahbilk" populations and in table 7 for the "Groningen" populations. Given N_e is about 2000 all loci show significant χ^2 values. Hence the allele frequency changes are much too rapid to be accounted for by random

TABLE 6
Changes in allele frequency between occasions—Chateau tahbilk population

Environment	<i>Adh</i>	<i>Est-6</i>	<i>G-6pdh</i>	<i>Pgm</i>	α - <i>Gpdh</i>	<i>6-Pdgh</i>	<i>Lap-D</i>	<i>Aph</i>	Average χ^2
AAA ₁	0.090 N.S.	16.098***	13.257***	6.944*	14.649***	57.016***	7.798*	2.383 N.S.	7.452
AAA ₂	3.584 N.S.	5.872 N.S.	0.219 N.S.	16.302***	2.064 N.S.	0.815 N.S.	1.136 N.S.	7.856	2.366
AAB ₁	47.532***	3.921*	11.608***	1.655 N.S.	3.171 N.S.	5.052*	3.377 N.S.	3.650 N.S.	9.996
AAB ₂	27.612***	0.034 N.S.	17.501***	0.132 N.S.	17.454***	0.685 N.S.	0.634 N.S.	14.562***	9.826
AAC ₁	2.196 N.S.	4.343 N.S.	2.633 N.S.	0.148 N.S.	3.305 N.S.	0.150 N.S.	0.274 N.S.	15.795***	3.606
AAC ₂	0.371 N.S.	0.050 N.S.	1.522 N.S.	0.935 N.S.	0.013 N.S.	0.427 N.S.	0.272 N.S.	4.459*	1.006
ABC ₁	3.266 N.S.	0.940 N.S.	8.582*	0.152 N.S.	17.159***	6.009*	2.542 N.S.	1.821 N.S.	2.529
ABC ₂	9.026*	0.629 N.S.	36.253***	4.487 N.S.	2.174 N.S.	16.781***	13.961***	8.182***	5.718
Average χ^2	7.806	2.657	7.631	2.563	4.999	7.245	2.583	3.576	5.580

The chi-squares from the AAA and ABC environments have 2 degrees of freedom, all others have 1 degree of freedom.

TABLE 7
Changes in allele frequency between occasions—Groningen population

Environment	Adh	Est-6	G-6pdh	Pgm	α -Gpdh	Lap-D	Aph	Average χ^2
AAA ₁	7.678*	8.713*	1.002 N.S.	27.274***	1.212 N.S.	17.972***	15.343***	5.689
AAA ₂	7.329*	6.970	7.024*	10.981**	8.231*	0.695 N.S.	3.782 N.S.	3.215
AAB ₁	1.233 N.S.	7.306*	1.712 N.S.	21.602***	1.164 N.S.	0.446 N.S.	6.978*	5.777
AAB ₂	0.858 N.S.	0.127 N.S.	9.676*	0.019 N.S.	0.824 N.S.	2.535 N.S.	0.585 N.S.	2.089
AAC ₁	5.790*	0.002 N.S.	6.832*	0.541 N.S.	5.710*	2.533 N.S.	0.335 N.S.	3.106
AAC ₂	2.269 N.S.	0.889 N.S.	0.632 N.S.	0.405 N.S.	0.074 N.S.	2.122 N.S.	0.778 N.S.	1.024
ABC ₁	3.271 N.S.	2.283 N.S.	0.317 N.S.	1.054 N.S.	1.411 N.S.	0.573 N.S.	1.004 N.S.	0.708
ABC ₂	0.363 N.S.	3.112 N.S.	2.159 N.S.	1.482 N.S.	1.839 N.S.	3.335 N.S.	2.656 N.S.	1.068
Average χ^2	2.399	2.450	2.446	5.317	1.872	2.518	2.622	2.803
ABB	1.233 N.S.	0.077 N.S.	0.220 N.S.	1.160 N.S.	0.297 N.S.	0.772 N.S.	3.102 N.S.	0.980
ACC	0.798 N.S.	0.907 N.S.	0.782 N.S.	12.133***	1.839 N.S.	5.360*	4.310*	3.733
BBB	0.686 N.S.	16.289***	3.949 N.S.	0.010 N.S.	9.089*	0.477 N.S.	12.034**	3.038
BBC	1.484 N.S.	0.465 N.S.	1.433 N.S.	0.323 N.S.	0.211 N.S.	1.390 N.S.	2.434 N.S.	1.106
BCC	6.405**	12.601***	0.034 N.S.	0.009 N.S.	1.178 N.S.	20.953***	0.010 N.S.	5.884
CCC	1.594 N.S.	6.474*	1.964 N.S.	7.107*	0.246 N.S.	11.142**	70.499***	7.073
Average χ^2	2.049	3.311	1.887	4.228	1.666	3.515	6.193	3.264

The chi-squares from the AAA, ABC, BBB and CCC environments have 2 degrees of freedom, all others have 1 degree of freedom.

genetic drift alone. If only environments common to both sets of populations are considered, the average χ^2 is greater in the "Chateau Tahbilk" (5.58) than in the Groningen population (2.80). Inspection of the average χ^2 for loci shows least evidence for selection (i.e. low average χ^2 values) at the *Lap-D*, *Est-6* and *Pgm* loci in "Chateau Tahbilk" and similarly at the *Adh*, *G6pdh* and α -*Gpdh* loci in the "Groningen" populations.

As predicted from earlier results replicate populations can differ considerably in behaviour. There are cases where a replicate population shows several highly significant χ^2 values and its counterpart none. Also allele frequencies can change in quite different directions; in the ABC environment of the "Chateau Tahbilk" population one replicate shows the common *G-6Pdh* allele to be at 0.811, 0.822 and 0.922 to be on the three respective sampling dates in one replicate and 0.814, 0.708 and 0.477 respectively in the duplicate population.

This example is, however, unusual, for of the 35 out of 60 cases in which there is a significant change in allele frequency in one or both duplicate populations, in 26 cases (when 3 sampling dates were included for analysis) the allele frequency changes in replicate cages were in the same direction. Hence there is a great deal of evidence to support the action of natural selection.

It might be argued that the effective size in experimental populations is substantially lower than that which we have used, and that the progeny of very few females contribute to the individual vial populations. However, in this experiment we have not detected significance between sub-samples within population cages implying that N_e is not small. Nonetheless we can suitably amend the analysis of data using the "Fisher-Ford" method by reducing N_e until the average χ^2 in the experiment is such as to give 5 per cent of the cases statistically significant, i.e., by chance alone. This would give an effective population size of approximately 150 in "Chateau Tahbilk" and 240 in Groningen populations, giving a value of N_e/N of about 0.06-0.1 which seems highly unlikely given the size and vigour of the adult populations.

Although random genetic drift must be invoked as at least the initiator of replicate differences within environments, there is little doubt that natural selection has been a major cause of the changes in allele frequencies seen during the course of this experiment.

(c) *Genic heterozygosity*

The design of this experiment provides an ideal opportunity to examine the relationship between environmental heterogeneity and genic heterozygosity. If, as has been predicted by some authors, there is a positive relationship between environmental heterogeneity and genic heterozygosity, environments containing three types of food medium should be on average more heterozygous than those containing two types of medium which should in turn be more heterozygous than those containing a single type of medium. In order to examine this hypothesis the average observed heterozygosity for each level of environmental heterogeneity on each occasion was calculated and is shown in table 8.

Inspection of Table 8 reveals no evidence for a simple positive relationship between environmental heterogeneity and genic heterozygosity.

TABLE 8

Average observed heterozygosity for each environmental type and level of heterogeneity for the "Chateau Tahbilk" and "Groningen" populations on sampling occasions 1 and 2

Level of Heterogeneity	Environment	"Chateau Tahbilk"		"Groningen"	
		Occasion 1	Occasion 2	Occasion 1	Occasion 2
1 type of Medium	AAA	0.3949	0.3682	0.2951	0.3163
2 types of Medium	AAB	0.3541	0.3878	0.3046	0.2965
	AAC	0.3207	0.3164	0.2693	0.3159
	Average	0.3374	0.3521	0.2870	0.3060
3 types of Medium	ABC	0.4239	0.3960	0.3185	0.3212

However, there is some evidence that heterozygosity varies with environmental heterogeneity as environments containing two types of food medium have the lowest average heterozygosity. Thus the metric warrants further investigation.

To facilitate analysis the average proportion of heterozygotes for every locus at every cage at each sampling occasion was transformed using the arcsine transformation. The analyses of variance of the transformed data are shown in table 9 for each combination of population ("Chateau Tahbilk" or "Groningen" derived) and sampling occasion. The variation between environments was partitioned into three orthogonal comparisons designed to locate the possible sources of any differences between the environments. The coefficients corresponding to the three comparisons C_1 , C_2 and C_3 are defined below:

Environment	Comparison		
	C_1	C_2	C_3
AAA	+1	+1	+1
AAB	-1	+1	-1
AAC	+1	-1	-1
ABC	-1	-1	+1

If increased environmental heterogeneity leads to an increased level of genic heterozygosity and this relationship is linear, then both C_1 and C_2 should be statistically significant and C_3 should not. If only one food medium affects the level of heterozygosity, for example, by maintaining larger population sizes and thus higher levels of heterozygosity, than either C_1 or C_2 may be significant but C_3 will not. If differences in the level of heterozygosity are associated with interactions between the types of food medium then the C_3 item may reach statistical significance.

In the analysis shows in table 9 it was never the case that both C_1 and C_2 were significant, as would be the case if environmental heterogeneity *per se* led to increased genic heterozygosity. The loci item is highly significant in all analyses but this just indicates that there are differences in heterozygosity between loci and is here of little interest. Additionally, the replicates item is significant in all analyses when compared with the expected binomial variance; this is hardly surprising in view of the extent of replicate divergence

TABLE 9

Analysis of variance of angular transformed, proportion heterozygosity for the "Chateau Tahbilk" and "Groningen" populations on sampling occasions 1 and 2.

Item	"Chateau Tahbilk"							
	Occasion 1				Occasion 2			
	df.	M.S.	F	p	df.	M.S.	F	p
C1	1	52.72	0.73	0.399	1	147.91	1.36	0.282
C2	1	0.03	0.00	0.999	1	20.59	0.50	0.485
C3	1	316.93	10.75	0.003	1	43.65	1.06	0.311
Loci	7	178.89	6.07	<0.001	7	144.01	3.49	0.007
C1×Loci	7	72.64	2.46	0.039	7	109.05	2.64	0.028
C2×Loci	7	8.13	0.28	0.957	7	17.79	0.43	0.876
C3×Loci	7	28.10	0.95	0.483	7	52.11	1.26	0.301
Replicates	32	29.48	$\chi^2_{(32)} = 101.60$	<0.001	32	41.28	$\chi^2_{(32)} = 148.09$	<0.001
Binomial								
Variance	—	9.28			—	8.92		

Item	"Groningen"							
	Occasion 1				Occasion 2			
	df.	M.S.	F	p	df.	M.S.	F	p
C1	1	63.54	4.41	0.045	1	9.90	0.38	0.543
C2	1	7.44	0.51	0.481	1	6.28	0.06	0.815
C3	1	31.73	2.20	0.148	1	2.79	0.11	0.743
Loci	6	978.83	67.88	<0.001	6	646.85	24.57	<0.001
C1×Loci	6	10.16	0.70	0.652	6	17.74	0.68	0.667
C2×Loci	6	23.94	1.66	0.168	6	114.06	4.35	0.003
C3×Loci	6	12.41	0.86	0.536	6	23.79	0.91	0.502
Replicates	28	14.42	$\chi^2_{(28)} = 44.31$	0.026	28	26.22	$\chi^2_{(28)} = 74.27$	<0.001
Binomial								
Variance	—	9.11			—	8.79		

observed in previous analyses. The C_1 comparison is significant at the 5 per cent level in the "Groningen" populations on occasion 1, indicating a difference in the level of heterozygosity between those environments which contained fig medium and those which did not. Table 8 shows that those environments which contained fig media support a higher level of heterozygosity. This effect is transitory; on occasion 2 the only between environment item significant is the $C_2 \times \text{Loci}$ item (at the 1 per cent level). This indicates a difference in heterozygosity between environments containing potato medium and those which do not which is not consistent for all loci. In the "Chateau Tahbilk" series of populations, C_3 is highly significant on sampling occasion 1, a consequence of the reduced genic heterozygosity in environments with two types of food medium compared with environments with only one type or with three types. Again this effect is transitory, not being found on occasion 2, but the $C_1 \times \text{loci}$ item is significant at the 5 per cent level on both sampling occasions, indicating a difference between those environments which contain fig medium and those which do not, which is not consistent over all loci. Thus these data provide little evidence for a general and sustained effect of environmental heterogeneity on genic heterozygosity at the loci under observation; the effects observed were either not consistent over occasions or over all loci.

4. DISCUSSION

Despite the large amount of effort expended in the study of electrophoretically detectable polymorphisms there is as yet no consensus as to the extent to which they are influenced by selection. However, this is unsurprising in view of the large proportion of work concentrated upon surveys of the extent and distribution of electrophoretic variation in natural populations. The lack of information on the ecology and dynamics of the surveyed populations has meant sets of data so derived are usually compatible with either extreme neutralist or selectionist viewpoints. Thus a more fruitful approach to the problem is provided by the laboratory investigation of electrophoretic polymorphisms.

Laboratory investigation of electrophoretic polymorphism are of two main types. Firstly, for individual, amenable loci, the *in vitro* properties, such as catalytic efficiency and thermostability, are determined. It may then be possible to relate these properties to the fitness of individual genotypes under selective conditions in the laboratory, or to the distribution of genotypes in the natural environment. This approach has met with some success in demonstrating that genotypic properties vary in ways which can be realistically related to environmental variation (de Jong and Scharloo, 1976; Hickey, 1977; van Delden, Boerema and Kamping, 1978; Kamping and van Delden, 1978). However, there has been less success in demonstrating that these polymorphisms are balanced in the natural environment.

In the long term, single locus studies will have to extend their scope to consider interactions between the locus under direct observation and associated loci (e.g., loci controlling expression), and between these loci and the rest of the genome and the environment. In the short term, it is necessary to broadly determine the limits of the response of genetic variation to natural selection. This need has led to the adoption of a second approach to the laboratory investigation of electrophoretic polymorphism which is to study the behaviour of several loci simultaneously in experimental populations. This approach, which is the one adopted in this study, has the additional advantage that it allows the investigation of general models of polymorphism maintenance, particularly those predicting associations between environmental heterogeneity and genic heterozygosity. The disadvantage of this approach is that it is not possible to ascribe selectively mediated changes at the loci under observation to selection acting directly on those loci. This disadvantage is outweighed, however, by the generality of the method, the observation of selection at a number of loci demonstrates that selection is of importance in effecting allele frequency changes over a large fraction of the genome.

The design of the experiment described here incorporates several advances over earlier studies. The main features of the study are:

- (i) The only environmental variation included in the design was spatial, and the range of environments encompassed three levels of spatial heterogeneity. The more heterogeneous environments contained discrete niches.
- (ii) In the main body of the experiment all environments were fully replicated, this facilitates comprehensive analysis of the data.
- (iii) The experiment was further replicated over two recently collected populations, which provides information on the generality of the results.

The outcome of this study was unexpected in some of its aspects. The results have re-affirmed that changes in allele frequency at electrophoretically detected loci may be mediated by selective forces, as it has been possible to demonstrate, for most loci under study, that systematic changes are found which can only have been caused by selection. However, rapid changes in allele frequency have occurred which are not shared by replicates, and, with the population size maintained in the experimental environments during the course of the study, these changes are unlikely to be due to random genetic drift. Nonetheless, random processes may play an important role in the initiation of such differences. The replicate divergence could come about through an accumulation of random environmental differences between replicates which initiate small genetic differences which are then magnified by selection. Alternatively, genetic differences between replicates for rare, but selectively important, alleles could have been created at the initiation of the populations. However, this seems unlikely to be the cause of much divergence as the rare alleles (either cryptic alleles, within an electrophoretic mobility class, at the loci under observation, or, alleles at a locus closely linked to the loci under observation) would have to be under intense selection to produce replicate differences. A more interesting possibility is that the loci under observation are part of a small segment of chromosome on which alleles at several loci are being selected in concert. Individual combinations of alleles will all be relatively rare, and the combination or combinations of alleles of selective significance may have been sampled initially or created *de novo* by recombination in one but not the other replicates. Another interesting possibility is that small genetic founder differences between replicates or small initial environmental differences between replicates placed the replicates at different positions on the fitness surface, this resulting in the replicates diverging to different selective peaks, or taking different routes to the same peak. Any of these mechanisms suggests a surprising subtlety in the response of genetic variability to natural selection.

In the systematic effects of selection demonstrated by the analyses of phenotype frequency (table 2) the effect of occasions was more common than the effects of experimental environments. The manipulated experimental environments produce some phenotypic differentiation, although there is no evidence for a positive relationship between environmental heterogeneity and genic heterozygosity, but the overall laboratory environment has had a greater effect. In the natural environment mortality can be caused by many different factors such as temperature, humidity, predation and competition; the laboratory environment, however, places a large emphasis on just one component of mortality, competition. The majority of mortality in the population cages is due to larval competition for resources and it is perhaps this which causes the strong selection obviously imposed by the laboratory environment. The effect of occasions is thus perhaps particularly prominent because the initial populations had only recently been collected. Indeed, it may be the particular vigour of the selective forces imposed by the laboratory environment which has resulted in substantial genomic rearrangement and thus replicate divergence initiated by rare recombinational events.

There is no general similarity between the responses of individual loci in the "Chateau Tahbilk" and "Groningen" populations. Those loci which

respond to the manipulated experimental environments in one population do not do so in the other, and in no case is there evidence for a similar directional selection in the two populations. The difference in response may arise because there are cryptic genetic differences (one electromorph representing different alleles in the two populations), or differences in the background genotype or different linkage relationships in the two populations. It is not so easy to suggest why there is a difference in the overall amount of change, as demonstrated by the phenotype analyses (tables 4 and 5) and the analyses of allele frequency (tables 6 and 7) between the two populations, the "Chateau Tahbilk" derived population showing a greater amount of change than the "Groningen" derived population. This difference cannot be due to population size differences, or to differences in the amount of linkage disequilibrium created at the initiation of the experiment. Thus, the extent of the response in the two populations must reflect genomic differences between the initial two populations, and may be some measure of the difference between the environment of the natural population and that of the laboratory population.

Selection has been demonstrated to affect a number of the loci under observation, but its action may be transmitted to those loci in a number of ways. The effects may result from selection solely at the loci under observation (albeit mediated by the background genotype) from selection on the locus under observation as part of a group of associated linked loci, or from selection on a linked locus or loci in linkage disequilibrium with the locus under observation. In this latter case, as the initial population samples were fairly large, this linkage disequilibrium will have been sampled from the original populations. This raises the question of why linkage disequilibrium is present in the original population (a survey of the "Chateau Tahbilk" population shows that inversions are not a complication, at least in this population, those that were present at the end of this study were small and rare: Middleton, pers. comm.). Furthermore, in this case, the observed selection is a fair reflection of what might have happened in the natural population had it been exposed to a novel environment. Thus, even if the alleles under observation are themselves neutral, it may not be possible to describe their behaviour in the natural population by application of the neutral theory.

Acknowledgements. The Science and Engineering Research Council are thanked for a research studentship (C. S. Haley). Statistical calculations were carried out on the University of Birmingham ICL 1906A and DEC 2060 computers.

REFERENCES

- AYALA, F. J. 1972. Darwinian *versus* non-Darwinian evolution in natural populations of *Drosophila*. *Proc. Sixth Berkeley Symp. Math. Stat. Prob.*, Vol. V, 211-236.
- BARKER, J. S. F. 1962. The estimation of generation interval in experimental populations of *Drosophila*. *Genet. Res.*, 3, 388-404.
- CROW, J. F. AND MORTON, N. E. 1955. Measurement of gene frequency drift in small populations. *Evolution*, 9, 202-214.
- DELLEN, VAN W., BOEREMA, A. C. AND KAMPING, A. 1978. The alcohol dehydrogenase polymorphism in populations of *Drosophila melanogaster*. I. Selection in different environments. *Genetics*, 90, 161-191.
- DEMPTER, E. 1955. Maintenance of genetic heterogeneity. *Cold Spring Harbour Symp. Quant. Biol.*, 20, 25-32.

- FISHER, R. A. AND FORD, E. B. 1947. The spread of a gene in natural conditions in a colony of the moth *Panaxia dominula* L. *Heredity*, 1, 143-174.
- GILLESPIE, J. H. 1974. The role of environmental grain in the maintenance of genetic variation. *Amer. Nat.*, 108, 831-836.
- GILLESPIE, J. H. 1976. A general model to account for enzyme variation in natural populations. II. Characterisation of the fitness function. *Amer. Nat.*, 110, 809-821.
- GILLESPIE, J. H. AND LANGLEY, C. H. 1974. A general model to account for enzyme variation in natural populations. *Genetics*, 76, 837-848.
- HALDANE, J. B. S. AND JAYAKAR, S. D. 1963. Polymorphism due to selection of varied direction. *J. Genet.*, 58, 291-317.
- HICKEY, D. A. 1977. Selection for amylase allozymes in *Drosophila melanogaster*. *Evolution*, 31, 800-804.
- DE JONG, G. AND SCHARLOO, W., 1976. Environmental determination of selective significance or neutrality of amylase variants in *Drosophila melanogaster*. 2. The relationship between ADH activity and adult mortality. *Biochem. Genet.*, 16, 541-551.
- KAMPING, A. AND DELDEN, VAN, W., 1978. The alcohol dehydrogenase polymorphism in populations of *Drosophila melanogaster*. 2. The relationship between ADH activity and adult mortality. *Biochem. Genet.*, 16, 541-551.
- KULLBACK, S. 1959. *Information theory and statistics*. Wiley, New York.
- LEVENE, H. 1953. Genetic equilibrium when more than one ecological niche is available. *Amer. Nat.*, 87, 331-333.
- LEVINS, R. 1968. *Evolution in changing environments*. Princeton University Press, Princetown, New Jersey.
- LEVINS, R. AND MACARTHUR, R. 1966. The maintenance of genetic polymorphism in a spatially heterogeneous environment: variations on a theme by Howard Levene. *Amer. Nat.*, 100, 585-589.
- MAYNARD SMITH, J. 1966. Sympatric speciation. *Amer. Nat.*, 100, 637-650.
- MCDONALD, J. F. AND AYALA, F. J. 1974. Genetic response to environmental heterogeneity. *Nature*, 250, 572-574.
- MINAWA, A. AND BIRLEY, A. J. 1975. Genetical and environmental diversity in *Drosophila melanogaster*. *Nature*, 255, 702-704.
- MINAWA, A. AND BIRLEY, A. J. 1978. The genetical response to natural selection by varied environments. I. Short-term observations. *Heredity*, 40, 39-50.
- POULIK, M. D. 1957. Starch gel electrophoresis in a discontinuous system of buffers. *Nature*, 180, 1477.
- POWELL, J. R. 1971. Genetic polymorphism in varied environments. *Science*, 174, 1035-1036.
- STROBECK, C. 1975. Selection in a fine grained environment. *Amer. Nat.*, 109, 419-425.
- TEMPLETON, A. R. AND ROTHMAN, E. D. 1974. Evolution in heterogeneous environments. *Amer. Nat.*, 108, 409-428.
- WALLIS, B. B. AND FOX, A. S. 1968. Genetic and developmental relationships between two alkaline phosphatases in *Drosophila melanogaster*. *Biochem. Genet.*, 2, 141-158.