BOOK REVIEW

THE ARCTIC SKUA. Peter O'Donald. Cambridge University Press. 1983. Pp. xvi + 324. Price: £25.00 HB.

The sub-title is "a study of the ecology and evolution of a sea-bird", and this bird, *Stercorarius parasiticus*, is the most strikingly dimorphic species on the British breeding list. The main thrust of the book is to explain this dimorphism, primarily by a systematic study of the demography of the Fair Isle population. So there is indeed quite a lot of ecology, also a fair amount of behaviour, with a score of delightful line drawings by Robert Gillmor. The discussion of formal genetics and life-history includes photographs of eggs and chicks. About half the book deals with the evidence for natural and sexual selection, and models of these processes; what I would call population genetics. The bits on evolution, as I understand the term, are thin, with an unconvincing cladogram. There are five living species of skua, four in the arctic and two in the antarctic, and all except the bipolar species are dimorphic, pale (rather like a dark adult gull) or melanic (like a dark immature gull). The frequency of the two forms varies widely. In the arctic skua there is a cline from about 100 per cent pale in the high-arctic to predominantly dark, up to 95 per cent or so in the Baltic, in temperate populations.

This dimorphism, in *S. parasiticus* at least, is controlled by a single gene, but the genetic basis of the considerable variation in the plumage of the two classes, of the chick's plumage, and of other characters like variation (within a morph) of age at first breeding is largely unknown. In the melanics, the darker birds are mostly homozygotes, the intermediates mostly heterozygotes, and this match is sufficient to help the analysis. Even with only two alleles and essentially two phenotypes, explaining the persistence and variation of the polymorphism is complicated.

Fair Isle is between Orkney and Shetland, and this book is largely based on work done there between 1958 and 1961 and from 1973 to 1979. The colony seems to have been started in the 1920's, 15 pairs were recorded in 1948, and then there was a consistent rise in numbers to a peak of 138 pairs in 1977. The frequency of melanics has been between 75 and 80 per cent. Rates of increase can be calculated separately for pales, intermediates, and darks. However, as the frequency of the forms hasn't changed, the differences between these rates of increase needs to be discussed somewhat more fully than is done in this book. The tentative explanation for the dimorphism relies on three sets of processes. There is natural selection favouring pales, because they start breeding younger, by half a year on average. This is counteracted by melanics producing more fledged chicks, 1.3 on average, compared with 1.1 for pales. The second process is sexual selection which favours melanics. Male melanics, by female choice, breed four or five days earlier in each season than pale birds. The combination of these two sets gives a net disadvantage to heterozygotes, an unstable equilibrium, and predicts fixation of melanics on Fair Isle. The third process is migration from the North; increasing the frequency of pales. There is,

regrettably, no discussion of why pales should be favoured overall further North, nor the extent to which this triple explanation is sufficient to explain the features of the cline as a whole, even though it is clearly consistent with the high frequency of melanics throughout the Baltic.

The evidence for all these phenomena is set out clearly, with lots of chi-squares and maximum likelihood estimates. The book is well written, well produced, and with a good index which is helpful when the argument gets complicated. I would have liked better maps, no Yates' corrections, graphical explanations of the models, a better definition of generation time, rates of increase estimated by computer, and data for morph numbers by years at Fair Isle and Foula. But this is an enjoyable book, and as a clear exposition of the care that is needed in field population genetics, and the thought required to combine demography and selection, one to be warmly welcomed.

MARK WILLIAMSON

Department of Biology, University of York