Book Reviews

Mechanisms of morphological evolution. Wallace Arthur. John Wiley & Sons Ltd., Chichester, West Sussex. 1984. Pp. xv + 275. £19.00.

There is at present considerable controversy as to the ability of neo-Darwinism to explain the pattern of evolution described by palaeontologists. Several palaeontologists have claimed that the fossil record shows short periods of rapid morphological change associated with speciation interspersed with long periods of relative stasis; furthermore they claim that this pattern could not have been brought about by the processes studied by ecological and population geneticists and have argued that novel, largely unspecified, evolutionary processes involving developmental saltations must be evoked. In this book Wallace Arthur has tried to address these problems and to develop an expanded neo-Darwinism that would be acceptable to geneticists, but at the same time explain apparent saltations. He restricts himself to the evolution of morphological characters because he believes that they are in some fundamental way different from molecular characters because of the developmental processes giving rise to them.

The first half of the book looks at microevolution, and considers selection on polygenic characters, the evolution of isolating barriers and character displacement. These are the processes which he believes to be involved in generating interspecific differences. Drift as an agent of genetic differentiation is given short shrift for two reasons: firstly because morphological differences are unlikely to be neutral, and secondly because, since morphology is generally polygenically controlled, drift at individual polygenes is unlikely to lead to much differentiation. Both assertions are open to argument; but it does give him the opportunity to omit completely the rather sterile debate and its concomitant mathematics between selectionists and neutralists. The first half is sound and unlikely to raise much blood pressure. I thought the treatment of heritability, though correct, unlikely to bring enlightenment to those who did not already understand it; and I found surprising, in view of some of the models developed later, his treatment of speciation, where he is very dismissive of non-allopatric models.

The second half is devoted to a model of how saltational evolution might occur. He believes this may have a role in megaevolution, while microevolution is sufficient to explain macroevolution. I am not convinced that the distinctions are all that useful, since I do not see how you measure the extent of any observed differences between taxa in the units that evolution itself works on. Arthur's model of saltation depends essentially on a novel form of selection acting on develop-

mental genes of large effect. He argues that morphological characters are produced by a developmental pathway, and that any model of morphological evolution must take account of that pathway. He postulates a bifurcating tree-like hierarchy of developmental (D) genes with changes in early acting genes having a much greater effect on final phenotype than late acting D genes, which are in effect conventional polygenes. He suggests that many major changes in body plan may have been produced by changes in early acting D genes. Because such a gene is likely to have manifold pleiotropic effects, it is very likely to be disadvantageous in conventional terms compared to wild type, and so would be removed by natural selection. To account for their occasional fixation, he postulates a new form of selection, *n*-selection (as opposed to *w*-selection). *n*-Selection operates where competitive pressures are absent so that the only constraint on a new form is that it should have a net reproductive rate of more than one. In order to achieve such uncompetitive conditions, the new mutant must have a sufficiently large effect on the phenotype that its possessors no longer compete for the same ecological resources as the normal type, and it must also give reproductive isolation. I found this argument unconvincing. The model seems in many ways similar to Maynard Smith's (1966) model of sympatric speciation through niche selection, and indeed would appear to be a special case of this model. Arthur is arguing that the mutation in the D locus produces individuals which are sufficiently different morphologically that they occupy a new niche: if this is so, then the selective coefficients are going to be sufficiently large that Maynard Smith's model will give a stable polymorphism. In other words, given Arthur's assumptions, standard *w*-selection models would appear competent to explain the spread of major mutants that shifted a species from one adaptive peak to the slopes of an adjacent peak. The question is not so much whether it could happen, but whether the spread of such large mutations does or has happened—and that is very difficult to obtain hard data on.

Arthur defines morphology in a rather strict way, so as to exclude, for instance, pattern. He also largely confines himself to animals, particularly flies and snails. He thus precludes discussion of a number of examples, for instance the evolution of mimicry in butterflies and of crop plants, which could have been very relevant to his basic thesis. The examples he does choose are interestingly discussed, though often of necessity inconclusive.

If my comments have so far been rather negative, this should not be taken to imply that the book is not of value. It has many good features, no least of which is the clarity of style. This book is a useful contribution

to the macroevolution debate, and deserves to be widely read.

REFERENCE

MAYNARD SMITH, J. (1966) Sympatric speciation. *Am. Nat.*, 100, 637–650.

MARK MACNAIR

*Department of Biological Sciences
University of Exeter*

Molecular Genetics of the Bacteria–Plant Interaction. Edited by A. Puhler. Springer. 1983. 393 pp. Price DM 114 (ca £30).

This book resulted from a symposium with the same title held in Germany in Autumn 1982. The papers are grouped in three sections: Rhizobium–plant interaction (22), Agrobacterium–plant interaction (13) and plant pathogenic bacteria (7). Some of the papers—in my opinion too few—are speculative. I especially enjoyed D. Tepfer's (Versailles) on the biology of genetic transformation of higher plants, which suggests that *A. rhizogenes* may play a role in evolution by providing a means for sexually isolated species to exchange genetic information. Tempé and Petit's (Orsay) account of "the opine trail" raises the interesting suggestion that a microcin producing strain of *A. radiobacter*, not itself a pathogen since it is unable to trigger opine synthesis, is a superparasite. It uses the opines induced by pathogenic Agrobacteria but controls their growth by its own microcin to which they are sensitive. The same paper also explores the speculation that Rhizobium may induce opines. In the welter of other short, technical papers I learned that free-living Rhizobia do not fix N₂ and grow at the same time (Ludwig, Santa Cruz) and that recent work on host plant genes involved in Rhizobium associations (Verma *et al.*, Montreal) shows that their functions are as poorly understood as other plant genes that govern the growth of plant pathogens. The section on pathogenic bacteria includes a first account of the molecular analysis of virulence genes (Staskawicz *et al.*, San Carlos) by using Tn5 induced mutagenesis, and a useful summary on cloning ice nucleation and siderophore genes (Orser *et al.*, Berkeley). These bacterial siderophores are fluorescent pigments that appear to enhance the growth of some plants and to control certain plant pathogenic microorganisms.

Although such volumes are ephemeral for the specialist, this one is especially valuable and useful as a source of recent information for people, like this reviewer, who are more concerned with the plant hosts than the bacteria.

P. R. DAY

*Plant Breeding Institute
Cambridge*

Evolutionary genetics of fishes. Edited by Bruce J. Turner. Plenum Press, New York. 1984. Pp. xviii + 636. \$85.00 (U.S.).

This volume consists of 12 chapters which are either reviews or extended papers covering a wide diversity of topics in the field of evolutionary genetics. All of the contributions are made by scientists from American research establishments.

The first and second chapters describe the evolutionary impact of polyploidy in two families of fish, the Salmonidea (Allendorf and Thorgaard) and the Catostomidea (Ferris). Polyploidy in these families has arisen in different ways (autopolyploidy and allopolyploidy) and the chapters describe the effects of rediploidization and gene silencing providing a useful comparison of the two polyploid events.

The following six chapters deal for the most part with the genetics of a single family, the Poeciliidae. This group of fishes is particularly interesting because it contains examples of naturally-occurring unisexual species and reproductively competent triploids. Kallman describes the mechanisms of sex determination in *Xiphophorus* species which involves the regulation of the male-determining gene by a closely-linked operator gene and by autosomal regulatory proteins. Molizot and Siciliano explore the linkage relationship of interspecific hybrids with the genus *Xiphophorus* in order to illuminate evolutionary processes by gene mapping. Borowsky provides copiously illustrated and highly detailed description of polymorphism in *Xiphophorus* and describes adaptations to "fine" and "coarse-grained" environmental conditions. Monaco, Rasch and Balsano present a fascinating account of the Amazon molly (*Poecilia formosa*) which describes the interdependence of the unisexual gynogenetic molly and the closely-related bisexual species. The evolutionary ecology of the genus *Poeciliopsis* investigated by Moore explains the adaptive value of parthenogenesis, the problems of coexistence between unisexual and bisexual members of the genus and the maintenance of clonal diversity. This latter subject is dealt with in more detail by Vrijenhoek in the final chapter on the Poeciliidae. He describes the effects of hybridogenesis on the evolution of *Poeciliopsis* species.

My initial concern that half of this volume dealt with a single family of fishes was dispelled by the realization of the importance of this group and the contribution it makes to the understanding of evolutionary genetics.

Bell's review of the genetics of the familiar three-spined stickleback (*Gasterosteus aculeatus*) and its related species presents a further dimension to the species which has been extensively used in behavioural research. He describes the effects of natural selection on several morphological features found in this species.

Williams and Koehn detail evidence which suggests the monospecific classification of the Atlantic eel. The biology, genetics and taxonomy of *Anguilla* species have been the subject of considerable controversy over many years. Their work is an important contribution but I