Original Article

Relationship of β_2 -Microglobulin to Arterial Stiffness in Japanese Subjects

Yasuaki SAIJO, Megumi UTSUGI, Eiji YOSHIOKA, Naoko HORIKAWA,
Tetsuro SATO, Yingyan GONG, and Reiko KISHI

β_2 -Microglobulin (β_2m) is related to inflammatory diseases, but there have been few reports of a relationship between β_2m and atherosclerosis. We have examined the influence of β_2m on brachial-ankle pulse wave velocity (baPWV) to clarify whether it is related to arterial stiffness. baPWV, β_2m , C-reactive protein (CRP), and conventional risk factors were measured in 614 males and 158 females. The adjusted means of baPWV were compared with the quartiles of β_2m , and significant differences in baPWV were observed across the quartiles of β_2m ($p=0.037$). After being adjusted for potential confounders, quartile 4 of β_2m , quartile 4 of CRP, and the combination of high β_2m plus high CRP were significantly associated with a high value of PWV (quartile 4 of β_2m : odds ratio [OR] 2.53, 95% confidence interval [CI], 1.31–4.89; quartile 4 of CRP: OR 2.27, 95% CI, 1.18–4.34; high β_2m plus high CRP: OR 5.60, 95% CI, 2.38–13.2). These results suggest that β_2m is associated with an increase of arterial stiffness. Further studies are needed to clarify whether β_2m is related to atherosclerotic diseases, and whether the combination of β_2m and CRP measurement is a useful predictor for the development of atherosclerosis. (*Hypertens Res* 2005; 28: 505–511)

Key Words: β_2 -microglobulin, C-reactive protein, glomerular filtration rate, pulse wave velocity, arterial stiffness

Introduction

Atherosclerosis is now generally accepted to be an inflammatory disorder in the arterial wall (1), and the C-reactive protein (CRP) level is a strong predictor of cardiovascular events (2–5). Meanwhile, it has been reported that β_2 -microglobulin (β_2m) is related to inflammatory diseases (6) and β_2m is now widely used in evaluation of many clinical conditions, such as dialysis-related amyloidosis (7), human immunodeficiency virus (HIV) disease (8), myeloma (9), leukemia (10), and collagen disease (11), for the estimation of the glomerular filtration rate (GFR) (12), and so on. However, there have been few

reports of a relationship between β_2m and atherosclerosis.

Pulse wave velocity (PWV) is known to be an indicator of arterial stiffness (13, 14), and there have been many reports on PWV and the development of atherosclerotic diseases (15–17). A simple noninvasive method for automatic measurement of brachial-ankle PWV (baPWV) has recently been developed. The technical simplicity and short sampling time of the new method make it more feasible for screening a large population than previous methods such as carotid-femoral PWV.

In this study, we have investigated the influences of β_2m on arterial stiffness to clarify whether β_2m is related to early stage atherosclerosis.

From the Department of Public Health, Hokkaido University Graduate School of Medicine, Sapporo, Japan.

This work was supported in part by a Grant-in-Aid for Young Scientists from the Ministry of Education, Culture, Sports, Science and Technology of Japan and a Grant-in-Aid for Scientific Research from the Ministry of Health, Labour and Welfare of Japan.

Address for Reprints: Yasuaki Saijo, M.D., Department of Public Health, Hokkaido University Graduate School of Medicine, Kita 15, Nishi 7, Kita-ku, Sapporo 060–8638, Japan. E-mail: y-saijo@med.hokudai.ac.jp

Received January 27, 2005; Accepted in revised form April 19, 2005.

Table 1. Characteristics in Quartiles of β_2 -Microglobulin

	β_2 -Microglobulin category				<i>p</i> -value
	Quartile 1 (<i>n</i> =223)	Quartile 2 (<i>n</i> =226)	Quartile 3 (<i>n</i> =185)	Quartile 4 (<i>n</i> =128)	
β_2 -Microglobulin range (mg/dl)	0.9–1.3	1.4–1.5	1.6–1.7	1.8–3.4	
Gender (male, %)	67.4	82.3	85.4	88.3	<0.00001
Age (years)	48.6±6.3	50.1±6.0	50.4±6.3	51.2±5.5	<0.0001
BMI (kg/m ²)	23.3±3.1	23.5±2.9	24.0±2.9	23.6±2.9	0.14
SBP (mmHg)	119.2±12.5	119.7±15.4	124.2±16.8	123.2±16.3	<0.01
DBP (mmHg)	75.2±11.0	75.6±10.9	78.3±11.6	77.5±11.3	<0.01
Heart rate (bpm)	60.7±9.5	61.1±10.0	61.7±9.7	61.3±9.2	0.77
Total cholesterol (mg/dl)	209.4±33.2	209.3±31.6	205.9±30.8	203.1±32.6	0.23
Triglycerides (mg/dl)	93 (62–146)	102 (76–145)	101 (73–164)	105 (75–147)	0.22
HDL-C (mg/dl)	59.9±15.0	57.4±14.1	55.6±14.4	53.2±13.2	<0.001
Fasting glucose (mg/dl)	97.3±22.6	98.8±25.0	95.9±13.3	95.2±15.1	0.33
Uric acid (mg/dl)	5.3±1.3	5.6±1.2	5.9±1.2	6.0±1.2	<0.00001
CRP (mg/dl)	0.035 (0.018–0.065)	0.034 (0.020–0.077)	0.046 (0.025–0.083)	0.055 (0.028–0.125)	<0.001
Estimated GFR (ml/min per 1.73 m ²)	122.3±19.1	105.2±18.6	102.7±19.4	97.8±17.0	<0.00001
Current smoker (%)	39.9	36.7	38.9	39.6	0.46
Drinker (%)	69.1	72.1	73.0	65.6	0.48
Frequency of exercise (%)					
Rarely or never	59.2	48.2	56.8	60.9	0.05
≥1 week	40.8	51.8	43.2	39.1	
Educational attainment (%)					
High school education or less	52.8	46.5	45.9	51.6	0.4
More than high school education	47.2	53.5	54.1	48.4	
Medication for					
Hypertension (%)	5.1	5.3	13.0	15.6	<0.0001
Hyperlipidemia (%)	6.9	5.7	5.9	2.3	0.34
Diabetes (%)	0.9	1.8	1.6	0.8	0.76
PWV (cm/s)	1,314±177	1,347±204	1,365±196	1,407±213	<0.001

Variables are presented as mean±SD, median (interquartile range) for skewed variables, or percentage. BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; HDL-C, high density lipoprotein cholesterol; CRP, C-reactive protein; GFR, glomerular filtration rate; PWV, pulse wave velocity.

Methods

Subjects

The subjects were local government employees (8,229 men and 2,194 women) aged 35 years or more who had their annual health checkup during the period from April 2003 through March 2004. We used a self-administered questionnaire including items on clinical history, family history, smoking, alcohol consumption, educational status, frequency of exercise, menopausal status, and hormone-replacement therapy. The questionnaire was distributed to the subjects in advance of their annual health checkup, and was collected at the checkup. Answers to the questionnaire and written informed consent to view health checkup data were obtained

from 3,907 men and 1,044 women (response rate: men 47.5%, women 47.6%). A total of 685 subjects (495 men, 190 women) were excluded for the following reasons: past history of coronary disease or stroke (*n*=136; 124 men, 12 women), low ankle/brachial pressure index (ABI; <0.9, *n*=12; 11 men, 1 woman), PWV not measured (*n*= 600; 416 men, 184 women), or blood samples not measured (*n*=3; 3 women). Among this original study group consisting of 3,412 male and 854 female subjects, we analyzed 614 male and 158 female subjects who requested optional examinations, including measurement of the serum β_2 m level.

This study was conducted with all the subjects' written informed consent and approved by the institutional ethical board for epidemiological studies of Hokkaido University Graduate School of Medicine.

Data Collection

Subjects were classified as either current smokers or non-smokers, with the latter group including both never- and ex-smokers. Drinkers were defined as those who consumed alcohol once a week or more. With regard to leisure-time exercise (with perspiration), subjects were categorized as exercising “rarely or never,” or “ ≥ 1 per week.” Finally, two groups were used to categorize subjects according to their educational attainment: “high school education or less” and “more than high school education.”

Anthropometric measures (height, body weight, and waist and hip circumferences) were recorded by a standardized protocol. The body mass index (BMI) was calculated as weight (kg)/height² (m²).

Blood samples were drawn from the antecubital vein of the seated subject with minimal tourniquet use after a 12-h fast. Specimens were collected in siliconized glass vacuum tubes containing sodium fluoride for blood glucose, and no additives for serum.

Total cholesterol (TC) levels were measured by an enzymatic method (Wako, Osaka, Japan). The triglyceride (TG) levels were measured by an enzymatic method (Daiichi Pure Chemicals, Tokyo, Japan), high density lipoprotein cholesterol (HDL-C) level by a direct method (Daiichi Pure Chemicals), uric acid (UA) by an enzymatic method (Daiichi Pure Chemicals), creatinine by an enzymatic method (KANTO KAGAKU, Tokyo, Japan), blood glucose levels by an amperometric method (ARKRAY, Kyoto, Japan), and β_2m by a latex immunoassay (Eiken Chemical, Tokyo, Japan).

The CRP levels were measured by nephelometry, with a latex particle-enhanced immunoassay (N Latex CRP II; Dade Behring, Tokyo, Japan). The assay could detect 0.004 mg/dl of CRP. Undetectable CRP values were recorded as 0.002 mg/dl.

All blood variables except for CRP were measured at Daiichi Clinical Laboratories, Inc. (Sapporo, Japan), a commercial hematology laboratory, where the measurements of TC and HDL-C were all standardized by the Lipid Standardized Program of the Centers for the Disease Control and Prevention (Atlanta, USA). CRP was measured at Mitsubishi Kagaku Bio-Clinical Laboratories, Inc. (Tokyo, Japan), a commercial hematology laboratory.

The estimated GFR was calculated using the Cockcroft-Gault formula (18) adjusted for body surface area (BSA) as follows:

$$\text{Estimated GFR} = (140 - \text{age}) / \text{Scr} \times \text{weight} / 72 \times 1.73 / \text{BSA},$$

where Scr is the serum creatinine concentration (mg/dl) and weight is measured in kg. In females, a correlation factor (0.85) was used. BSA was estimated using the DuBois formula (19).

baPWV was measured using a volume-plethysmographic apparatus (Form PWV/AVI; model BP-203RPEII, Colin Co., Komaki, Japan). Details about this instrument and its use

have been described elsewhere (20–23). The subjects were examined in the supine position. This device records the phonocardiogram, electrocardiogram, and volume pulse form and arterial blood pressure at both the left and right brachia and ankles.

Blood pressure, heart rate (HR), and ABI were measured using the pulse-wave velocimeter at the same time that PWV was measured. ABI was the ratio of ankle systolic blood pressure (SBP) to brachial SBP, and the right and left ABIs were measured simultaneously. In all the studies, baPWV was obtained after an at least 5-min rest.

Statistical Analysis

The subjects were categorized according to quartiles of β_2m values. The data are presented as the mean $\pm$ SD, the median (and interquartile range) for variables with a skewed distribution, or percentages, and analysis of variance (ANOVA), the Kruskal-Wallis test, or the χ^2 -test was used to compare data for these groups. The adjusted mean of PWV was compared among the quartiles of β_2m , with analysis of covariance (ANCOVA) with age, gender, BMI, SBP, HR, TC, HDL-C, blood glucose, log TG, UA, estimated GFR, log CRP, smoking status (smoker/nonsmoker), alcohol consumption (drinker/rarely or never), frequency of exercise (≥ 1 per week/rarely or never), educational attainment (high school education or less/more than high school education), medication for hypertension, medication for hyperlipidemia, and medication for diabetes. Logistic regression analyses were used to evaluate whether quartiles of β_2m and CRP were related to a high value of PWV (tertile three). As the next step, combined variables (low β_2m [≤ 1.7 mg/dl] plus low CRP [≤ 0.080 mg/dl]; low β_2m [≤ 1.7 mg/dl] plus high CRP [≥ 0.081 mg/dl]; high β_2m [≥ 1.8 mg/dl] plus low CRP [≤ 0.080 mg/dl]; and high β_2m [≥ 1.8 mg/dl] plus high CRP [≥ 0.081 mg/dl]) were created, and their association with the high value of PWV was evaluated. Odds ratios (OR) and 95% confidence intervals (95% CI) were calculated before and after adjustment for potential confounders. All of the above-mentioned potential confounders except log CRP were included in the multivariate logistic regression models as independent variables. To avoid multicollinearity, diastolic blood pressure (DBP) was not included in these models.

p-values < 0.05 were considered to be statistically significant. All analyses were conducted using the SPSS software package Version 12 for Windows (SPSS Inc., Chicago, USA).

Results

Characteristics of the groups in the β_2m category are shown in Table 1. Gender, age, SBP, DBP, HDL-C, UA, CRP, estimated GFR, medication for hypertension, and PWV were significantly different in the group in the β_2m category. Also, in crude regression analysis, β_2m was significantly associated

Fig. 1. Adjusted means of baPWV compared among quartiles of β_2 -microglobulin. Adjusted for age, gender, BMI, SBP, HR, TC, HDL-C, FBS, log TG, UA, log CRP, estimated GFR, smoking status, alcohol consumption, frequency of exercise, educational attainment, medication for hypertension, medication for hyperlipidemia, and medication for diabetes. *p value for difference (p for trend)

with age (Pearson's coefficient: 0.15; $p < 0.0001$).

Next, the adjusted means of baPWV were compared with the quartiles of β_2m (Fig. 1). Significant differences in baPWV were observed across the quartiles of β_2m ($p = 0.037$; p for trend = 0.069).

In unadjusted logistic regression analysis (Table 2), quartile 4 of β_2m (reference quartile 1 of β_2m), quartiles 2, 3 and 4 of CRP (reference quartile 1 of CRP), and the combinations of "high β_2m plus high CRP" and "high β_2m plus high CRP" (reference: low β_2m plus low CRP) were significantly associated with a high value of PWV. After being adjusted for age, BMI, SBP, HR, TC, HDL-C, log TG, UA, smoking status, alcohol consumption, frequency of exercise, educational attainment, medication for hypertension, medication for hyperlipidemia, and medication for diabetes, the associations with quartiles 2 and 3 of CRP disappeared, but quartile 4 of β_2m , quartile 4 of CRP, and the combinations of "high β_2m plus high CRP" and "high β_2m plus high CRP" were significantly associated with a high value of PWV (quartile 4 of β_2m : OR 2.53, 95% CI, 1.31–4.89; quartile 4 of CRP: OR 2.27, 95% CI, 1.18–4.34; high β_2m plus high CRP: OR 1.86, 95% CI, 1.18–2.95; high β_2m plus high CRP: OR 5.60, 95% CI, 2.38–13.2). These results were not substantially affected even if we used DBP as an independent variable instead of SBP.

Discussion

A significant relationship between CRP and PWV has been reported (24, 25), but, to the best of our knowledge, this is the first study to clarify the significant association between β_2m and PWV.

The end-stage renal disease (ESRD) population has increased arterial stiffness, and the PWV level is a strong independent predictor of all-cause and cardiovascular mortal-

ity (26). It has been reported that elevated PWV is significantly associated with reduced GFR (27), and that β_2m is a marker of GFR (12). Thus, GFR is a strong confounder in analyses of the association between β_2m and arterial stiffness, and our analyses were adjusted for estimated GFR. We speculate therefore that the inflammatory factor of β_2m is related to arterial stiffness.

In addition, we showed that the combination of high β_2m plus high CRP was significantly related to a high value of PWV with a higher OR (5.60). Since, in some inflammatory disorders, β_2m is regarded as necessary for, or as a discriminative marker of, inflammation (12–15, 28–30), this might indicate the inflammation that can not fully be estimated using only CRP.

β_2m has been identified as the light chain common to the HLA-A, -B, and -C major histocompatibility complex antigens, and is expressed on the surface of virtually all normal nucleated cells. The surfaces of lymphocytes and monocytes are particularly rich in β_2m , and lymphocytic synthesis and expression are further augmented by stimulation with mitogens or with interferons (31). Viral infections such as infectious mononucleosis, cytomegalovirus (CMV), and influenza A are associated with pronounced increases in the serum β_2m concentration (32). CMV-seropositive individuals have endothelial dysfunction and impaired responses to nitric oxide (33). Thus, chronic persistent viral infections may be related to the β_2m concentration and arterial stiffness.

Meanwhile, it has been reported that β_2m inhibits the growth of, and induces apoptosis or necrosis in tumor cells such as leukemia and myeloma cells (34, 35). Xie *et al.* suggested that it would be of interest to examine whether β_2m at high concentrations could also induce apoptosis or necrosis in normal cells, including endothelial cells and fibroblasts, because apoptotic or necrotic bodies and released enzymes and cytokines could act as chemoattractants for mononuclear cells, and they speculated that β_2m may be a potential initiator of the inflammatory response (36).

Diets and exercise inducing weight loss lower the CRP level (37, 38), and smoking and alcohol consumption are related to the CRP level (39, 40). Exercise induces an increase in the rate of β_2m excretion into the urine (41). But the relationships between β_2m , diet, and lifestyle have not been fully investigated. It is therefore necessary to elucidate the influences of diet and lifestyle on β_2m .

The present study has several limitations. First, this study could not identify a causal role for β_2m in the pathogenesis of arterial stiffness. Second, we measured only estimated GFR, using the Cockcroft-Gault formula. Since the direct assessment of GFR is rather complicated, we believe that estimated GFR is sufficient for a large population study. Third, only 4,951 of the 10,423 subjects that participated in the original study completed the questionnaire required for participation in this study. Since all of the present subjects requested optional examinations at their annual health checkup, they might have been more worried about their health than the

Table 2. Association of β_2 -Microglobulin (β_2 m), C-Reactive Protein (CRP) Levels, and Their Combination with High Values of Pulse Wave Velocity (Tertile 3 Category)

Parameter	Unadjusted OR (95% CI)	p-value	Adjusted OR* (95% CI)	p-value
β_2 m				
Quartile 1 (0.9–1.3 mg/dl)	reference		reference	
Quartile 2 (1.4–1.5 mg/dl)	1.50 (0.99–2.26)	0.06	1.45 (0.82–2.57)	0.20
Quartile 3 (1.6–1.7 mg/dl)	1.30 (0.84–2.01)	0.24	0.73 (0.40–1.36)	0.32
Quartile 4 (1.8–3.4 mg/dl)	2.40 (1.51–3.83)	<0.001	2.53 (1.31–4.89)	<0.01
CRP				
Quartile 1 (<0.004–0.021 mg/dl)	reference		reference	
Quartile 2 (0.022–0.040 mg/dl)	1.81 (1.14–2.89)	<0.05	1.39 (0.74–2.61)	0.30
Quartile 3 (0.041–0.080 mg/dl)	2.03 (1.28–3.23)	<0.01	1.24 (0.65–2.39)	0.52
Quartile 4 (0.081–8.36 mg/dl)	3.14 (2.00–4.96)	<0.00001	2.27 (1.18–4.34)	<0.05
Combination				
Low β_2 m (≤ 1.7 mg/dl) plus low CRP (≤ 0.080 mg/dl)	reference		reference	
High β_2 m (≥ 1.8 mg/dl) or High CRP (≥ 0.081 mg/dl)	1.78 (1.28–2.49)	<0.001	1.86 (1.18–2.95)	<0.01
High β_2 m (≥ 1.8 mg/dl) plus high CRP (≥ 0.081 mg/dl)	4.86 (2.54–8.91)	<0.00001	5.60 (2.38–13.2)	<0.0001

OR, odds ratio; CI, confidence interval. *Adjusted for age, gender, body mass index, systolic blood pressure, heart rate, total cholesterol, high density lipoprotein cholesterol, fasting blood glucose, log triglycerides, uric acid, estimated glomerular filtration rate, smoking status, alcohol consumption, frequency of exercise, educational attainment, medication for hypertension, medication for hyperlipidemia, and medication for diabetes.

general population. The age of subjects who requested the optional examinations was significantly higher than that of subjects who did not request the optional examinations (50 years vs. 48 years). And the baPWV of subjects who requested the optional examinations was higher than that of subjects who did not request the optional examinations (1,351 cm/s vs. 1,343 cm/s), but the difference was not significant. Thus, this study's subjects had slightly higher age and baPWV, but because the analyses were adjusted for many possible confounders, we believe that β_2 m was actually related to the high value of PWV. Fourth, since the subjects requested the optional examinations at their annual health checkup, they might have been more worried about their health than the general population. But the subjects who had past histories of coronary disease, stroke, or low ankle/brachial pressure were excluded, and the analyses were adjusted for many possible confounders. Fifth, conventional methods of measurement of PWV are carotid femoral and heart-ankle PWV, and the significance of baPWV for the prediction of cardiovascular events has not been published. The carotid femoral and heart-ankle PWV mainly reflect a property of the aorta (elastic artery), but baPWV involves properties of both the aorta and lower limb arteries (muscular artery). However, the validity and reliability of baPWV have been reported (42). Yamashina *et al.* (20) reported that baPWV was significantly correlated with aortic PWV measured directly by a catheter pressure transducer ($n=41$, $r=0.87$, $p<0.01$); the coefficient of variation of interobserver reproducibility was 8.4% in their study, and that of intraobserver reproducibility was 10.0%. The path length was estimated from the height of each subject

based on the superficial measurements in the Japanese population, suggesting possible errors. However, use of the equation should not have seriously biased the reliability of the PWV measurements, because the Pearson's correlation coefficient between the estimated length and the actual surface measurement was higher than 0.9 (43). And baPWV can be measured noninvasively and automatically. Therefore, we believe that baPWV is useful for population-based studies. Sixth, the sample size was relatively small. The lack of a significant relationship between quartile 3 of β_2 m and a high value of PWV would seem to have been due to the small sample size. In addition, when the logistic regression analyses were performed separately for men and women, the OR of men were consistently significant. However, the OR of women were not significant, even though the OR of women were similar to those of men. Finally, we could not obtain data on the subjects' income, although all the subjects worked for one local government. We therefore believe that the subjects were socioeconomically similar, and our data were adjusted for educational attainment, so it was considered that the influence of socioeconomic status on the adjusted analysis was practically nil.

In summary, our results suggest that β_2 m is associated with an increase of arterial stiffness. Because β_2 m is measured easily and is in widespread use, further studies are needed to clarify whether β_2 m is related to atherosclerotic diseases, and to elucidate whether the combination of β_2 m and CRP measurement is a useful predictive strategy for the development of atherosclerosis.

Acknowledgements

We thank Mr. Manabu Shojiguchi, Mr. Hiyoruki Arizuka, Ms. Toyoko Enomoto, Mr. Takanori Mogi, Mr. Naoto Sasaki, Mr. Takeshi Tsuda, Ms. Tomoko Arihara, Dr. Toshiyuki Hayashi, Ms. Chizuko Sato, and Dr. Takehito Nakabayashi for their excellent assistance with the data collection, and Ms. Akemi Onodera, Ms. Maki Fukushima, and Ms. Aki Yasuike for their assistance with the baPWV measurement.

References

- Ross R: Atherosclerosis—an inflammatory disease. *N Engl J Med* 1999; **340**: 115–126.
- Ridker PM, Hennekens CH, Buring JE, Rifai N: C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med* 2000; **342**: 836–843.
- Koenig W, Sund M, Frohlich M, *et al*: C-reactive protein, a sensitive marker of inflammation, predicts future risk of coronary heart disease in initially healthy middle-aged men: results from the MONICA (Monitoring Trends and Determinants in Cardiovascular Disease) Augsburg Cohort Study, 1984 to 1992. *Circulation* 1999; **99**: 237–242.
- Danesh J, Whincup P, Walker M, *et al*: Low grade inflammation and coronary heart disease: prospective study and updated meta-analyses. *BMJ* 2000; **321**: 199–204.
- Ridker PM, Rifai N, Rose L, Buring JE, Cook NR: Comparison of C-reactive protein and low-density lipoprotein cholesterol levels in the prediction of first cardiovascular events. *N Engl J Med* 2002; **347**: 1557–1565.
- Betha M, Forman DT: Beta 2-microglobulin: its significance and clinical usefulness. *Ann Clin Lab Sci* 1990; **20**: 163–168.
- Winchester JF, Salsberg JA, Levin NW: Beta-2 microglobulin in ESRD: an in-depth review. *Adv Ren Replace Ther* 2003; **10**: 279–309.
- Ullum H, Lepri AC, Katzenstein TL, *et al*: Prognostic value of single measurements of beta-2-microglobulin, immunoglobulin A in HIV disease after controlling for CD4 lymphocyte counts and plasma HIV RNA levels. *Scand J Infect Dis* 2000; **32**: 371–376.
- Diem H, Fateh-Moghadam A, Lamerz R: Prognostic factors in multiple myeloma: role of beta 2-microglobulin and thymidine kinase. *Clin Investig* 1993; **71**: 918–923.
- Sadamori N, Mine M, Hakariya S, *et al*: Clinical significance of beta 2-microglobulin in serum of adult T cell leukemia. *Leukemia* 1995; **9**: 594–597.
- Castro J, Jimenez-Alonso J, Sabio JM, *et al*, Grupo Lupus Virgen de las Nieves: Salivary and serum beta2-microglobulin and gamma-glutamyl-transferase in patients with primary Sjogren syndrome and Sjogren syndrome secondary to systemic lupus erythematosus. *Clin Chim Acta* 2003; **334**: 225–231.
- Jovanovic D, Krstivojevic P, Obradovic I, Durdevic V, Dukanovic L: Serum cystatin C and beta2-microglobulin as markers of glomerular filtration rate. *Ren Fail* 2003; **25**: 123–133.
- Lehmann ED: Clinical value of aortic pulse-wave velocity measurement. *Lancet* 1999; **354**: 528–529.
- Asmar R, Benetos A, Topouchian J, *et al*: Assessment of arterial distensibility by automatic pulse wave velocity measurement. Validation and clinical application studies. *Hypertension* 1995; **26**: 485–490.
- Laurent S, Katsahian S, Fassot C, *et al*: Aortic stiffness is an independent predictor of fatal stroke in essential hypertension. *Stroke* 2003; **34**: 1203–1206.
- Laurent S, Boutouyrie P, Asmar R, *et al*: Aortic stiffness is an independent predictor of all-cause and cardiovascular mortality in hypertensive patients. *Hypertension* 2001; **37**: 1236–1241.
- Guerin AP, Blacher J, Pannier B, Marchais SJ, Safar ME, London GM: Impact of aortic stiffness attenuation on survival of patients in end-stage renal failure. *Circulation* 2001; **103**: 987–992.
- Cockcroft DW, Gault MH: Prediction of creatinine clearance from serum creatinine. *Nephron* 1976; **16**: 31–41.
- DuBois D, DuBois EF: A formula to estimate the approximate surface area if height and weight be known. *Arch Intern Med* 1916; **17**: 863–871.
- Yamashina A, Tomiyama H, Takeda K, *et al*: Validity, reproducibility, and clinical significance of noninvasive brachial-ankle pulse wave velocity measurement. *Hypertens Res* 2002; **25**: 359–364.
- Yamashina A, Tomiyama H, Arai T, *et al*: Brachial-ankle pulse wave velocity as a marker of atherosclerotic vascular damage and cardiovascular risk. *Hypertens Res* 2003; **26**: 615–622.
- Tomiyama H, Yamashina A, Arai T, *et al*: Influences of age and gender on results of noninvasive brachial-ankle pulse wave velocity measurement—a survey of 12517 subjects. *Atherosclerosis* 2003; **166**: 303–309.
- Saijo Y, Utsugi M, Yoshioka E, *et al*: Relationship of *Helicobacter pylori* infection to arterial stiffness in Japanese subjects. *Hypertens Res* 2005; **28**: 283–292.
- Okamura T, Moriyama Y, Kadowaki T, Kanda H, Ueshima H: Non-invasive measurement of brachial-ankle pulse wave velocity is associated with serum C-reactive protein but not with α -tocopherol in Japanese middle-aged male workers. *Hypertens Res* 2004; **27**: 173–180.
- Yasmin, McEniery CM, Wallace S, Mackenzie IS, Cockcroft JR, Wilkinson IB: C-reactive protein is associated with arterial stiffness in apparently healthy individuals. *Arterioscler Thromb Vasc Biol* 2004; **24**: 969–974.
- Blacher J, Guerin AP, Pannier B, Marchais SJ, Safar ME, London GM: Impact of aortic stiffness on survival in end-stage renal disease. *Circulation* 1999; **99**: 2434–2439.
- Mourad JJ, Pannier B, Blacher J, *et al*: Creatinine clearance, pulse wave velocity, carotid compliance and essential hypertension. *Kidney Int* 2001; **59**: 1834–1841.
- Chiou YY, Chiu NT, Chen MJ, Cheng HL: Role of beta 2-microglobulinuria and microalbuminuria in pediatric febrile urinary tract infection. *Acta Paediatr Taiwan* 2001; **42**: 84–89.
- Vraetz T, Ittel TH, van Mackelenbergh MG, Heinrich PC, Sieberth HG, Graeve L: Regulation of beta2-microglobulin expression in different human cell lines by proinflammatory cytokines. *Nephrol Dial Transplant* 1999; **14**: 2137–2143.
- Mogi M, Otogoto J, Ota N, Inagaki H, Minami M, Kojima

- K: Interleukin 1 beta, interleukin 6, beta 2-microglobulin, and transforming growth factor-alpha in gingival crevicular fluid from human periodontal disease. *Arch Oral Biol* 1999; **44**: 535–539.
31. Azocar J, Essex M, Watson A, Gazit E, Anderson D, Yunis EJ: Changes in the expression of HLA and beta 2-microglobulin by cultured lymphoid cells. *Hum Immunol* 1982; **5**: 283–293.
 32. Cooper EH, Forbes MA, Hambling MH: Serum beta 2-microglobulin and C reactive protein concentrations in viral infections. *J Clin Pathol* 1984; **37**: 1140–1143.
 33. Grahame-Clarke C, Chan NN, Andrew D, et al: Human cytomegalovirus seropositivity is associated with impaired vascular function. *Circulation* 2003; **108**: 678–683.
 34. Mori M, Terui Y, Ikeda M, et al: Beta(2)-microglobulin identified as an apoptosis-inducing factor and its characterization. *Blood* 1999; **94**: 2744.
 35. Min R, Li Z, Epstein J, Barlogie B, Yi Q: Beta(2)-microglobulin as a negative growth regulator of myeloma cells. *Br J Haematol* 2002; **118**: 495–505.
 36. Xie J, Yi Q: Beta2-microglobulin as a potential initiator of inflammatory responses. *Trends Immunol* 2003; **24**: 228–229.
 37. Esposito K, Pontillo A, Di Palo C, et al: Effect of weight loss and lifestyle changes on vascular inflammatory markers in obese women: a randomized trial. *JAMA* 2003; **289**: 1799–1804.
 38. Okita K, Nishijima H, Murakami T, et al: Can exercise training with weight loss lower serum C-reactive protein levels? *Arterioscler Thromb Vasc Biol* 2004; **24**: 1868–1873.
 39. Saito M, Ishimitsu T, Minami J, Ono H, Ohnri M, Matsuoka H: Relations of plasma high-sensitivity C-reactive protein to traditional cardiovascular risk factors. *Atherosclerosis* 2003; **167**: 73–79.
 40. Albert MA, Glynn RJ, Ridker PM: Alcohol consumption and plasma concentration of C-reactive protein. *Circulation* 2003; **107**: 443–447.
 41. Poortmans JR, Blommaert E, Baptista M, De Broe ME, Nouwen EJ: Evidence of differential renal dysfunctions during exercise in men. *Eur J Appl Physiol Occup Physiol* 1997; **76**: 88–91.
 42. Munakata M, Ito N, Nunokawa T, Yoshinaga K: Utility of automated brachial ankle pulse wave velocity measurements in hypertensive patients. *Am J Hypertens* 2003; **16**: 653–657.
 43. Kobayashi K, Akishita M, Yu W, Hashimoto M, Ohni M, Toba K: Interrelationship between non-invasive measurements of atherosclerosis: flow-mediated dilation of brachial artery, carotid intima-media thickness and pulse wave velocity. *Atherosclerosis* 2004; **173**: 13–18.