

## COMMENTARY

# Normotension, blood pressure variability and early target organ damage

Luis M Ruilope

*Hypertension Research* (2011) 34, 1075; doi:10.1038/hr.2011.122; published online 28 July 2011

Arterial hypertension is an important public health challenge. Worldwide, nearly 8 million premature deaths, 54% of stroke cases and 47% of ischemic heart disease cases were attributable to high blood pressure (BP, >115 mm Hg systolic) in 2001. About half of these cases develop in people with arterial hypertension (>140/90 mm Hg); the remaining cases affect patients with BP levels that are normal or high normal.<sup>1</sup> Wide intra-individual variability in BP influences prognosis and contributes to the burden of cardiovascular (CV) disease. In fact, data collected over the past 50 years through ambulatory BP monitoring (ABPM)<sup>2</sup> have shown that BP is a highly variable parameter in both normotensive and hypertensive subjects.

The potential role of BP variability as an independent predictor of organ damage and CV events was initially described years ago,<sup>3,4</sup> but most recommendations related to hypertension management are still based on isolated clinic measurements, and BP variability is not recognized as an important prognostic parameter in recent guidelines.<sup>5</sup>

Recent data have increased interest in the prognostic significance of this parameter in the context of CV prognosis.<sup>6–9</sup> Some of these studies have investigated more refined estimates of BP variability based on ABPM.<sup>6,7</sup> However, the recent demonstration in hypertensive patients<sup>8</sup> and the general population<sup>9</sup> that variability in BP measurement occurring over days or months—obtained in clinical visits in the case of hypertensive patients<sup>8</sup> and in iterative BP estimations in the general

population followed in the NHANES study<sup>9</sup>—has greatly expanded the scope of investigations of the role of BP variability in CV prognosis.

In this issue of *Hypertension Research*, Schutte *et al.*<sup>10</sup> report interesting data that correlate short-term BP variability and electrocardiographic left ventricular mass in normotensive African patients, including both normo- and hypertensive subjects. As a measure of short-term reading-to-reading BP variability, the authors used averages of daytime and nighttime SDs weighted for the duration of the daytime and nighttime intervals, as proposed previously.<sup>5</sup> These data confirm the role of BP variability in normotensive subjects. The burden of disease attributable to high BP has already been established in normotension (BP <140/90 mm Hg).<sup>1</sup> The data of Schutte *et al.*<sup>10</sup> demonstrate that BP variability is involved in the development of target organ damage in normotensive blacks. African populations are known to have a two- to threefold greater prevalence of left ventricular hypertrophy,<sup>11</sup> which contributes more to the risk of CV mortality in African-Americans than in Caucasians.<sup>12</sup>

Interestingly, the same electrocardiographic findings were not obtained in Caucasian patients or in hypertensive African patients. In hypertensive Caucasians, an independent relationship between carotid cross-sectional wall area and 24-h systolic BP variability was found.

These data confirm a relationship between BP variability and target organ damage, but also that there are ethnic differences that require additional studies to clarify when the analysis and treatment of BP variability can be useful in clinical practice.

### CONFLICT OF INTEREST

The author declares no conflict of interest.

- 1 Lawes CMM, Hoorn SV, Rodgers A, for the International Society of Hypertension. Global burden of blood-pressure-related disease. *Lancet* 2008; **371**: 1513–1518.
- 2 Mancia G, Ferrari A, Gregorini L, Parati G, Pomidossi G, Bertinieri G, Grassi G, de Rienzo M, Pedotti A, Zanchetti A. Blood pressure and heart rate variability in normotensive and hypertensive human beings. *Circ Res* 1983; **53**: 96–104.
- 3 Parati G, Pomidossi G, Albini F, Lallaspina D, Mancia G. Relationship of 24-h blood pressure mean and variability to severity of target organ damage. *J Hypertens* 1987; **5**: 93–98.
- 4 Pringle E, Phillips C, Thijs L, Davidson C, Staessen JA, De Leeuw P, Jaaskivi M, Nachev C, Parati G, O'Brien ET, Tuomilehto J, Webster J, Bulpitt CJ, Fagard RH, on behalf of the Syst-Eur study investigators, On behalf of the Syst-Eurinvestigators. Systolic blood pressure variability as a risk factor for stroke and cardiovascular mortality in the elderly in the elderly hypertensive population. *J Hypertens* 2003; **21**: 2251–2257.
- 5 2007 Guidelines for the Management of Arterial Hypertension. The Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and the European Society of Cardiology (ESC). *J Hypertens* 2007; **25**: 1105–1187.
- 6 Mancia G. Prognostic value of long-term blood pressure variability. The evidence is growing. *Hypertension* 2011; **57**: 141–143.
- 7 Bilo G, Parati G. Rate of blood pressure changes assessed by 24 h ambulatory blood pressure monitoring: another meaningful index of blood pressure variability. *J Hypertens* 2011; **29**: 1054–1058.
- 8 Rothwell PM, Howard SC, Dolan E, O'Brien E, Dobson JE, Dahlof B, Sever PS, Poulter NR. Prognostic significance of visit-to-visit variability, maximal systolic blood pressure, and episodic hypertension. *Lancet* 2010; **375**: 895–905.
- 9 Muntner P, Shimbo D, Tonelli M, Reynolds K, Arnett DK, Oparil S. The relationship between visit-to-visit variability in systolic blood pressure and all-cause mortality in the general population: findings from III NHANES, 1988 to 1994. *Hypertension* 2011; **57**: 160–166.
- 10 Schutte AE, Schutte R, Huisman HW, van Rooyen JM, Fourie CMT, Malan NT, Malan L. Blood pressure variability is significantly associated with ECG Left ventricular mass in normotensive Africans: The SABPA Study. *Hypertens Res* 2011; **34**: 1127–1134.
- 11 Drazner MH, Dries DL, Peshock RM, Cooper RS, Klassen C, Kafi F, Willett D, Victor RG. Left ventricular hypertrophy is more prevalent in blacks than in whites in the general population: the Dallas Heart Study. *Hypertension* 2005; **46**: 124–129.
- 12 Havranek EP, Froshaug DB, Emserman CD, Hanratty R, Krantz MJ, Masoudi FA, Dickinson LM, Steiner JF. Left ventricular hypertrophy and cardiovascular mortality by race and ethnicity. *Am J Med* 2008; **121**: 870–875.