

## ORIGINAL ARTICLE

# The effects of subtherapeutic antibiotic use in farm animals on the proliferation and persistence of antibiotic resistance among soil bacteria

Sudeshna Ghosh and Timothy M LaPara

Department of Civil Engineering, University of Minnesota, Minneapolis, MN, USA

**The use of antibiotics at subtherapeutic concentrations for agricultural applications is believed to be an important factor in the proliferation of antibiotic-resistant bacteria. The goal of this study was to determine if the application of manure onto agricultural land would result in the proliferation of antibiotic resistance among soil bacteria. Chlortetracycline-resistant bacteria were enumerated and characterized from soils exposed to the manure of animals fed subtherapeutic concentrations of antibiotics and compared to the chlortetracycline-resistant bacteria from soils at farms with restricted antibiotic use (dairy farms) and from non-agricultural soils. No significant differences were observed at nine different study sites with respect to the numbers and types of cultivated chlortetracycline-resistant bacteria. Genes encoding for tetracycline resistance were rarely detected in the resistant bacteria from these sites. In contrast, soils collected from a tenth farm, which allowed manure to indiscriminately accumulate outside the animal pen, had significantly higher chlortetracycline-resistance levels. These resistant bacteria frequently harbored one of 14 different genes encoding for tetracycline resistance, many of which (especially *tet(A)* and *tet(L)*) were detected in numerous different bacterial species. Subsequent bacterial enumerations at this site, following the cessation of farming activity, suggested that this farm remained a hotspot for antibiotic resistance. In conclusion, we speculate that excessive application of animal manure leads to the spread of resistance to soil bacteria (potentially by lateral gene transfer), which then serve as persistent reservoir of antibiotic resistance.**

*The ISME Journal* (2007) 1, 191–203; doi:10.1038/ismej.2007.31; published online 24 May 2007

**Subject Category:** microbial population and community ecology

**Keywords:** antibiotic resistance; chlortetracycline; concentrated animal feeding operation; lateral gene transfer; subtherapeutic

## Introduction

Antibiotics play an important role in controlling infectious disease. Many antibiotics used in clinical practice are naturally occurring, such that resistance to antibiotics among bacteria predates their modern clinical use. The past few decades, however, have witnessed a steady increase in the number and diversity of antibiotic-resistant bacteria – rendering some bacterial infections virtually untreatable (Livermore, 2003). The cost of resistance associated with the treatment of infections caused by antibiotic-resistant bacteria has increased many-fold (Levy and Marshall, 2004). The pervasive use of

antibiotics for both therapeutic and non-therapeutic purposes has been blamed for the widespread occurrence of antibiotic-resistant bacteria, because the use of antibiotics creates a selective pressure in favor of resistant bacteria.

Agricultural use accounts for at least half of the antibiotics produced in the United States (Lipsitch *et al.*, 2002). Antibiotics are used in agriculture for the treatment of sick animals, disease prophylaxis, growth promotion and crop dusting. Growth promotion involves the addition of subtherapeutic doses of antibiotics to animal feed to induce faster weight gain, whereas prophylaxis involves antibiotic use to prevent disease outbreaks. There is considerable controversy regarding the necessity of such subtherapeutic antibiotic use compared to the threat it poses to human health (Ferber, 2003; Livermore, 2003; Arnold *et al.*, 2004; Phillips *et al.*, 2004). While it is agreed that subtherapeutic antibiotic use leads to an increase in antibiotic-resistant fecal bacteria in the animals (Levy, 1978; Aarestrup

---

Correspondence: Dr TM LaPara, Department of Civil Engineering, University of Minnesota, 500 Pillsbury Drive SE, Minneapolis, MN, 55455-0116, USA.

E-mail: lapar001@umn.edu

Received 14 February 2007; revised 27 March 2007; accepted 3 April 2007; published online 24 May 2007

*et al.*, 2001; Hayes *et al.*, 2004; Jackson *et al.*, 2004), the role of subtherapeutic antibiotic use in the global spread of antibiotic resistance remains ambiguous.

The disposal of animal waste could be a possible route for the spread of antibiotic resistance from animal farms. Animal waste generated at farms is typically applied to agricultural fields as a fertilizer; however, many animal farms are too large compared to the quantity of available land (United States Environment Protection Agency, 2004). This problem has become more pertinent over the last few decades because of a gradual transition from small farms to concentrated animal feeding operations (CAFOs). Thus, there is an increased risk of adding persistent antibiotic residues and resistant fecal bacteria to soil, potentially leading to the proliferation of resistance among indigenous bacteria (Chee-Sanford *et al.*, 2001; De Liguoro *et al.*, 2003; Sørnum *et al.*, 2006).

While numerous studies have focused on the antibiotic resistance and survival of specific commensal bacteria (for example, *Enterococci*) originating from farms using subtherapeutic concentrations of antibiotics in their animal feed (for examples, see Aarestrup *et al.*, 2001, 2002; Hayes *et al.*, 2004), few studies have investigated the proliferation of antibiotic resistance among diverse species of bacteria (for examples, see Aminov *et al.*, 2002; Sengelov *et al.*, 2003). In the present study, antibiotic-resistant bacteria were enumerated and characterized from soils fertilized with manure at farms using subtherapeutic concentrations of antibiotics in their animal feed. We hypothesized that the application of

manure from farms using subtherapeutic antibiotics on agricultural land would result in the proliferation of antibiotic resistance among the indigenous soil bacteria. As a basis of comparison, antibiotic-resistant bacteria were also enumerated and characterized from soils fertilized with manure from dairy farms (DFs) (where antibiotic use is more restricted) and from non-agricultural soils. Our results suggest that excessive application of animal manure leads to the spread of resistance to soil bacteria (potentially by lateral gene transfer), which then serve as persistent reservoir of antibiotic resistance.

## Methods

### *Study sites and sample collection*

Soil samples from 10 sites were studied, including four swine feeding operations (SFOs), three dairy farms and three non-agricultural areas (NAGs) (Table 1). Site SFO1 was a small swine farm, where animal waste was allowed to overflow the animal pen and accumulate. The other three SFOs were large CAFOs (>1000 animal units per year) housing pigs, where manure was stored in underground pits for 6–12 months before land application. At these sites, liquid manure was injected into agricultural fields approximately six inches below the soil surface, once every 2–4 years. Antibiotic use at the three DFs was restricted to veterinary treatment of diseased animals. Manure at the DFs was stored for up to a year before it was applied to the soil surface.

**Table 1** Description of the 10 sites from which soil samples were collected to determine antibiotic resistance levels

Site name	Location	Site type	No. of animal units/year	Subtherapeutic antibiotic use at SFO/therapeutic use at DF	Manure application time	Manure application rate gallons/acre	Sample collection time
SFO1	South-central Minnesota	Farrowing to finish pig farm	40	Chlortetracycline	NA	NA	Nov-03
SFO2	South-central Minnesota	Wean to finish pig farm	1200	Tylosin, bacitracin, chlortetracycline	Late fall-03	5000	Apr-04
SFO3	South-central Minnesota	Wean to finish pig farm	3000	Chlortetracycline, carbadox	Apr-04	Data not available	Aug-04
SFO4	South-central Minnesota	Wean to finish pig farm	2800	Tylosin, chlortetracycline	Nov-04	4500	Nov-04
DF1	South-central Minnesota	Dairy farm	77	Amoxicillin, penicillin, cephalosporin	Winter-04	3500	May-04
DF2	Wisconsin	Dairy farm	350	Penicillin, tetracycline, excenel	Spring-04	12000	Sep-04
DF3	Wisconsin	Dairy farm	196	Penicillin, streptomycin	May-04	8000	Nov-04
NAG1	Quetico provincial park, Canada	Non-agricultural					July-04
NAG2	BWCAW Minnesota	Non-agricultural					Aug-04
NAG3	University of Minnesota	Non-agricultural					Aug-04

Abbreviations: DF, dairy farm; NAG, non-agricultural; SFO, swine feeding operations.

One animal unit is defined as  $0.4 \times$  the number of swine over 25 kg or as  $1.4 \times$  the number of mature dairy cattle. Manure was not directly applied to soils at Site SFO1, rather it was allowed to overflow the animal pen as it accumulated.

The three NAGs were Quetico Provincial Park (ON, Canada), the Boundary Waters Canoe Area Wilderness (BWCAW; Minnesota, MN, USA) and the University of Minnesota (Minneapolis, MN, USA). The BWCAW and Quetico Provincial Park are undeveloped wilderness areas situated west of Lake Superior near the Canada–US border comprised of pristine forests and lakes that receive approximately 200 000 and 125 000 recreational visitors each year, respectively (Beymer, 2000; Ontario Parks, 2006).

Soil samples were collected from three different locations within 2 m of each other. Samples (~500 g) were collected from the soil surface to which manure had been applied in the case of the DFs and from depths of 1–6 inches below the manure treated soil surface at the SFOs except Site SFO1. At Site SFO1, soil samples were collected near (~20 m) the pen from the edge of an adjacent corn field where there was no visible manure accumulation. Soil samples at the NAGs were collected from grassland (University of Minnesota) and forested land (BWCAW and Quetico Provincial Park) immediately below the O-horizon. Soil samples were immediately transported to the University of Minnesota and processed within 24 h, except for the BWCAW and Quetico Provincial Park samples, which were processed within 72 h.

#### Enumeration of resistant soil bacteria

Bacteria were dislodged from soil by vortexing 0.5 g of soil (wet weight) suspended in 9 ml of phosphate-buffered saline (PBS, 10 mM, pH 7). Samples were then serially diluted 10-fold in PBS and applied to agar plates using standard spread plating techniques. Agar plates were amended with either 20 mg l<sup>-1</sup> of chlortetracycline, 20 mg l<sup>-1</sup> tylosin, or 40 mg l<sup>-1</sup> of carbadox to enumerate resistant bacteria. Antibiotic resistance levels were calculated as the ratio of bacteria growing on plates supplemented with antibiotics compared to the number of bacteria growing on plates without antibiotics. All initial plate counts were done in triplicate and averaged before calculating resistance levels for each soil sample. The average resistance levels for each of three soil samples collected at a site were subsequently used to calculate mean and s.d. of resistance levels at a site.

Two different nutrient media were used to enumerate bacteria. PYT80 agar plates contained 80 mg l<sup>-1</sup> (each) of peptone, tryptone and yeast extract as well as 400 mg l<sup>-1</sup> delvovid salt (DSM Food Specialties, Netherlands) and 15 g l<sup>-1</sup> granulated agar. Delvovid salt was added as a fungicide; it contained 50% natamycin (w/w). Luria–Bertani (LB) growth medium was composed of 10 g l<sup>-1</sup> tryptone, 5 g l<sup>-1</sup> yeast extract, 4.9 g l<sup>-1</sup> NaCl, 200 mg l<sup>-1</sup> delvovid salt and 15 g l<sup>-1</sup> granulated agar. PYT80 plates were incubated at 25°C until bacterial colonies were visible (3–5 days); LB plates were incubated for

1–2 days at 37°C. All growth media were sterilized by autoclaving (20 min; 121°C; 15 psig).

#### Identification of resistant bacterial isolates

Antibiotic-resistant colonies were randomly picked from plates containing 30–300 colonies. Bacteria collected from plates were regrown in liquid culture or collected directly from agar plates. Cells were suspended in 0.5 ml of lysis buffer (5% sodium dodecyl sulfate, 120 mM sodium phosphate, pH 8) followed by three freeze–thaw cycles and a 90 min incubation at 70°C to lyse the cells. Genomic DNA was then purified using a Fast DNA Spin Kit (MP Biomedicals LLC, Irvine, CA, USA).

Bacterial isolates were initially screened by ribosomal intergenic spacer analysis (RISA) to identify unique isolates. Polymerase chain reaction (PCR) was performed using a PTC 100 thermal cycler (MJ Research Inc., Watertown, MA, USA). The ribosomal intergenic spacer region was amplified using primers 1522F (5'-TGCGGCTGGATCCCCTCCTT-3') and 132R (5'-CCGGGTTTCCCCATTCCG-3') (Ranjard *et al.*, 2001). The final 50 µl reaction mixture contained: 1 × PCR buffer (Promega, Madison, WI, USA), 4 nmol deoxynucleoside triphosphates, 25 pmol forward and reverse primers, 1.25 units of *Taq* polymerase (Promega) and ~1 ng of template DNA. The PCR protocol included a 5 min initial denaturation at 95°C, followed by 35 cycles of 94°C for 1 min, 55°C for 1 min, and 72°C for 2 min and a final extension for 10 min at 72°C. Amplified products were resolved by electrophoresis on 2% (wt/vol) agarose gels stained with ethidium bromide.

PCR was used to amplify 16S rRNA gene fragments of isolates having distinct RISA patterns using primer set 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1522R (5'-AAGGAGGTGATCCANCCCA-3') or primer set 338F (5'-ACTCCTACGGGAGG CAGCAG-3') and 907R (5'-CCGTCAATTCCTTTRAGTTT-3') or primer set 338F and 518R (5'-ATTA CCGCGGCTGCTGCTGG-3') (Edwards *et al.*, 1989; Lane, 1991; Muyzer *et al.*, 1993). Different primer sets were used because some primer sets would generate either no PCR products or multiple products of different sizes. PCR mixtures and conditions were as described above. PCR products were purified using a GeneClean kit (MP Biomedicals) and sequenced using 338F and 907R or 338F and 518R as sequencing primers. Sequencing was performed at the Advanced Genetic Analysis Center at the University of Minnesota using an ABI 3100 Genetic Analyzer (Applied Biosystems, Foster City, CA, USA).

#### Multiplex PCR

Chlortetracycline-resistant isolates were examined by multiplex PCR for the presence of 14 different genes encoding for resistance to tetracycline (Ng

*et al.*, 2001). These genes confer resistance to tetracycline by three different mechanisms: efflux (*tet(A)*, *tet(B)*, *tet(C)*, *tet(D)*, *tet(E)*, *tet(G)*, *tet(K)*, *tet(L)*, *tetA(P)*); ribosomal protection (*tet(M)*, *tet(O)*, *tet(S)*, *tet(Q)*); enzymatic degradation (*tet(X)*) (Chopra and Roberts, 2001). Multiplex PCR was performed in four separate groups (Group I: *tet(B)*, *tet(C)*, and *tet(D)*; Group II: *tet(A)*, *tet(E)*, and *tet(G)*; Group III: *tet(K)*, *tet(L)*, *tet(M)*, *tet(O)* and *tet(S)*; and Group IV: *tetA(P)*, *tet(Q)* and *tet(X)*). PCR conditions included a 5 min initial denaturation at 94°C, followed by 35 cycles of 94°C for 1 min, 55°C for 1 min and 72°C for 1.5 min and a final extension for 10 min at 72°C. PCR products were separated on 2% (wt/vol) agarose gels stained with ethidium bromide. Isolates putatively containing genes encoding for tetracycline resistance were identified by comparison with positive controls (Ng *et al.*, 2001) that were simultaneously resolved on 2% (w/vol) agarose gels. Multiplex PCR product specificity was initially confirmed by performing PCR targeting a single gene encoding resistance to tetracycline. About 10% of these PCR products targeting a single gene were randomly selected for further confirmation by nucleotide sequence analysis. In addition, specific PCR products were subjected to nucleotide sequence analysis when genes encoding resistance to tetracycline were detected in an organism not previously known to harbor that gene (for example, *tet(A)* in a *Bacillus* sp).

#### Data analysis

Analysis of variance (ANOVA) was performed to determine the significance of the differences between antibiotic resistance levels at the 10 sites. Differences in chlortetracycline resistance levels at the sites were further examined by pairwise comparison of resistance levels using Tukey honest significant difference. This method used a stringent Type I error rate and used the Studentized range distribution to construct simultaneous confidence intervals for differences of all pairs of means. These statistical analyses were performed using MacAnova

software (Version of 02/05/03 Win32s (BCPP5.0), Department of Applied Statistics, University of Minnesota, MN, USA (<http://www.stat.umn.edu/macanova>).

Principal component analysis (PCA) was performed to evaluate differences between the 10 sites based on the presence or absence of the types of chlortetracycline-resistant bacteria (based on 16S rRNA gene sequences) and their relative abundance. Bacteria having different RISA patterns but at least 98% partial (~550 bp) 16S rRNA gene sequence identity were grouped together as a single type and their abundance was measured as the percentage of the total number of chlortetracycline-resistant isolates from a particular site on a particular growth medium. PCA was performed using NTSYSpc version 2.11S (Applied Biostatistics Inc., Setauket, NY, USA).

The 16S rRNA gene fragment sequences were compared with sequences obtained from the GenBank database using the BLASTn program (Benson *et al.*, 1999) to determine the phylogenetic affiliation of the isolate. Nucleotide sequences (partial 16S rRNA genes and genes encoding tetracycline resistance) have been deposited in the GenBank database under accession numbers DQ910178-DQ910203, DQ910205-DQ910237, DQ910239-DQ910317 and EF055265-EF055292.

## Results

#### Enumeration of resistant bacteria in soil

Bacteria resistant to tylosin and carbadox were enumerated on nutrient-poor (PYT80) agar plates from soils at nine different sites (Table 2). Mean resistance levels were highly variable for both of these compounds, ranging between 3–55% and 10–47%, respectively, of the plate count values on unamended growth media. However, ANOVA revealed that there were no statistically significant differences between the 10 sites based on tylosin ( $P > 0.15$ ) and carbadox ( $P > 0.20$ ) resistance levels.

**Table 2** Antibiotic resistance levels as percentages of plate count enumerations on growth media amended with antibiotic compared to plate count enumerations on unamended growth media

Site	LB+chlortetracycline	PYT80+chlortetracycline	PYT80+tylosin	PYT80+carbadox
SFO1	0.15 ± 0.05	0.2 ± 0.03	17 ± 10	42 ± 13
SFO2	0.001 ± 0.002	0.1 ± 0.08	7 ± 3	21 ± 5
SFO3	0.006 ± 0.003	0.003 ± 0.002	5 ± 2	32 ± 33
SFO4	0.1 ± 0.04	0.02 ± 0.008	12 ± 6	24 ± 13
DF1	0.01 ± 0.007	0.01 ± 0.009	16 ± 12	15 ± 2
DF2	0.09 ± 0.08	0.04 ± 0.02	3 ± 2	15 ± 6
DF3	0.02 ± 0.02	0.003 ± 0.003	12 ± 6	47 ± 30
NAG1	n.d.	0.3 ± 0.4	6 ± 3	10 ± 5
NAG2	n.d.	0.05 ± 0.02	55 ± 60	24 ± 18
NAG3	0.03 ± 0.04	0.007 ± 0.01		

Abbreviations: DF, dairy farm; NAG, non-agricultural; SFO, swine feeding operations.

Results are presented as the arithmetic means of enumerations of triplicate soil samples at each site ± one s.d. of the mean. n.d., no chlortetracycline-resistant bacteria were detected.

Bacteria resistant to chlortetracycline were cultivated on both nutrient rich and nutrient poor growth media from all 10 sites (Table 2), with the exception of Site NAG2 where no chlortetracycline-resistant bacteria could be enumerated on nutrient-rich media. In contrast to the tylosin and carbadox resistance levels, ANOVA suggested that there were statistically significant differences between bacterial resistance to chlortetracycline among the sites, as enumerated on both nutrient-rich (LB;  $P < 0.0005$ ) and nutrient-poor ( $P < 0.05$ ) agar plates. More rigorous statistical analysis (Tukey's honest significant difference;  $P < 0.05$ ) suggested that only the enumerations on nutrient-rich media from sites SFO1 and SFO4 were significantly higher than the other sites. This same test suggested that there were no statistically significant differences among the bacterial resistance levels enumerated on nutrient-poor media.

#### Identification of chlortetracycline-resistant bacteria

Because differences were detected among the resistance levels to chlortetracycline, more than 750 bacterial strains were isolated from the 10 sites and analyzed in further detail. Bacterial isolates were initially screened by RISA to identify unique strains. Although about 10% of the isolates failed to generate a fingerprint, RISA was able to identify successfully unique bacterial strains among the rest of the isolates. All of the unique strains, as well as the ones that failed to generate a RISA fingerprint, were then identified by partial 16S rRNA gene sequence analysis (Tables 3 and 4).

The type of growth media had a substantial impact on the type of chlortetracycline-resistant bacteria isolated from the different sites. Low G+C Gram-positive (*Firmicutes*) and high G+C Gram-positive (*Actinobacteria*) bacteria were the most prominent chlortetracycline-resistant bacteria isolated on nutrient-rich media, although a few members of the *Proteobacteria* were also detected (Table 3). Particularly common among these isolates were *Streptomyces* spp., which were the most abundant chlortetracycline-resistant organisms at two of the swine farms (Sites SFO2 and SFO3), two of the DFs (Sites DF2 and DF3) and two of the non-agricultural sites (Sites NAG1 and NAG3). Site SFO1 had the highest diversity of resistant bacteria (16 of the 47 isolates were unique) with a high prevalence of *Firmicutes* and *Proteobacteria* and an absence of *Streptomyces* spp.

The types of chlortetracycline-resistant bacteria enumerated on nutrient-poor growth media were predominantly *Chryseobacteria* spp. and different types of *Proteobacteria*, with a bias towards *Variovorax* spp. (Table 4). Site NAG1 was distinctive compared to the other sites because only *Streptomyces* spp. were detected on nutrient-poor media. Site SFO1 again contained the most diverse group of resistant bacteria enumerated on nutrient-poor media (11 of 37 colonies were unique).

PCA was performed separately on chlortetracycline-resistant isolates on nutrient-rich and nutrient-poor media to evaluate differences between sites based on the types of resistant bacteria and their abundance (Figure 1). Site SFO1 strongly separated from all the other sites on the first principal component axis for both media types. Site SFO4 separated on the second principal component axis with only the nutrient-rich media (Figure 1a). Similarly, Site NAG2 separated from the rest on the second principal component axis with the nutrient-poor media (Figure 1b).

#### Multiplex PCR of genes encoding for resistance to tetracycline

All of the chlortetracycline-resistant bacterial isolates (that is, even those identified as redundant by RISA) were characterized by the presence or absence of 14 different genes that encode for tetracycline resistance. Numerous genes encoding for tetracycline resistance were detected, predominantly among the *Firmicutes* and *Proteobacteria* that had been isolated from different sites on nutrient-rich media (Table 3). The highest frequencies of detection of these genes encoding for tetracycline resistance were found in bacteria isolated from Sites SFO1 (43 of 47 isolates, with some of them harboring up to three different genes) and SFO4 (26 of 49 isolates). Genes encoding for tetracycline resistance were also commonly detected among bacteria isolated from Site SFO1 on nutrient-poor media (28 of 37 isolates) (Table 4). In contrast, bacteria isolated from the other nine sites that were isolated on nutrient-poor media rarely (2 of 293 isolates) contained one of these 14 genes encoding for tetracycline resistance.

#### Persistence of chlortetracycline resistance at Sites SFO1 and SFO4

Coincidental to the onset of our study, farming operations ceased at Site SFO1, allowing us to study the long-term persistence of antibiotic resistance. Bacterial enumerations on both nutrient-rich and nutrient-poor media suggested that resistance levels were sustained for at least 18 months and there was no significant temporal variation in resistance levels (Figure 2a). Furthermore, soil samples collected in July 2005 at three different locations at Site SFO1 (at distances of 5, 20 and 100 m from the animal pen) suggested that the former location of the animal pen remained a hot spot for chlortetracycline resistance (Figure 2b).

Approximately 300 chlortetracycline-resistant bacteria were isolated and characterized from soil samples collected from three different locations at Site SFO1 in July 2005. Of the bacteria isolated on nutrient-rich media, there was considerable overlap in the types of resistant bacteria detected at 5 m from the animal pen compared to samples collected 20 m

**Table 3** Chlortetracycline-resistant bacteria isolated from nine sites on nutrient-rich growth medium including April 2004 samples from Site SFO1

Site	Bacterial division	Best phylogenetic match (accession no.)	Length (bp)	% identity	No. of isolates	Presence of tet gene			
SFO1	Firmicutes	<i>Bacillus</i> sp. (CP000001)	551	100	7	<i>tet</i> (L)			
		<i>Bacillus</i> sp. (AB098575)	550	100	7	<i>tet</i> (L)			
		<i>Bacillus</i> sp. (AY822613)	534	99.8	1	<i>tet</i> (L)			
		<i>Bacillus</i> sp. (AF202056)	533	99.4	1				
		<i>Bacillus</i> sp. (BGA535638)	531	100	1	<i>tet</i> (L)			
		<i>Bacillus</i> sp. (AB211020)	547	98.2	1				
		<i>Oceanobacillus</i> sp. (AB188089)	550	99.3	3	<i>tet</i> (L)			
		<i>Oceanobacillus</i> sp. (AY553089)	550	100	1	<i>tet</i> (L)			
		<i>Staphylococcus</i> sp. (AF527483)	548	100	2	<i>tet</i> (A), <i>tet</i> (L)			
	$\alpha$ -Proteobacteria	<i>Enterococcus</i> sp. (AB188089)		550	99.3	1	<i>tet</i> (M)		
						1	<i>tet</i> (M), <i>tet</i> (L)		
						1	<i>tet</i> (M)		
		<i>Brevundimonas</i> sp. (D49422)		521	99.7	5	<i>tet</i> (G)		
			<i>Ochrobactrum</i> sp. (AY331580)		523	100	2	<i>tet</i> (G)	
							1	<i>tet</i> (G), <i>tet</i> (L)	
	$\gamma$ -Proteobacteria	<i>Pseudomonas</i> sp. (DQ084459)		532	99.6	4	<i>tet</i> (A)		
						1	<i>tet</i> (A), <i>tet</i> (L)		
						1	<i>tet</i> (A), <i>tet</i> (L)		
						1	<i>tet</i> (A), <i>tet</i> (M)		
		<i>Pseudomonas</i> sp. (PH16SRNA1)		528	99.1	1			
			<i>Serratia</i> sp. (AF286868)	548	99.3	3	<i>tet</i> (B)		
<i>Acinetobacter</i> sp. (AC17008)			550	99.8	1	<i>tet</i> (M)			
SFO2	Actinobacteria	<i>Streptomyces</i> sp. (AJ781328)	531	100	10				
		<i>Streptomyces</i> sp. (AY277559)	531	99.8	26				
SFO3	Actinobacteria	<i>Streptomyces</i> sp. (AY465336)	449	100	5				
		<i>Streptomyces</i> sp. (SAL494865)	534	98.5	2				
		<i>Streptomyces</i> sp. (AF503493)	498	100	1				
		<i>Bacillus</i> sp. (AF286486)	538	100	1	<i>tet</i> (L)			
	$\gamma$ -Proteobacteria	<i>Bacillus</i> sp. (BSI6086)		521	99.8	2	<i>tet</i> (M)		
			<i>Bacillus</i> sp. (AB211020)	547	98.2	2	<i>tet</i> (L)		
			<i>Paenibacillus</i> sp. (PCI575659)	550	99.6	2			
			<i>Paenibacillus</i> sp. (AB073188)	550	99.1	1			
			<i>Sporosarcina</i> sp. (AF506059)	552	97.8	1	<i>tet</i> (L)		
			<i>Xanthomonas</i> sp. (XSP244722)	523	98.9	1			
			<i>Serratia</i> sp. (AY394724)	538	97.6	1			
		SFO4	Actinobacteria	<i>Streptomyces</i> sp. (AF128874)	531	100	1		
				<i>Firmicutes</i>	<i>Aerococcus</i> sp. (AF076639)	548	100	9	<i>tet</i> (M)
			$\gamma$ -Proteobacteria	<i>Staphylococcus</i> sp. (AB009936)		548	100	5	<i>tet</i> (L)
					<i>Staphylococcus</i> sp. (AY161045)	550	99.6	2	<i>tet</i> (M)
					1	<i>tet</i> (L)			
<i>Staphylococcus</i> sp. (AF527483)	548	100			2	<i>tet</i> (L), <i>tet</i> (M)			
					1	<i>tet</i> (L)			
<i>Kurthia</i> sp. (KZ16RNAA)	548	100			5	<i>tet</i> (L), <i>tet</i> (M)			
<i>Vagococcus</i> sp. (AY179329)		548		99.1	2				
	<i>Streptococcus</i> sp. (AF459431)	548		100	1				
	<i>Psychrobacter</i> sp. (AJ871083)	539		100	1	<i>tet</i> (O)			
					7				
	<i>Acinetobacter</i> sp. (AY167273)	550		100	4				
DF1	Actinobacteria	<i>Acinetobacter</i> sp. (ASP551155)	550	98.7	2				
		<i>Streptomyces</i> sp. (SB16SRRN)	531	100	2	<i>tet</i> (L)			
		<i>Streptomyces</i> sp. (AY094368)	531	99.6	1				
	Firmicutes	<i>Lactococcus</i> sp. (AY675242)	545	100	5	<i>tet</i> (S)			
					1				
		<i>Oceanobacillus</i> sp. (AY553089)	550	100	2	<i>tet</i> (L)			
					1				
		<i>Virgibacillus</i> sp. (AY422988)	550	100	2	<i>tet</i> (L)			
					1				
		<i>Staphylococcus</i> sp. (AY030342)	546	100	1	<i>tet</i> (K)			
DF2	$\gamma$ -Proteobacteria	<i>Bacillus</i> sp. (BSP302941)	551	100	1	<i>tet</i> (L)			
		<i>Pseudomonas</i> sp. (AB091837)	547	100	4				
	Actinobacteria	<i>Streptomyces</i> sp. (AY277559)	531	99.8	17				
		<i>Firmicutes</i>	<i>Bacillus</i> sp. (AY456223)	553	98.6	3			
	$\gamma$ -Proteobacteria	<i>Paenibacillus</i> sp. (AY308758)	550	99.8	1				
		<i>Serratia</i> sp. (AY337583)	439	96.6	6				
<i>Morganella</i> sp. (AF461011)		553	91.9	1	<i>tet</i> (D)				

**Table 3** (Continued)

Site	Bacterial division	Best phylogenetic match (accession no.)	Length (bp)	% identity	No. of isolates	Presence of tet gene	
DF3	Actinomycetes	<i>Streptomyces</i> sp. (AY277559)	531	99.8	19		
		<i>Streptomyces</i> sp. (AY741287)	533	99.2	1		
	Firmicutes	<i>Paenibacillus</i> sp. (AY308758)	550	99.8	1		
		<i>Variovorax</i> sp. (AJ746113)	547	99.3	1		
		<i>Obesumbacterium</i> sp. (OPR233422)	532	99.6	1		
NAG1	Actinobacteria	<i>Streptomyces</i> sp. (AF128874)	531	100	16		
NAG3	Actinobacteria	<i>Streptomyces</i> sp. (AJ621602)	531	99.8	16		
		<i>Streptomyces</i> sp. (AY277559)	531	99.8	7		
		<i>Streptomyces</i> sp. (AY572485)	495	100	5		
		<i>Streptomyces</i> sp. (AY396146)	498	100	3		
		<i>Streptomyces</i> sp. (AJ781328)	531	100	1		
		<i>Streptomyces</i> sp. (AB119009)	498	99.0	1		
		Firmicutes	<i>Virgibacillus</i> sp. (AY422988)	550	100	2	<i>tet(L)</i>
			<i>Paenibacillus</i> sp. (AY308758)	550	99.6	1	<i>tet(L)</i>
						1	
				<i>Bacillus</i> sp. (AF071858)	504	100	1
			<i>Bacillus</i> sp. (AB211020)	547	98.2	1	<i>tet(L)</i>
			<i>Bacillus</i> sp. (AY148429)	484	98.3	1	
	γ-Proteobacteria	<i>Serratia</i> sp. (AY498856)	509	100	1		
		Bacteroidetes	<i>Chryseobacterium</i> sp. (AY468451)	503	99.6	2	

Abbreviations: DF, dairy farm; NAG, non-agricultural; SFO, swine feeding operations

Length (column 4) refers to the length of the 16S rRNA gene sequence used to determine the percent identity (column 5) to the best phylogenetic match.

from the pen (Table 5). These bacteria were phylogenetically similar to the bacteria that were previously detected (Table 3), although there was an increase in the frequency of *Bacillus* spp., which correlated to a decrease in the diversity of resistant bacteria. In contrast, *Streptomyces* spp. were most prevalent (21 of 41 isolates) among the chlortetracycline-resistant bacteria isolated from the soil collected about 100 m from the animal pen.

Of the bacteria isolated on nutrient-poor media, the bacteria detected at 5 and 20 m from the animal pen in July 2005 (Table 6) were phylogenetically similar to those bacteria previously isolated on nutrient-poor media (Table 4). Further away from the animal pen (100 m), *Variovorax* spp. were the most commonly detected chlortetracycline-resistant bacteria (18 of 24 isolates). There was also a shift in the prevalence of genes encoding for tetracycline resistance as a function of distance from the animal pen. More than half of the bacteria (31 of 45 isolates) isolated 5 m from the animal pen possessed at least one of 14 genes encoding for tetracycline resistance. In contrast, only a small fraction (4 of 24 isolates) of the bacteria isolated 100 m from the animal pen harbored one of the 14 genes encoding for tetracycline resistance.

Although marginally elevated chlortetracycline resistance levels were originally detected at Site SFO4, we speculated that this result was biased because these soil samples were collected only 1 week subsequent to manure application. Subsequent samples were therefore collected after an additional 10 months of time. Chlortetracycline-resistant bacteria were not detected on the nutrient-

rich media, while resistance levels on the nutrient-poor media were statistically similar to all of the other sites (except Site SFO1) (data not shown).

## Discussion and conclusions

The subtherapeutic use of antibiotics in agriculture has been a controversial subject for decades (Levy, 1978). Along with rising concerns about the failure of many antibiotics in treating diseases, evidence suggests that the use of agricultural antibiotics is contributing to the spread of antibiotic resistance in the environment (Ferber, 2003; Onan and LaPara, 2003; Wegener, 2003; Rooklidge, 2004). There is also speculation that agricultural use is responsible for continued prevalence of resistance to certain antibiotics after discontinuing their clinical use (Livermore, 2003). However, the lack of causal linkage between agricultural use and the proliferation of antibiotic resistance among pathogenic bacteria has led some people to believe that the economic benefit of antibiotic use in agriculture is sufficient to continue this practice (Phillips *et al.*, 2004).

In our study, we quantified and characterized antibiotic-resistant bacteria in soils at farms using antibiotics for non-therapeutic purposes as well as at farms using antibiotics predominantly for veterinary purposes. The goal of this study was to discern the impact of antibiotic use on the occurrence and abundance of cultivable antibiotic-resistant bacteria in nearby soils exposed to animal waste. Elevated levels of chlortetracycline resistance were quantified in the soils at some, but not all, of the farms that

**Table 4** Chlortetracycline-resistant bacteria isolated from soils at 10 different sites on nutrient-poor growth medium

Site	Bacterial division	Best phylogenetic match (accession no.)	Length (bp)	% identity	No. of isolates	Presence of tet gene	
SFO1	Firmicutes	<i>Bacillus</i> sp.(AY305275)	551	100	1	<i>tet</i> (L)	
		$\alpha$ -Proteobacteria	<i>Ochrobactrum</i> sp. (AY331580)	523	100	3	<i>tet</i> (G)
	$\gamma$ -Proteobacteria		<i>Brevundimonas</i> sp. (AY571823)	519	99.0	1	<i>tet</i> (G)
		<i>Brevundimonas</i> sp. (D49422)	521	99.6	1	<i>tet</i> (G)	
		<i>Pseudomonas</i> sp. (AY263482)	545	100	10	<i>tet</i> (A)	
		<i>Pseudomonas</i> sp. (AF326378)	547	100	3	<i>tet</i> (A)	
		<i>Pseudomonas</i> sp. (D85999)	547	100	1	<i>tet</i> (A)	
		<i>Pseudomonas</i> sp. (AY653222)	547	99.5	1	<i>tet</i> (A)	
		<i>Serratia</i> sp. (AF286868)	548	99.3	5	<i>tet</i> (B)	
		<i>Stenotrophomonas</i> sp. (AY689084)	533	99.8	1		
		<i>Sphingobacterium</i> sp. (AY635870)	538	99.8	2	<i>tet</i> (X)	
	SFO2	$\alpha$ -Proteobacteria	<i>Caulobacter</i> sp. (CHE7805)	419	99.5	2	
			<i>Variovorax</i> sp. (AF214127)	548	100	25	
Bacteriodetes		<i>Chryseobacterium</i> sp. (AF375840)	147	100	16		
		<i>Chryseobacterium</i> sp. (AY468464)	298	99.3	4		
SFO3	$\beta$ -Proteobacteria	<i>Variovorax</i> sp. (AF214127)	546	100	1		
SFO4	Bacteriodetes	<i>Chryseobacterium</i> sp. (AJ874979)	530	99.1	20		
	Firmicutes	<i>Kurthia</i> sp. (KZ16RNAA)	548	100	1		
SFO4	$\alpha$ -Proteobacteria	<i>Brevundimonas</i> sp. (AJ717390)	523	100	1	<i>tet</i> (G)	
		<i>Variovorax</i> sp. (AF214127)	546	100	45		
	$\beta$ -Proteobacteria	<i>Burkholderia</i> sp. (AY268162)	548	100	2		
DF1	$\beta$ -Proteobacteria	<i>Variovorax</i> sp. (AF214127)	546	100	13		
		<i>Herbaspirillum</i> sp. (AB027694 )	150	97.3	3		
	$\gamma$ -Proteobacteria	<i>Pseudomonas</i> sp. (AB091837)	149	98.7	1		
DF2	Bacteriodetes	<i>Chryseobacterium</i> sp. (AY751083)	145	100	13		
	$\alpha$ -Proteobacteria	<i>Caulobacter</i> sp. (CHE7805)	511	99.6	1		
	$\beta$ -Proteobacteria	<i>Variovorax</i> sp. (AF214127)	546	100	45		
DF3	$\gamma$ -Proteobacteria	<i>Pseudomonas</i> sp. (AB091837)	149	98.7	1		
	$\beta$ -Proteobacteria	<i>Variovorax</i> sp.(AJ746113)	547	99.3	14		
	$\gamma$ -Proteobacteria	<i>Lysobacter</i> sp. (AB083480)	550	99.6	1		
NAG1	Actinobacteria	<i>Streptomyces</i> sp. (AF128874)	531	100	16		
NAG2	$\beta$ -Proteobacteria	<i>Variovorax</i> sp. (AF532867)	534	95.1	2		
		<i>Variovorax</i> sp. (AF214127)	546	100	1		
	$\gamma$ -Proteobacteria	<i>Luteibacter</i> sp. (LJ9580499)	546	99.1	9		
		<i>Stenotrophomonas</i> sp. (AY512625)	550	100	7		
		<i>Stenotrophomonas</i> sp. (AY040357)	550	100	1		
	<i>Xanthomonas</i> sp. (AY841369)	550	100	4			
	<i>Serratia</i> sp. (AF286868)	547	99.5	1	<i>tet</i> (C)		
	Bacteriodetes	<i>Chryseobacterium</i> sp. (AF531766)	523	94.7	14		
		<i>Chryseobacterium</i> sp. (AF375840)	144	100	3		
		<i>Rhizobium</i> sp. (RS16SCIAM)	493	100	1		
NAG3	$\alpha$ -Proteobacteria	<i>Variovorax</i> sp. (AF214127)	546	100	12		
	$\beta$ -Proteobacteria	<i>Chryseobacterium</i> sp. (AY468471)	356	100	11		
	Bacteriodetes	<i>Chryseobacterium</i> sp. (AY468451)	503	99.6	1		


Abbreviations: DF, dairy farm; NAG, non-agricultural; SFO, swine feeding operations.

Length (column 4) refers to the length of the 16S rRNA gene sequence used to determine the percent Identity (column 5) to the best phylogenetic match.

used antibiotics for non-veterinary purposes. When elevated levels of antibiotic-resistant bacteria were enumerated, significant shifts were also observed in the types of chlortetracycline-resistant bacteria as well as the types of genes that encode resistance within these organisms.


Contrary to our expectations, the farm with the highest chlortetracycline resistance levels was the smallest swine farm (Site SFO1), which housed no more than 50 animals at a time. These elevated

resistance levels were concomitantly associated with a higher frequency of 14 genes encoding for resistance to tetracycline. We speculate that the excessive exposure to animal manure was responsible for the observed result. While the other farms in our study collected and applied manure to soil at agronomically appropriate rates, Site SFO1 allowed manure to spill uncontrolled out of the animal pen. Previous studies have shown that manure from animals fed subtherapeutic concentrations of anti-



**Figure 1** Principal component analysis of the types (based on partial 16S rRNA gene sequence) and abundance of chlortetracycline-resistant isolates from the 10 sites on (a) nutrient-rich and (b) nutrient-poor growth media. For the nutrient-rich medium, principal component 1 and principal component 2 explain 34 and 22% of variation in the data, respectively. For the nutrient-poor medium, principal component 1 and principal component 2 explain 32 and 17% of variation in the data, respectively.

biotics contains numerous resistant bacteria (Levy, 1978; Aarestrup *et al.*, 2002; Sørum *et al.*, 2006), but that resistance among soil bacteria returned to preapplication levels within 6 months of manure application (Sengelov *et al.*, 2003). We hypothesize, however, that if an excessive quantity of these bacteria is applied to soil, then antibiotic resistance can spread among the indigenous soil bacteria because of lateral gene transfer of the resistance determinants (Andrews *et al.*, 2004). The spread of



**Figure 2** (a) Chlortetracycline resistance levels at SFO1 between November 2003 and July 2005. Animal feeding operations ceased in November 2003. (b) Chlortetracycline resistance levels as measured in July 2005 at SFO1 at varying distances from the location of the animal pen. Solid bars represent the percentage of chlortetracycline-resistant bacteria enumerated on nutrient-rich medium; open bar represent the percentage of chlortetracycline-resistant bacteria enumerated on nutrient-poor medium.

genes encoding for resistance to indigenous soil bacteria is potentially pertinent because these organisms should be better adapted for survival in the soil, thus potentially serving as a long-term reservoir of antibiotic resistance.

More detailed analysis of the genes encoding tetracycline resistance further suggested that lateral

**Table 5** Chlortetracycline-resistant bacteria isolated from Site SFO1 in July 2005 on nutrient-rich growth medium

Site	Bacterial division	Best phylogenetic match (accession no.)	Length (bp)	% identity	No. of isolates	Presence of tet gene	
5 m	Firmicutes	<i>Bacillus</i> sp. (CP000001)	551	100	16	<i>tet(L)</i>	
		<i>Bacillus</i> sp. (AY548950)	534	100	8	<i>tet(L)</i>	
						6	<i>tet(L)</i> , <i>tet(M)</i>
		<i>Bacillus</i> sp. (AB098575)	550	100	10	<i>tet(L)</i>	
		<i>Bacillus</i> sp. (AY988598)	551	100	1		
	$\gamma$ -Proteobacteria	<i>Bacillus</i> sp. (AF513458)	547	99.3	1	<i>tet(L)</i>	
		<i>Paenibacillus</i> sp. (PSP297713)	549	100	3	<i>tet(M)</i>	
		<i>Enterobacter</i> sp. (AB098582)	533	99.2	1	<i>tet(A)</i>	
						1	<i>tet(B)</i>
		<i>Morganella</i> sp. (AF461011)	553	91.9	1	<i>tet(D)</i>	
20 m	Firmicutes	<i>Pseudomonas</i> sp. (AY972267)	509	99.8	1	<i>tet(A)</i> , <i>tet(L)</i>	
		<i>Bacillus</i> sp. (CP000001)	551	100	27	<i>tet(L)</i>	
		<i>Bacillus</i> sp. (AB098575)	550	100	7	<i>tet(L)</i>	
	$\gamma$ -Proteobacteria	<i>Bacillus</i> sp. (AY548950)	534	100	3	<i>tet(L)</i>	
						2	<i>tet(L)</i> , <i>tet(M)</i>
		<i>Pantoea</i> sp. (AY691543)	533	99.8	3	<i>tet(B)</i>	
		<i>Pseudomonas</i> sp. (DQ182328)	532	100	1	<i>tet(A)</i> , <i>tet(L)</i>	
		<i>Pseudomonas</i> sp. (AB091837)	547	100	1		
		<i>Pseudomonas</i> sp. (DQ192039)	525	99.8	1	<i>tet(A)</i> , <i>tet(L)</i>	
		<i>Enterobacter</i> sp. (AB098582)	533	99.2	1	<i>tet(B)</i>	
100 m	Actinobacteria	<i>Providencia</i> sp. (AM040495)	532	100	1		
		<i>Streptomyces</i> sp. (AY741286)	524	99.6	18		
		<i>Streptomyces</i> sp. (AY094368)	531	99.6	3		
	Firmicutes	<i>Rhodococcus</i> sp. (AB210967)	528	100	1		
		<i>Bacillus</i> sp. (AF513458)	547	99.3	5	<i>tet(L)</i>	
						1	<i>tet(L)</i> , <i>tet(M)</i>
		<i>Bacillus</i> sp. (AB098575)	550	100	5	<i>tet(L)</i>	
		<i>Bacillus</i> sp. (DQ198162)	545	100	1	<i>tet(L)</i>	
						1	<i>tet(L)</i> , <i>tet(M)</i>
		<i>Bacillus</i> sp. (CP000001)	551	100	1	<i>tet(L)</i>	
$\gamma$ -Proteobacteria	<i>Bacillus</i> sp. (DQ166809)	547	99.8	1	<i>tet(L)</i>		
	<i>Bacillus</i> sp. (AY548950)	534	100	1	<i>tet(L)</i> , <i>tet(M)</i>		
	<i>Paenibacillus</i> sp. (PSP297713)	549	100	1	<i>tet(M)</i>		
	<i>Paenibacillus</i> sp. (AY266990)	545	99.8	1			
	<i>Pseudomonas</i> sp. (DQ073452)	549	100	1	<i>tet(A)</i> , <i>tet(L)</i>		

Abbreviation: SFO, swine feeding operations.

Length (column 4) refers to the length of the 16S rRNA gene sequence used to determine the percent identity (column 5) to the best phylogenetic match.

gene transfer was a pertinent factor in the persistence of chlortetracycline resistance at Site SFO1. Numerous bacteria harbored the same gene encoding for tetracycline resistance, similar bacteria (based on 16S rRNA gene sequences) harbored different genes encoding for tetracycline resistance, and individual bacteria harbored multiple types of genes encoding for tetracycline resistance. Particularly noteworthy is the detection of *tet(A)* in *Bacillus* spp. and *tet(L)* in different types of *Proteobacteria*. Both *tet(A)* and *tet(L)* genes encode for efflux pumps that are predominantly associated with gram-negative and gram-positive bacteria, respectively (Chopra and Roberts, 2001; Aminov *et al.*, 2002). The detection of *tet(X)* in a *Sphingobacteria* spp., which encodes for a tetracycline-degrading mono-oxygenase (Yang *et al.*, 2004), is also interesting because this gene, thus far, has been found only in *Bacteroides* spp. (Whittle *et al.*, 2001), which are obligate anaerobes, perhaps suggesting that we have found the natural host of this gene. We are currently characterizing this organism in more detail.

The results from our study have important practical implications. Three of the swine farms were able to subtherapeutically use antibiotics without leading to significantly different numbers or types of chlortetracycline-resistant bacteria compared to the dairy farm soils. These swine farms temporarily stored their animal waste in covered, underground storage pits until it was applied to an appropriate surface area of land per Minnesota state regulations (Minnesota Pollution Control Agency, 2000). We conclude, therefore, that a relatively small quantity of manure containing antibiotic-resistant bacteria can be spread on agricultural soils without significant lateral gene transfer of resistance determinants to the indigenous soil bacteria, allowing the resistant bacteria in the manure to slowly decay in the soil without leading to the proliferation of resistance.

An ancillary goal of our research was to compare resistant bacteria from soils at farms using antibiotics with resistant bacteria from soils unaffected by antibiotic use. Because many antibiotics are

**Table 6** Chlortetracycline resistant bacteria isolated from site SFO1 in July 2005 on nutrient-poor growth medium

Site	Bacterial division	Best phylogenetic match (accession no.)	Length (bp)	% identity	No. of isolates	Presence of tet gene		
5 m	Firmicutes	<i>Bacillus</i> sp. (CP000001)	551	100	7	<i>tet(L)</i>		
		<i>Pseudomonas</i> sp. (DQ182328)	532	100	1	<i>tet(A)</i> , <i>tet(L)</i>		
	$\gamma$ -Proteobacteria	<i>Pseudomonas</i> sp. (D85999)	<i>Pseudomonas</i> sp. (DQ073452)	549	100	7	<i>tet(A)</i>	
			<i>Pseudomonas</i> sp. (PFL308320)	549	100	1	<i>tet(A)</i> , <i>tet(B)</i>	
			<i>Stenotrophomonas</i> sp. (AM402950)	481	100	1	<i>tet(A)</i>	
			<i>Serratia</i> sp. (DQ207558)	549	100	1	<i>tet(G)</i> , <i>tet(L)</i>	
			Bacterioidetes	<i>Chryseobacterium</i> sp. (AY468469)	522	98.7	9	<i>tet(A)</i>
				<i>Chryseobacterium</i> sp. (AY468471)	541	99.8	2	<i>tet(A)</i>
				<i>Chryseobacterium</i> sp. (AY468469)	522	99.8	3	<i>tet(A)</i>
	20 m	Firmicutes	<i>Bacillus</i> sp. (CP000001)	551	100	1		
			<i>Ochrobactrum</i> sp. (AY331580)	523		3	<i>tet(G)</i>	
		$\alpha$ -Proteobacteria	<i>Brevundimonas</i> sp. (D49422)	<i>Brevundimonas</i> sp. (AB091837)	521	99.1	2	<i>tet(B)</i> , <i>tet(A)</i>
				<i>Pseudomonas</i> sp. (AY263482)	545	100	1	
<i>Pseudomonas</i> sp. (AF326378)				547	100	8	<i>tet(A)</i>	
<i>Pseudomonas</i> sp. (AY271792)				532	100	2	<i>tet(A)</i>	
<i>Lysobacter</i> sp. (AB083480)				532	100	2	<i>tet(G)</i>	
<i>Klebsiella</i> sp. (AY786181)				532	99.6	6	<i>tet(A)</i>	
Bacterioidetes				<i>Chryseobacterium</i> sp. (AY468469)	522	98.7	1	<i>tet(A)</i>
		<i>Chryseobacterium</i> sp. (AY468475)	524	99.4	7			
		<i>Chryseobacterium</i> sp. (AY468469)	522	99.8	2			
100 m		Firmicutes	<i>Bacillus</i> sp. (CP000001)	551	100	1		
			<i>Variovorax</i> sp. (AF214127)	548	100	2		
	$\beta$ -Proteobacteria	<i>Pseudomonas</i> sp. (DQ182328)	<i>Pseudomonas</i> sp. (AY263482)	545	100	18	<i>tet(A)</i>	
			<i>Pseudomonas</i> sp. (DQ182328)	532	100	1	<i>tet(A)</i>	
			<i>Pseudomonas</i> sp. (AY263482)	545	100	2	<i>tet(A)</i>	
			<i>Pseudomonas</i> sp. (DQ182328)	532	100	1	<i>tet(A)</i>	
			<i>Pseudomonas</i> sp. (AY263482)	545	100	1	<i>tet(A)</i>	

Abbreviation: SFO, swine feeding operations.

Length (column 4) refers to the length of the 16S rRNA gene sequence used to determine the percent identity (column 5) to the best phylogenetic match.

natural compounds, numerous soil bacteria would be expected to be resistant even if humans had not been mass-producing various antibiotics since the 1940s (Davelos *et al.*, 2004; Guardabassi and Dalsgaard, 2004). It was somewhat surprising, therefore, that the soil bacteria from the three DFs and two of the farms using antibiotics at subtherapeutic concentrations had similar quantities and types of resistant bacteria compared to the soils that had not been affected by agricultural activity. It is particularly interesting that the bacteria isolated from Quetico Provincial Park (that is, the site least affected by human activity) were exclusively *Streptomyces* spp., which naturally produce many antibiotics and are equipped with various defense mechanisms.

A principal limitation of the present study is that a cultivation-based approach was used to quantify and characterize chlortetracycline-resistant bacteria in our soil samples. Cultivation-based approaches are known to detect only a small fraction (usually

about 1% of soil bacteria) of the total bacterial community (Amann *et al.*, 1995). Although two different growth media were used for chlortetracycline to help limit this bias, we assume that our approach was able to detect only a small fraction of the chlortetracycline-resistant bacteria in these soils. In contrast, cultivation-independent analysis of chlortetracycline resistance would have its own limitations and biases. Cultivation-independent community analysis of chlortetracycline resistance would require that the genes encoding for resistance be analyzed directly (for example, by quantitative PCR), which would have provided a dataset of limited usefulness because the organisms that would have harbored these genes would have remained unknown. Second, cultivation-independent analysis of chlortetracycline resistance would be especially complex because there are now more than 40 different genes that are known to encode for tetracycline resistance (Roberts, 2005).

The diversity and complexity of the genetics of tetracycline resistance also represents another substantial limitation of the present study. Although the multiplex PCR approach used herein was able to reliably detect 14 common genes that encode for resistance to tetracycline, we were unable to detect any gene that encoded for resistance to tetracycline in the majority of our isolates – particularly among the strains isolated from soil samples other than Site SFO1. These bacteria, which did not harbor any of the genes for which we assayed, could have been intrinsically resistant to tetracycline (especially the *Streptomyces* spp.), harbored one of the other previously discovered genes that encodes for resistance to tetracycline, or harbored a novel gene encoding for resistance to tetracycline. Because soil bacterial communities are particularly diverse (Torsvik *et al.*, 1990), they can also potentially serve as a critical reservoir of genetic material for antibiotic resistance (D'Costa *et al.*, 2006). Additional research is needed to further elucidate the diversity of resistance determinants in environmental hotspots of antibiotic resistance.

Our hypothesis is that the proliferation of antibiotic resistance is principally modulated by the number of antibiotic-resistant bacteria, the genes harbored by these bacteria, the relative ease by which genes encoding for resistance can be laterally transferred among different bacterial strains, and any selective pressures that provide a competitive advantage for antibiotic-resistant bacteria. At the present time, the scientific community appears to be particularly focused on the latter factor (selective pressures), which are primarily linked to antibiotic use. Although we agree that antibiotic use is the most pertinent factor in the global spread of resistance, our research demonstrates that the proliferation of antibiotic resistance is not intrinsically linked to antibiotic use and that the other three factors can also be critically important. We recommend, therefore, substantially more stringent control of fecal waste (both of animal and of human origin) as a viable approach to slow the proliferation of antibiotic-resistant bacteria.

In conclusion, our results suggest that the sub-therapeutic use of antibiotics can lead to the proliferation of antibiotic resistance among soil bacteria if an excessive quantity of manure is applied to an insufficient surface area of land. Additional research is needed, however, to confirm or refute this conclusion. In contrast, our results revealed no increase in the quantity of resistant bacteria when animal manure was applied at agronomically acceptable rates. Of particular importance is that the proliferation of antibiotic resistance appears to be associated with specific genes that confer antibiotic resistance. Because these genes were detected among numerous different bacterial populations, lateral genetic exchange appears to be a pertinent mechanism by which resistance develops among soil bacteria.

## Acknowledgements

This work was financially supported by the National Research Initiative of the United States Department of Agriculture Cooperative State Research, Education and Extension Service (Grant number 2003–35107–13830).

## References

- Aarestrup FM, Hasman H, Jensen LB, Moreno M, Herrero IA, Dominguez L *et al.* (2002). Antimicrobial resistance among enterococci from pigs in three European countries. *Appl Environ Microbiol* **68**: 4127–4129.
- Aarestrup FM, Seyfarth AM, Emborg HD, Pedersen K, Hendriksen RS, Bager F. (2001). Effect of abolishment of the use of antimicrobial agents for growth promotion on occurrence of antimicrobial resistance in fecal enterococci from food animals in Denmark. *Antimicrob Agents Chemother* **45**: 2054–2059.
- Amann RI, Ludwig W, Schleifer KH. (1995). Phylogenetic identification and *in situ* detection of individual microbial cells without cultivation. *Microbiol Rev* **59**: 143–169.
- Aminov RI, Chee-Sanford JC, Garrigues N, Teferedegne B, Krapac IJ, White BA *et al.* (2002). Development, validation, and application of PCR primers for detection of tetracycline efflux genes of gram-negative bacteria. *Appl Environ Microbiol* **68**: 1786–1793.
- Andrews RE, Johnson WS, Guard AR, Marvin JD. (2004). Survival of *Enterococci* and Tn916-like conjugative transposons in soil. *Can J Microbiol* **50**: 957–966.
- Arnold S, Gassner B, Giger T, Zwahlen R. (2004). Banning antimicrobial growth promoters in feedstuffs does not result in increased therapeutic use of antibiotics in medicated feed in pig farming. *Pharmacoepidemiol Drug Safety* **13**: 323–331.
- Benson DA, Boguski MS, Lipman DJ, Ostell J, Ouellette BF, Rapp BA *et al.* (1999). GenBank. *Nucleic Acids Res* **27**: 12–17.
- Beymer R. (2000). *Boundary Waters Canoe Area – Western Region*, 6th edn. Wilderness Press: Berkeley, CA.
- Chee-Sanford JC, Aminov RI, Krapac IJ, Garrigues-Jeanjean N, Mackie RI. (2001). Occurrence and diversity of tetracycline resistance genes in lagoons and groundwater underlying two swine production facilities. *Appl Environ Microbiol* **67**: 1494–1502.
- Chopra I, Roberts M. (2001). Tetracycline antibiotics: mode of action, applications, molecular biology, and epidemiology of bacterial resistance. *Microbiol Mol Biol Rev* **65**: 232–260.
- D'Costa VM, McGrann KM, Hughes DW, Wright GD. (2006). Sampling the antibiotic resistome. *Science* **311**: 374–377.
- Davelos AL, Kinkel LL, Samac DA. (2004). Spatial variation in frequency and intensity of antibiotic interactions among Streptomycetes from prairie soil. *Appl Environ Microbiol* **70**: 1051–1058.
- De Liguoro M, Cibir V, Capolongo F, Halling-Sorensen B, Montesissa C. (2003). Use of oxytetracycline and tylosin in intensive calf farming: evaluation of transfer to manure and soil. *Chemosphere* **52**: 203–212.
- Edwards U, Rogall T, Blocker H, Emde M, Botzger EC. (1989). Isolation and direct complete nucleotide determination of entire genes. Characterization of a gene coding for 16S ribosomal RNA. *Nucleic Acids Res* **17**: 7843–7853.

- Ferber D. (2003). Antibiotic resistance – WHO advises kicking the livestock antibiotic habit. *Science* **301**: 1027.
- Guardabassi L, Dalsgaard A. (2004). Occurrence, structure, and mobility of Tn1546-like elements in environmental isolates of vancomycin-resistant enterococci. *Appl Environ Microbiol* **70**: 984–990.
- Hayes JR, English LL, Carr LE, Wagner DD, Joseph SW. (2004). Multiple-antibiotic resistance of *Enterococcus* spp. isolated from commercial poultry production environments. *Appl Environ Microbiol* **70**: 6005–6011.
- Jackson CR, Fedorka-Cray PJ, Barrett JB, Ladely SR. (2004). Effects of tylosin use on erythromycin resistance in *Enterococci* isolated from swine. *Appl Environ Microbiol* **70**: 4205–4210.
- Lane DJ. (1991). 16S/23S rRNA sequencing. In: Stackebrandt E, Goodfellow M (eds). *Nucleic Acid Techniques in Bacterial Systematics*. John Wiley & Sons: New York, pp 115–175.
- Levy SB. (1978). Emergence of antibiotic-resistant bacteria in the intestinal flora of farm inhabitants. *J Infect Dis* **137**: 689–690.
- Levy SB, Marshall B. (2004). Antibacterial resistance worldwide: causes, challenges and responses. *Nat Med* **10**: S122–S129.
- Lipsitch M, Singer RS, Levin BR. (2002). Antibiotics in agriculture: when is it time to close the barn door? *Proc Natl Acad Sci USA* **99**: 5752–5754.
- Livermore DM. (2003). Bacterial resistance: origins, epidemiology, and impact. *Clin Infect Dis* **36**: S11–S23.
- Minnesota Pollution Control Agency. (2000). Revised feedlot rules at a glance, Minnesota Rules Chapter 7020. Minnesota Pollution Control Agency, St Paul, MN.
- Muyzer G, de Waal EC, Uitterlinden AG. (1993). Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl Environ Microbiol* **59**: 695–700.
- Ng L-K, Martin I, Alfa M, Mulvey M. (2001). Multiplex PCR for the detection of tetracycline resistant genes. *Mol Cell Probes* **15**: 209–215.
- Onan LJ, LaPara TM. (2003). Tylosin-resistant bacteria cultivated from agricultural soil. *FEMS Microbiol Lett* **220**: 15–20.
- Ontario Parks. (2006). Quetico Provincial Park management plan review terms of service. [www.ontarioparks.com](http://www.ontarioparks.com).
- Phillips I, Casewell M, Cox T, De Groot B, Friis C, Jones R *et al.* (2004). Does the use of antibiotics in food animals pose a risk to human health? A critical review of published data. *J Antimicrob Chemother* **53**: 28–52.
- Ranjard L, Poly F, Lata JC, Mougel C, Thioulouse J, Nazaret S. (2001). Characterization of bacterial and fungal soil communities by automated ribosomal intergenic spacer analysis fingerprints: biological and methodological variability. *Appl Environ Microbiol* **67**: 4479–4487.
- Roberts MC. (2005). Update on acquired tetracycline resistance genes. *FEMS Microbiol Lett* **245**: 195–203.
- Rooklidge SJ. (2004). Environmental antimicrobial contamination from terraccumulation and diffuse pollution pathways. *Sci Tot Environ* **325**: 1–13.
- Sengelov G, Agero Y, Halling-Sorensen B, Baloda SB, Andersen JS, Jensen LB. (2003). Bacterial antibiotic resistance levels in Danish farmland as a result of treatment with pig manure slurry. *Environ Int* **28**: 587–595.
- Sørum M, Johnsen PJ, Aasnes B, Rosvoll T, Kruse H, Sundsfjord A *et al.* (2006). Prevalence, persistence, and molecular characterization of glycopeptide-resistant *Enterococci* in Norwegian poultry and poultry farmers 3 to 8 years after the ban on avoparcin. *Appl Environ Microbiol* **72**: 516–521.
- Torsvik V, Goksøyr J, Daae FL. (1990). High diversity in DNA of soil bacteria. *Appl Environ Microbiol* **56**: 782–787.
- United States Environment Protection Agency. (2004). *Risk assessment evaluation for concentrated animal feeding operations* EPA 600/R-04/042. National Environmental Publications Internet Site.
- Wegener HC. (2003). Antibiotics in animal feed and their role in resistance development. *Curr Opin Microbiol* **6**: 439–445.
- Whittle G, Hund BD, Shoemaker NB, Salyers AA. (2001). Characterization of the 13-kilobase *ermF* region of the *Bacteroides* conjugative transposons CTnDOT. *Appl Environ Microbiol* **67**: 3488–3495.
- Yang W, Moore IF, Koteva KP, Bareich DC, Hughes DW, Wright GD. (2004). TetX is a flavin-dependent mono-oxygenase conferring resistance to tetracycline antibiotics. *J Biol Chem* **279**: 52346–52352.