ORIGINAL ARTICLE

[FeFe]-hydrogenase in Yellowstone National Park: evidence for dispersal limitation and phylogenetic niche conservatism

Eric S Boyd¹, Trinity L Hamilton¹, John R Spear², Matthew Lavin³ and John W Peters¹

¹Department of Chemistry and Biochemistry and the Astrobiology Biogeocatalysis Research Center, Montana State University, Bozeman, MT, USA; ²Division of Environmental Science and Engineering, Colorado School of Mines, Golden, CO, USA and ³Department of Plant Sciences, Montana State University, Bozeman, MT, USA

Hydrogen (H₂) has an important role in the anaerobic degradation of organic carbon and is the basis for many syntrophic interactions that commonly occur in microbial communities. Little is known, however, with regard to the biotic and/or abiotic factors that control the distribution and phylogenetic diversity of organisms which produce H₂ in microbial communities. In this study, we examined the [FeFe]-hydrogenase gene (*hydA*) as a proxy for fermentative bacterial H₂ production along physical and chemical gradients in various geothermal springs in Yellowstone National Park (YNP), WY, USA. The distribution of *hydA* in YNP geothermal springs was constrained by pH to environments co-inhabited by oxygenic phototrophs and to environments predicted to have low inputs of abiotic H₂. The individual HydA assemblages from YNP springs were more closely related when compared with randomly assembled communities, which suggests ecological filtering. Model selection approaches revealed that geographic distance was the best explanatory variable to predict the phylogenetic relatedness of HydA communities. This evinces the dispersal limitation imposed by the geothermal spring environment on HydA phylogenetic diversity even at small spatial scales. pH differences between sites is the second highest ranked explanatory variable of HydA phylogenetic relatedness, which suggests that the ecology related to pH imposes strong phylogenetic niche conservatism. Collectively, these results indicate that pH has imposed strong niche conservatism on fermentative bacteria and that, within a narrow pH realm, YNP springs are dispersal limited with respect to fermentative bacterial communities.

The ISME Journal (2010) 4, 1485–1495; doi:10.1038/ismej.2010.76; published online 10 June 2010

Subject Category: microbial population and community ecology

Keywords: dispersal limitation; [FeFe]-hydrogenase; HydA; niche conservatism; pH; Yellowstone National Park

Introduction

Hydrogen (H₂) is ubiquitous in the biosphere and has the highest energy content per unit mass of all the naturally occurring fuels. High concentrations of H₂ have been reported in a variety of environments that include the deep subsurface (Stevens and McKinley, 1995), hydrothermal vents (Jannasch and Mottl, 1985), salt-evaporation ponds (Hoehler *et al.*, 2001) and a range of terrestrial geothermal features (Conrad *et al.*, 1985; Inskeep and McDermott, 2005; Spear *et al.*, 2005). H₂ is an important source

of energy for the maintenance and growth of microbial populations (Wolin, 1982; Morita, 2000) and has recently been postulated to be the primary fuel supporting a number of microbial assemblages (Stevens and McKinley, 1995; Chapelle *et al.*, 2002), including those inhabiting geothermal spring environments (Conrad *et al.*, 1985; Spear *et al.*, 2005).

H₂ can be produced by both geological and biological processes. Geologically, the production of H₂ occurs through the hydrolysis of reduced meteoric water when in contact with iron-bearing basalts (Apps and van de Kamp, 1993), through the reduction of water during serpentinization reactions in the presence of ultramafic minerals such as olivine (Oze and Sharma, 2007), and through the radiolysis of water (Lin *et al.*, 2005). Basalt-catalyzed H₂ production occurs readily in the presence of reduced acidic fluids (Stevens and McKinley, 2000), whereas H₂ production from serpentinization reactions is more prevalent at alkaline pH (Oze and

Correspondence: M Lavin, Department of Plant Sciences, 308 Plant BioScience Building, Montana State University, Bozeman, MT 59717, USA.

E-mail: mlavin@montana.edu or JW Peters, Department of Chemistry and Biochemistry, 103 Chemistry Research Building, Montana State University, Bozeman, MT 59717, USA.

E-mail: John.peters@chemistry.montana.edu

Received 28 January 2010; revised 20 April 2010; accepted 23 April 2010; published online 10 June 2010

Sharma, 2005; Cardace and Hoehler, 2009). Thus, geological H_2 production through serpentinization can be expected to predominate in alkaline environments rich in ultramafic minerals, whereas geological H_2 production by hydrolysis can be expected to be predominant in acidic and basalt-rich environments. Radiolytic production of H_2 is likely an important abiotic source of H_2 in subsurface environments rich in radioactive minerals (Lin *et al.*, 2005).

Biologically, [FeFe]- and [NiFe]-hydrogenase enzymes catalyze the transfer of excess electrons generated during fermentative or photosynthetic metabolism to protons to form H_2 (Vignais *et al.*, 2001). The H_2 evolution activity of [FeFe]-hydrogenase is 10- to 100-fold greater than that of the [NiFe]-hydrogenase (Frey, 2002), an observation that is consistent with the physiological role of most [NiFe]-hydrogenase in H_2 uptake (Vignais *et al.*, 2001). Recently, PCR primers for the specific amplification of the large subunit of the [FeFe]-hydrogenase gene (*hydA*) were developed to examine the diversity of *hydA* as a proxy for fermentative bacteria in a H_2 -generating saline microbial-mat system (Boyd *et al.*, 2009b) and in a H_2 -generating anaerobic bioreactor (Xing *et al.*, 2008). The results of these studies, as well as those from a number of metagenomic investigations that include the deep subsurface (Chivian *et al.*, 2008) and termite hindgut environments (Warnecke *et al.*, 2007), collectively indicate a widespread distribution and diversity of [FeFe]-hydrogenase in nature. However, our understanding of the environmental constraints that control the distribution and diversity of [FeFe]-hydrogenase-mediated H_2 production in natural systems is confounded by the tight coupling between H_2 production and consumption and by the abiological reactions that contribute to H_2 production and consumption in natural systems. In this study, we describe the distribution and phylogenetic diversity of [FeFe]-hydrogenase along physical and geochemical gradients in Yellowstone National Park (YNP), WY, USA. The results suggest that pH imposes strong phylogenetic niche conservatism (Harvey and Pagel, 1991; Wiens, 2004) and that the YNP geothermal complex imposes dispersal limitation (Hubbell, 2001) on fermentative bacterial communities, as assessed using *hydA*-deduced amino acid sequences as a functional marker.

Materials and methods

Sample collection

Approximately 500 mg samples of microbial mat or sediment were collected from 65 locations in YNP, including 16 geothermal spring locations in the Witch's Creek area of the Heart Lake Geyser Basin (HG), 12 locations in and around Imperial Geyser (IG) (Lower Geyser Basin), 21 locations in the 100 Springs Plain area of the Norris Geyser Basin (NG)

and 15 locations in the thermal field at the north end of Nymph Lake (NL) in July 2007. All samples were collected aseptically, placed into sterile microcentrifuge tubes, and were flash frozen on-site using an ethanol and dry ice slurry. Samples were kept on dry ice during transport to the laboratory, where they were kept at -80°C until they were further processed. The field sites were initially chosen to sample the range of environmental pH and temperature combinations commonly encountered in YNP springs across a variety of geographical locations. pH and temperature were determined on-site at all of the 65 sites with a model 59002-00 Cole-Parmer temperature compensated pH meter (Vernon Hills, IL, USA). Temperature was confirmed using an alcohol thermometer.

Chemical analyses

Nine sites were chosen for more detailed chemical and biological examination based on the detection of *hydA* (see below) together with the pH and temperature combination of the sites. Concentrations of ferrous iron (Fe^{2+}) and total sulfide (S^{2-}) were quantified on-site with a Hach DR/2000 spectrophotometer (Hach Company, Loveland, CO, USA), and Hach ferrozine pillows and sulfide reagents, respectively. For both Fe^{2+} and S^{2-} determinations, water samples were filtered ($0.22\ \mu\text{m}$) before addition of reagents. A separate 10 ml sample of water was filtered ($0.22\ \mu\text{m}$) into sterile tubes and was immediately frozen on dry ice for use in determining the concentrations of nitrate (NO_3^-), nitrite (NO_2^-), ammonia (NH_4^+), and phosphate (PO_4^{3-}) using a SEALQuAAtro (West Sussex, UK), calibrated daily with freshly prepared standards. Salinity was determined using a YSI model 33 S-C-T meter (Yellow Springs Instrument Company, Yellow Springs, OH, USA).

DNA extraction and PCR amplification of *hydA*

Approximately 100 mg of microbial mat or sediment was subjected to community genomic DNA extraction, purification, and quantification as earlier described (Boyd *et al.*, 2007). All DNA extracts were screened for the presence of 16S rRNA genes to ensure the presence of PCR-amplifiable DNA by using 10 ng of DNA, and primers bacterial 1070F (5'-ATGGCTGTCGTCAGCT-3') and universal 1492R (5'-GGTTACCTTGTTACGACTT-3') (Boyd *et al.*, 2009a). Approximately 500 bp fragments of *hydA* were PCR-amplified from 10 ng of environmental genomic DNA as template and primer pair FeFe-272F and FeFe-427R using previously established reagent concentrations and reaction conditions (Boyd *et al.*, 2009b). Twenty-nine of the 65 sites yielded amplicons of the correct size and out of those 9 were selected for further sequence analysis. These nine sites were chosen to sample the range of pH (1.90–9.84) and temperature combinations

(~32–65 °C) in YNP where *hydA* was detected. For each of these nine sites, new samples of microbial mat were sampled in July 2008 such that a more comprehensive assessment of geochemical data could be collected concurrently. The physical, chemical, and biological attributes of each of these sites reflect data derived from samples collected during July 2008. Community DNA extraction and quantification was performed as described above. *hydA* was amplified in triplicate and equal volumes of each replicate amplification were pooled. Pooled amplicons were purified using the Wizard PCR Preps DNA purification system (Promega, Madison, WI, USA), quantified using the Low DNA Mass Ladder (Invitrogen, Carlsbad, CA, USA), cloned using the pGem-T Easy Vector System (Promega), and sequenced using the M13F-M13R primer pair as described previously (Boyd *et al.*, 2009a).

chlL/bchL primer design and PCR amplification of chlL/bchL

Protochlorophyllide oxidoreductase subunit L genes sequences encoded by *chlL/bchL* were compiled from the GenBank database using tBLASTn with the *Anabaena variabilis* str. ATCC 29413 *ChlL* sequence as bait. The corresponding putative *chlL/bchL* sequences were imported into MEGA4 (version 4.0.1), translated in frame, and the inferred amino-acid sequences were aligned using the ClustalW application (Gonnet substitution matrix, default parameters) within the MEGA4 program. Putative ChlL/BchL sequences were screened for the presence of conserved signature catalytic residues as defined previously (Burke *et al.*, 1993). Sequences that lacked these domains were discarded without further consideration, resulting in 94 sequences for use in identifying consensus regions. Reverse translation of aligned ChlL/BchL sequences facilitated the design of primers containing minimal degeneracy.

Degenerate PCR primers corresponding to positions 54–60 (QIGCDPKHAD) and positions 167–174 (NGFDALFA) in the *Anabaena variabilis* str. ATCC 29413 *ChlL* sequence (YP_322845) were selected. Primer specificity and PCR condition optimization was performed in reactions containing genomic DNA from positive amplification control cultures of *Synechococcus* sp. JA-3-3Ab, *Roseiflexus* sp. Rs-1 and using genomic DNA from *Thermatoga maritima* MSB8, *Azotobacter vinelandii* and *Methanobacterium thermoautotrophicum* str. Delta H as negative amplification controls. Primers *chlL/bchL*-54F (5'-CARATYGGHTGYCAYCCNAARCAYGA-3', where H=A, C, or T; Y=C or T; N=A, C, G, or T; R=A or G; 768-fold degeneracy) and *chlL/bchL*-167R (5'-AAYGRYTTYGAYDSBWTHTTYGC-3', where M=A or C; S=C or G; B=C, G or T; 4608-fold degeneracy) were used to amplify a ~340 bp fragment of *chlL/bchL*. Fifty microliter PCRs contained ~10 ng of genomic DNA as template, 2 mM MgCl₂ (Invitrogen), 200 μM of each deoxyribonucleotide

triphosphate (Promega), 1 μM of both the forward and reverse primer, 200 nM molecular-grade bovine serum albumin (Roche, Indianapolis, IN, USA) and 0.25 U Taq DNA Polymerase (Invitrogen) in 1 × PCR buffer (Invitrogen). Step-down PCR conditions included an initial 4 min denaturation at 94 °C, followed by 4 cycles of denaturation at 94 °C (1 min), annealing at 64, 62, 60 and 58 °C (1 min), and primer extension at 72 °C (1.5 min). The conditions of the final 30 cycles consisted of denaturation at 94 °C (1 min), annealing at 56.5 °C (1 min) and primer extension at 72 °C (1.5 min), with a final extension step at 72 °C (20 min).

Phylogenetic analyses

hydA sequences were translated using the translate tool available on the Expasy proteomics server (<http://www.expasy.ch/tools/dna.html>). ClustalX (version 2.0.8) (Larkin *et al.*, 2007) was used to align inferred amino-acid sequences using the Gonnet 250 protein weight matrix with a pairwise alignment gap opening penalty of 13 and gap extension penalty of 0.05. ClustalX was also used to create a pairwise sequence identity matrix. The pairwise sequence identity matrix was then used in the program DOTUR (Schloss and Handelsman, 2005) to identify and group operational taxonomic units and to perform rarefaction analyses.

The phylogenetic position of putative HydA was assessed using MrBayes (version 3.1) (Huelsenbeck and Ronquist, 2001). Tree topologies were sampled every 500 generations for 2×10^6 generations using the Whelton and Goldman (WAG) evolutionary model with fixed amino-acid frequencies and gamma-shaped rate variation with a proportion of invariable sites as recommended by ProtTest (Abascal *et al.*, 2005). The *Saccharomyces cerevisiae* 'Narf' protein, a distantly related homolog of HydA, served as the outgroup in the phylogenetic analysis. A consensus phylogeny was generated from 2000 trees sampled at stationarity (standard deviation between split frequencies <0.03). The HydA phylogram was rate-smoothed using the non-parametric rate smoothing approach implemented in version 1.71 of the program r8s (Sanderson, 2002).

Community ecology

Phylocom (version 4.0.1) (Webb *et al.*, 2008) was used to calculate various metrics of phylogenetic community structure, including Rao's quadratic entropy (D_p), the net relatedness index (NRI) and the nearest taxon index (NTI) using the non-parametric-rate smoothed Bayesian chronogram. D_p represents an abundance-weighted measure of the pairwise differences in branch length of sequences in a community and is thus a metric describing phylogenetic diversity. A higher D_p index for an assemblage is indicative of higher phylogenetic diversity relative to the total *HydA* pool. The

NRI metric is a measure of tree-wide phylogenetic clustering of sequences from a given community, whereas the NTI metric is a standardized measure of the phylogenetic distance to the nearest taxon for each taxon within a community and is more sensitive to terminal (branch tip) clustering (Webb, 2000). Increasingly positive NRI and NTI scores indicate that co-occurring species are more phylogenetically related than expected by chance (phylogenetic clustering). On the contrary, increasingly negative NRI and NTI scores indicate that co-occurring species are less phylogenetically related than expected by chance (phylogenetic overdispersion) (Webb, 2000). Statistical confidence in NRI and NTI values was assessed by comparison with randomly generated phylogenies (null communities) using the 'phylogeny shuffle' model in Phylocom.

A Rao phylogenetic community distance matrix for all pairwise comparisons of sampled communities was constructed from site abundance data and an NRPS *HydA* cladogram using Phylocom. Euclidean distance matrices derived from the nine environmental variables listed in Table 1 as well as geographic distances (GDs) and elevation differences between sites were constructed using the base package within R (version 2.10.1) (R Development Core Team, 2010). Environmental parameters that were below the analytical detection limit were given a value of 0 for the purpose of this study. With Rao phylogenetic distance as the response variable, Model selection through Akaike Information Criteria adjusted for small sample size (AICc), Mantel regressions, and principle coordinate ordinations (PCO) were performed using the R packages Ecodist (Goslee and Urban, 2007), pgirmess (version 1.4.3) (<http://pagesperso-orange.fr/giraudoux/>), vegan (<http://vegan.r-forge.r-project.org/>), and labdsv (<http://ecology.msu.montana.edu/labdsv/R>). Traditional Mantel regressions were performed on the data because of the inherent redundancy of the environmental and phylogenetic distance data, whereas Model selection and regression was performed to more closely evaluate the relative performance of individual and combina-

tions of parameters in describing the phylogenetic data. We utilized PCO analysis to visually inspect *HydA* community phylogenetic relationships, as arrayed on two-dimensional plots, for patterns of clustering. We considered the model with the lowest AICc value to be the best and evaluated the relative plausibility of each model by examining differences between the AICc value for the best model and values for every other model (ΔAICc) (Johnson and Omland, 2004). Models with $\Delta\text{AICc} < 2$ were considered strongly supported by the data, and models with $\Delta\text{AICc} > 10$ and/or Mantel significance (*P*) values > 0.05 can be considered to have essentially no support from the data.

Results

The geothermal springs selected for this study represent the broad range of temperature (25–93 °C) and pH (1.90–9.84) combinations that are commonly encountered in YNP springs (Figure 1, Table 1). In addition, the range of measured chemical parameters was broad and generally reflects the ranges of concentrations that are encountered in YNP springs. Of the 65 mat or sediment samples collected in the present study, 29 yielded *hydA* PCR products of the predicted size. These 29 putative *hydA* amplicons clustered within a defined pH and temperature range (Figure 1a), indicating that the distribution of *hydA* in YNP is non-random. *hydA* was not identified in any spring with temperature greater than 65 °C. At this temperature, *hydA* was restricted to environments with alkaline pH (pH > 8.9). In contrast, the upper temperature limit for the detection of *hydA* in acidic environments (pH < 3.0) was 36 °C, indicating that the upper temperature limit for *hydA* in YNP springs is pH-dependent.

Bacteriochlorophyll biosynthesis genes *bchL/chlL* were detected in 41 of the 65 mat or sediment samples collected from YNP springs in the present study. Like *hydA*, the putative *bchL/chlL* amplicons

Table 1 Geographical, physical and chemical^a data for nine selected environments in Yellowstone National Park

Site ^b (GPS)	Elevation (m)	pH	Temp. (°C)	Salinity (%)	S ²⁻ ^a (μM)	Fe ²⁺ ^a (μM)	NO ₂ ⁻ (nM)	NO ₃ ⁻ ^a (nM)	NH ₄ ⁺ ^a (μM)	PO ₄ ³⁻ (μM)
NL1 (44.75192, -110.72856)	2289	1.90	32.0	2.6	BD	30.1	268.6	BD	26.7	238.5
NG (44.73303, -110.70922)	2278	3.30	36.0	2.3	BD	6.8	35.0	912.9	87.8	16419.4
NL2 (within 10 m of NL1)	2289	5.52	43.0	0.5	1.6	5.3	497.9	21.4	2.4	158.6
IG1 (44.53183, -110.87592)	2238	5.80	45.0	0.5	4.7	16.4	51.1	63.7	70.8	0.2
IG2 (within 10 m of IG1)	2238	6.75	55.0	0.9	BD	BD	276.4	432.9	8.2	1843.5
IG3 (within 10 m of IG1)	2238	7.13	33.0	0.6	1.3	BD	94.3	49.4	9.5	0.6
IG4 (within 10 m of IG1)	2238	7.23	45.0	0.8	0.0	BD	97.2	73.2	6.5	1.3
HL1 (44.30478, -110.52331)	2344	8.88	65.0	3.1	2.3	0.2	196.1	143.7	3.8	0.5
HL2 (44.30436, -110.52283)	2344	9.84	40.0	1.9	2.2	0.1	62.2	116.6	3.2	0.5

Abbreviations: BD, below detection; HL, Heart Lake Geyser Basin; IG, Imperial Geyser within the Lower Geyser Basin; NG, Norris Geyser Basin; NL: Nymph Lake.

^aMethod detection limits for S²⁻ = ~150 nM; Fe²⁺ = ~180 nM; NO₃⁻ = ~7 nM.

^bNL, Nymph Lake.

Figure 1 Distribution of *hydA* (a) and *bchL* (b) in sediment and mat sampled from 65 geothermal springs from four geographic locations in Yellowstone National Park as a function of spring water pH and temperature. Red squares denote environments where amplicons were detected and blue triangles denote environments where amplicons were not detected.

clustered within a defined pH and temperature range (Figure 1b). In springs with alkaline pH (pH > 9.0), the upper temperature limit for the detection of *bchL/chlL* genes was 70 °C. On the contrary, the upper temperature limit for the detection of *bchL/chlL* genes at acidic pH (< 3.0) was ~50 °C, indicating that the upper temperature limit for *bchL/chlL* in YNP springs is also pH-dependent. *bchL/chlL* genes were detected in 27 of the 29 environments where *hydA* was detected, with the exceptions being the samples collected from springs with low pH (< 3.0) and low temperature (< 30 °C) (Figure 1a). In contrast, *bchL/chlL* genes were detected in 13 environments in YNP where *hydA* was not detected.

Of the 29 environments that yielded putative *hydA* amplicons, nine were chosen for further analysis on the basis of their geochemistry and geographic location in order to span the broad geochemical and geographical spectrum available in YNP (Table 1). A total of 160 *hydA* clones were sequenced (Supplementary Table 1) from these nine environments such that the average depth of sequence coverage, as indicated by the predicted number of phylotypes in each environment by rarefaction analysis, was approximately the same (data not shown). Rao's phylogenetic diversity index (D_p), a β -diversity metric which incorporates abundance weights for branch lengths associated with each assemblage, varied significantly with environmental pH ($R^2 = 0.85$, $P < 0.01$, second-order polynomial (SOP)). The highest D_p indices were observed in *HydA* assemblages sampled from slightly acidic to circumneutral springs whereas the lowest indices were observed in assemblages sampled from alkaline environments, indicating higher diversity in assemblages sampled from acidic environments. No other measured environmental variable exhibited significant correlation with D_p .

Mantel regression and Model selection approaches evaluated using the difference in Akaike Information Criteria ($\Delta AICc$) indicated that the *HydA* phylogeny shows evidence for both ecological and geographic structure. Both statistical approaches indicated that GD between sites was the best

Table 2 Phylogenetic diversity metrics computed for each of the nine assemblages

Plot	Phylogenetic diversity metrics				
	D_p	NRI	P-value	NTI	P-value
NL1	0.37	16.13	<0.01	0.23	0.44
NG	0.75	-0.20	0.53	1.88	0.03
NL2	0.60	6.80	<0.01	1.17	0.12
IG1	0.60	7.28	<0.01	1.82	0.03
IG2	0.60	2.61	0.02	1.25	0.11
IG3	0.50	9.62	<0.01	3.24	<0.01
IG4	0.62	5.20	<0.01	2.13	<0.01
HL1	0.04	8.47	<0.01	3.47	<0.01
HL2	0.03	6.48	<0.01	3.21	<0.01

Abbreviations: D_p , Rao's quadratic entropy; HL, Heart Lake Geyser Basin; IG, Imperial Geyser within the Lower Geyser Basin; NG, Norris Geyser Basin; NL, Nymph Lake; NRI, net relatedness index; NTI, nearest taxon index.

*Statistical confidence (P-value) for NRI and NTI metrics was determined by comparison with randomly generated phylogenies.

individual explanatory variable for predicting the phylogenetic relatedness of *HydA* assemblages ($\Delta AICc = 1.05$; Mantel $R^2 = 0.45$, $P < 0.01$) (Table 2). Importantly, pH differences between sites was also a strong individual explanatory variable for predicting the phylogenetic relatedness of *HydA* assemblages ($\Delta AICc = 6.48$; Mantel $R^2 = 0.36$, $P < 0.01$). Differences in site elevation and salinity also ranked as important explanatory variables of *HydA* phylogenetic relatedness by Mantel regression approaches, but not in model selection approaches. Site elevation was autocorrelated with GD (adj $R^2 = 0.83$, $P < 0.01$), and is thus of negligible importance in predicting *HydA* phylogenetic relatedness. Importantly, other models comprising a single explanatory variable had no statistical support from the data ($\Delta AICc > 10.0$ and/or Mantel $P > 0.05$) in predicting the relatedness of *HydA* assemblages.

Models incorporating various combinations of GD and pH distance with additional explanatory variables were often stronger predictors of the phylogenetic relatedness of *HydA* assemblages than single variable models. The combined GD and pH distance model was the best explanatory model

Figure 2 Scatter plots of *HydA* Rao phylogenetic distance between sites as a function of between-site geographic distance, pH differences, salinity differences, and phosphate differences. Mantel R^2 and associated P -values are reported for each plot. Environmental characteristics associated with each environment are described in Table 1.

for predicting *HydA* phylogenetic relatedness ($\Delta\text{AICc} = 0.00$; Mantel $R^2 = 0.51$, $P < 0.01$) (Table 2) although models incorporating GD and PO_4^{3-} differences between sites ($\Delta\text{AICc} = 0.15$; Mantel $R^2 = 0.50$, $P < 0.01$), and models incorporating pH and PO_4^{3-} differences between sites ($\Delta\text{AICc} = 0.44$; Mantel $R^2 = 0.50$, $P < 0.01$) were also strong predictors of *HydA* phylogenetic relatedness. Furthermore, a model incorporating GD and salinity differences between sites ($\Delta\text{AICc} = 1.42$; Mantel $R^2 = 0.49$, $P < 0.01$) was also as strong of a predictor of *HydA* phylogenetic relatedness as models incorporating PO_4^{3-} differences. In contrast to GD, pH, and salinity differences between sites, PO_4^{3-} differences between sites alone was not a statistically significant predictor of *HydA* relatedness between sites (Table 2). Linear regressions of Rao among community *HydA* relatedness plotted as a function of between site GD, pH, and salinity differences all resulted in positive trending and statistically significant relationships (Figure 2). In contrast, a linear regression of Rao among community *HydA* relatedness plotted as a function of between-site PO_4^{3-} differences yielded a slightly negative trending and statistically insignificant relationship. Such a negative relationship would suggest phylogenetic over-dispersion or competitive exclusion with respect to PO_4^{3-} . Given that this negative relationship is weak, invoking competitive exclusion would be tenuous at best and PO_4^{3-} parameter is therefore discounted. Thus, models, which incorporate GD, pH, and salinity differences between sites as explanatory variables of community phylogenetic relatedness, are more strongly supported by the data.

The phylogenetic relatedness of *HydA* assemblages in relation to environmental and geographic characteristics was further examined using PCO of Rao among-community distances (Figure 3). PCO also indicated a strong relationship between the Rao among-community phylogenetic relatedness of *HydA* assemblages, GD and spring pH differences. The first axis (72.1% variance explained) was

Figure 3 PCO analysis of *hydA* deduced amino acid sequences from nine environments. Assemblages sampled from acidic environments (NL1 and NG) are denoted in orange, circumneutral environments (IG1, IG2, IG3, IG4, and NL2) in green, and alkaline environments (HL1 and HL2) in blue. IG, Imperial Geyser within the Lower Geyser Basin; HL, Heart Lake Geyser Basin; NG, Norris Geyser Basin; NL, Nymph Lake.

significantly correlated to geographic distance (adj $R^2 = 0.88$, $P < 0.01$) and to pH differences between sites (adj $R^2 = 0.53$, $P = 0.02$). The second axis (22.5% variance explained) was significantly correlated to salinity (adj $R^2 = 0.55$, $P = 0.01$). Significant correlations were not observed between PCO axes and other variables. Thus, the results of PCO analysis corroborate the model selection approaches and indicate that the phylogenetic relatedness of *HydA* assemblages is likely controlled through the interaction between site geographic and pH differences and to a lesser extent, salinity differences.

HydA assemblages sampled from YNP geothermal springs contain lower taxonomic diversity and are more likely to be phylogenetically clustered when compared with randomly generated HydA assemblages as indicated by statistical confidence in NRI and NTI (Table 3). HydA from alkaline environments exhibited both terminal (as indicated by NTI) and tree-wide (as indicated by NRI) phylogenetic clustering whereas the extent of phylogenetic clustering in assemblages sampled from acidic

Table 3 Model ranking using $\Delta AICc$ and Mantel correlation coefficients (R^2) where HydA Rao among community phylogenetic distance is the response variable

Model	$\Delta AICc$	R^2	P-value
GD, pH	0.00	0.51	<0.01
pH, PO_4^{3-}	0.15	0.50	<0.01
GD, PO_4^{3-}	0.44	0.50	<0.01
GD	1.05	0.45	<0.01
GD, salinity	1.42	0.49	<0.01
GD, NO_3^-	1.49	0.48	<0.01
pH, NH_4^+	2.16	0.47	<0.01
GD, Fe^{2+}	2.76	0.47	<0.01
GD, Temperature	2.98	0.46	<0.01
GD, NH_4^+	3.25	0.46	<0.01
GD, NO_2^-	3.33	0.46	<0.01
GD, S^{2-}	3.56	0.45	<0.01
pH, elevation	5.70	0.42	<0.01
pH, NO_3^-	6.23	0.41	<0.01
PH	6.48	0.36	<0.01
pH, Fe^{2+}	6.75	0.40	0.01
pH, salinity	7.59	0.39	0.00
pH, temperature	7.62	0.39	<0.01
pH, S^{2-}	8.60	0.37	0.01
pH, NO_2^-	8.87	0.37	<0.01
Elevation	10.29	0.29	<0.01
Salinity	15.27	0.19	0.02
PO_4^{3-}	20.91	0.05	0.29
Temperature	21.54	0.03	0.26
Fe^{2+}	22.14	0.02	0.48
NH_4^+	22.33	0.01	0.45
NO_3^-	22.51	0.01	0.69
NO_2^-	22.60	0.00	0.71
S^{2-}	22.61	0.00	0.78

Abbreviations: $\Delta AICc$, Akaike Information Criteria adjusted for small sample size; GD, geographic distance. Explanatory variables of each model are associated with the criteria by which they were ranked.

environments was variable. Phylogenetic clustering as measured by NTI was greater in HydA assemblages sampled from alkaline environments than assemblages sampled from circumneutral to acidic environments. Statistically significant and positive NRI and/or NTI for all YNP HydA assemblages suggests a role for ecological filtering, rather than competition in structuring the phylogenetic diversity of HydA in the Yellowstone geothermal complex.

The dominant HydA phyla in YNP springs reflected spring pH (Figure 4), a finding that is consistent with the results of Mantel tests, PCO analysis, and phylogenetic diversity metrics (see above) which taken together indicate an important role for pH in controlling the phylogenetic composition of HydA assemblages. HydA assemblages sampled from acidic springs (NL1, NG) were dominated by sequences affiliated with the *Thermotogae* whereas assemblages sampled from circumneutral springs (IG1, IG2, IG3, IG4, NL2) were dominated by sequences affiliated with the *Firmicutes*. The sole sequence obtained from both alkaline springs (HL1, HL2) was affiliated with the *Elusimicrobia* (candidate division Termite Group 1). The abundance of sequences affiliated with *Thermotogae* and *Elusimicrobia* in each of the nine assemblages was positively correlated with pH ($r^2 = 0.71$ and 0.58 , respectively, SOP) (data not shown). In contrast, the abundance of sequences affiliated with the *Firmicutes* was not correlated strongly to pH ($r^2 = 0.08$, SOP) but was significantly correlated to both sulfide and salinity ($r^2 = 0.85$ and 0.70 , respectively, SOP).

Discussion

The distribution and phylogenetic diversity of protein-encoding genes provides insight into the factors, both biotic and abiotic, that have constrained the evolution of those genes in a given environment. In this study, *hydA* served as a proxy for examining the factors that have influenced the ecology and evolution of fermentative bacteria

Figure 4 Phylum-level phylogenetic composition of HydA sequences from three spring types as determined by BLASTp analysis to closest cultivated representative. Spring types were defined as acidic (NL1 and NG), circumneutral (IG1, IG2, IG3, IG4, and NL2), and alkaline (HL1 and HL2). The relative abundance of sequences within each phylum for each spring type represents the percent of total sequences for that spring type. The phylum *Elusimicrobia* is also known as candidate division Termite Group 1. IG, Imperial Geyser within the Lower Geyser Basin; HL, Heart Lake Geyser Basin; NG, Norris Geyser Basin; NL, Nymph Lake.

putatively involved in H₂ production in YNP. Our results indicate that the distribution of *hydA* in YNP is non-random and is pH- dependent, suggesting a role for pH in the ecology and evolution of *hydA* in YNP. The pH-dependent distribution of *hydA* in YNP is similar to the distribution of phototrophs in YNP as evinced by the distribution of *bchL/chlL* genes where the upper temperature limit for oxygenic phototrophs in alkaline environments was observed to decrease from ~70 °C in alkaline pH 9.0 systems to ~48 °C in acidic pH 3.0 systems, a set of observations that are consistent with the qualitative trends noted in earlier studies (Cox and Shock, 2003; Spear *et al.*, 2005; Lehr *et al.*, 2007). The relationship between the presence of *hydA* and *bchL/chlL* could be explained, in part, by the temporal dynamics of phototrophic mat communities, which cycle from oxygen saturation during the day to anoxia at night (Revsbech and Ward, 1984). At night, oxygenic phototrophs ferment stored polyglucose and excrete metabolites such as acetate, butyrate, and propionate which are rapidly consumed under periods of anoxia by fermentative organisms (Anderson *et al.*, 1987). Thus, the pH-dependent distribution of *hydA* in YNP springs might reflect the distribution of oxygenic phototrophs as a result of their production and excretion of fermentable organic carbon.

hydA was not detected in any YNP spring with temperature >65 °C, suggesting that fermentative bacteria are unlikely to be an important source of H₂ in these environments. Since the upper temperature limit for photosynthesis is ~72 °C (Shock and Holland, 2007), photosynthetically derived H₂ is also an unlikely source for H₂ in these environments. However, elevated concentrations of H₂ are routinely measured in springs in YNP where temperatures exceed 65 °C (Inskeep and McDermott, 2005; Spear *et al.*, 2005). H₂ in these systems is likely of abiotic origin, derived from subsurface basalt-catalyzed water hydrolysis (Spear *et al.*, 2005) and could further support the notion of an H₂ driven 'deep hot biosphere' (Gold, 1992). Basalt-catalyzed H₂ production is sensitive to both solution pH and reaction temperature, with a near doubling of H₂ production rates with a doubling of incubation temperature (Stevens and McKinley, 2000). Similarly, the rate of basalt-catalyzed H₂ production increases with increasing water acidity (lower pH) (Stevens and McKinley, 2000). Coincidentally, *hydA* was not detected in YNP springs predicted to have elevated geological H₂ production (for example, acidic pH, >36 °C; alkaline pH, >65 °C), an observation that might reflect the unfavorable thermodynamics of bacterial organic carbon fermentation in the presence of high H₂ partial pressure (Schink, 1997). Thus, environments with elevated inputs of geological H₂ might select against bacterial fermentative metabolisms, which in turn might constrain the distribution of *hydA* in YNP. Further examination of the distribution of *hydA* across a gradient of environmental H₂

concentrations and across the physical and chemical limits for photosynthesis ('photosynthetic fringe') (Shock and Holland, 2007) in YNP will continue to provide insight into the role of biological and abiological factors in controlling the distribution of *hydA* and bacterial fermentation in YNP.

Of all of the measured environmental parameters, GD was the best predictor of HydA assemblage relatedness, suggesting dispersal limitation even across the small spatial scale sampled in the present study (maximum between-site distance is 53.3 km). These findings are consistent with a number of recent studies focused on microbial communities ([Martiny *et al.*, 2006) and references therein] (Whitaker *et al.*, 2003)), which together challenge the long-held notion that 'everything is everywhere and the environment selects' and implicate the fundamental role of geographic constraints in the assembly of microbial lineages in natural communities. For example, an examination of endemic populations of the hyperthermophilic archaeon *Sulfolobus* using multi-locus sequence analysis revealed dispersal limitation in geothermal environments across large geographical scales spanning the Northern hemisphere (Whitaker *et al.*, 2003). Similarly, Papke *et al.* (Papke *et al.*, 2003), showed geographic isolation in cyanobacterial populations sampled from geothermal springs from geographically distinct environments across the globe. More recently, Takacs-Vesbach *et al.* (Takacs-Vesbach *et al.*, 2008), identified historical patterns in the phylogenetic diversity of the *Aquificales* genus *Sulfurihydrogenibium* as revealed by the similarity of 16S rRNA gene sequences within a given volcanic caldera in YNP but not between calderas. This suggests that geographic provinces structure population diversity of *Sulfurihydrogenibium* at very small spatial scales, a finding that is consistent with the results presented herein. Thus, dispersal limitation should be considered in all investigations of the ecological factors that influence microbial community structure, even when sample sites span small spatial scales.

As expected, pH was also a strong predictor of HydA phylogenetic diversity, with the maximum diversity observed in assemblages sampled from environments with slightly acidic pH (~4–6). Thus, environments with slightly acidic pH are likely to harbor higher than average HydA functional variation, if functional variation is positively correlated with genetic variation. Increased functional variation, whether driven by biotic or abiotic processes, enables members of a community to respond differentially to changes in their environment (McCann, 2000) and has been linked to increased ecosystem function (Tilman *et al.*, 1997). In the YNP subsurface, the boiling of fluids under high pressure leads to the partitioning of various solutes and volatile gases into either the vapor phase or hot-water phase (Fournier, 2005). Vapor-dominated springs are generally acidic whereas hot-water

phase dominated springs are generally neutral to alkaline. Springs with circumneutral to slightly acidic pH (for example, pH 4–6) are the result of dilution of vapor-dominated water with meteoric fluid, the extent of which can be affected by water-table drawdown as a function of seasonal changes in precipitation and hydrology (Nordstrom *et al.*, 2005). Thus, the high species variability associated with slightly acidic springs might enable members of a community to better respond to change in both physical and chemical conditions in these environments due to seasonal hydrological and chemical changes. The end-result is a more resilient community, which is able to maintain a relatively steady evolutionary tempo in these geothermal environments.

The composition of the HydA assemblages varied substantially over the three spring types sampled in the present study, with each spring type dominated by a HydA affiliated with a different bacterial phylum. The abundance of dominant sequence lineages in each of the spring types was correlated strongly with environmental pH or salinity, a hypothesis which would be consistent with the inferred role of ecological filtering in YNP HydA assemblages as assessed by the NRI and/or NTI metrics. Previous studies have identified pH as an important driver of the distribution and phylogenetic diversity of soil and freshwater lake bacterial and fungal communities in several geographic locales (Newton *et al.*, 2007; Lauber *et al.*, 2009; Dumbrell *et al.*, 2010), whereas salinity has been suggested to represent the predominant factor controlling the global phylogenetic diversity of bacterial assemblages (Lozupone and Knight, 2007). For the first time, the results of the present study indicate a role for pH and to a lesser extent salinity as filters of the phylogenetic diversity of a functional gene as a proxy for a metabolic guild (bacterial fermentation) in natural environments. Importantly, springs in YNP are inherently low salinity environments and thus the role of salinity in controlling the phylogenetic diversity and relatedness of HydA assemblages on a larger scale may be underestimated. Further cross-system comparisons of HydA assemblages in environments that span a gradient of salinity and pH regimes will continue to provide insight into the relative role of pH and salinity in structuring HydA communities and phylogeny.

In summary, the results presented here indicate that both geographic factors such as the fragmented nature of hot spring systems and ecological factors such as spring pH, acting in concert, explain phylogenetically structured HydA communities in YNP springs. Our observations indicate that spring pH has imposed strong phylogenetic niche conservatism on HydA assemblages in YNP springs. Such niche conservatism, [that is, the tendency of lineages to maintain their ancestral ecological niche (Wiens, 2004)] signifies that the constituents of a HydA bacterial community are likely to inherit their

pH affinity from their ancestors and furthermore pass on this same pH predilection to their descendants. This also suggests restricted gene flow across pH extremes. Within a narrow pH realm, the detection of dispersal limitation signifies not just that geographically proximal communities are more closely related phylogenetically but that local adaptation could be limited to isolated and persistent local communities, following the arguments of Hubbell (2001). These results are consistent with earlier examinations of species endemism in geothermal environments (Papke *et al.*, 2003; Whitaker *et al.*, 2003; Takacs-Vesbach *et al.*, 2008) and show endemism of bacterial lineages at small spatial scales. In addition to the role of pH and GD, the results presented here allude to the importance of interspecies or interguild relationships (that is, phototrophs and fermentative bacteria) in structuring the genetic diversity of communities. Further examination of HydA phylogenetic diversity of samples collected from locations that span even smaller geographical and geochemical scales in combination with high resolution genetic analysis of multiple loci will continue to provide insight into the relative roles of dispersal limitation, niche conservatism, and interspecies interactions that structure fermentative bacterial communities and microbial communities in general. A further extension of the present study would be to examine the phylogenetic structure of phenotypic traits, such as inferred protein stability in the presence of gradients of salt and temperature, in order to determine which traits facilitate ecological filtering among geochemically- and geographically-disparate environments. Such studies will provide insight into our understanding of the evolutionary forces that shape the phylogenetic structure of geothermal spring microbial communities and how this relates to function at the enzyme, population, or ecosystem level.

Acknowledgements

This work was supported by grants from the US Air Force Office of Scientific Research to JWP and JRS and by the NASA Astrobiology Institute (JWP). ESB is graciously supported by a NASA Astrobiology Institute Postdoctoral Fellowship and TLH was supported by a NSF-Integrated Graduate Educational and Research and Training fellowship. We are grateful to Christie Hendrix in the Yellowstone Center for Resources for permitting the research activity and for support in all facets of the research.

References

- Abascal F, Zardoya R, Posada D. (2005). ProtTest: selection of best-fit models of protein evolution. *Bioinformatics* **21**: 2104–2105.
- Anderson KL, Tayne TA, Ward DM. (1987). Formation and fate of fermentation products in hot spring cyanobacterial mats. *Appl Environ Microbiol* **53**: 2343–2352.

- Apps JA, van de Kamp PC. (1993). Energy gases of abiogenic origin in the Earth's crust. The future of energy gases. *US Geol Survey* **1570**: 81–132.
- Boyd ES, Jackson RA, Encarnacion G, Zahn JA, Beard T, Leavitt WD. *et al.* (2007). Isolation, characterization, and ecology of sulfur-respiring *Crenarchaea* inhabiting acid-sulfate-chloride geothermal springs in Yellowstone National Park. *Appl Environ Microbiol* **73**: 6669–6677.
- Boyd ES, Leavitt WD, Geesey GG. (2009a). CO₂ uptake and fixation by a thermoacidophilic microbial community attached to precipitated sulfur in a geothermal spring. *Appl Environ Microbiol* **75**: 4289–4296.
- Boyd ES, Spear JR, Peters JW. (2009b). [FeFe]-hydrogenase genetic diversity provides insight into molecular adaptation in a saline microbial mat community. *Appl Environ Microbiol* **75**: 4620–4623.
- Burke DH, Hearst JE, Sidow A. (1993). Early evolution of photosynthesis: clues from nitrogenase and chlorophyll iron proteins. *Proc Natl Acad Sci USA* **90**: 7134–7138.
- Cardace D, Hoehler TM. (2009). Serpentinizing fluids craft microbial habitat. *Northeastern Naturalist* **16**: 272–284.
- Chapelle FH, O'Neill K, Bradley PM, Methé BA, Ciufo SA, Knobel LL *et al.* (2002). A hydrogen-based subsurface microbial community dominated by methanogens. *Nature* **415**: 312–315.
- Chivian D, Brodie EL, Alm EJ, Culley DE, Dehal PS, DeSantis TZ *et al.* (2008). Environmental genomics reveals a single-species ecosystem deep within Earth. *Science* **322**: 275–278.
- Conrad R, Bonjour F, Aragno M. (1985). Aerobic and anaerobic microbial consumption of hydrogen in geothermal spring water. *FEMS Microbiol Lett* **29**: 201–205.
- Cox AD, Shock EL. (2003). American Geophysical Union: San Francisco, abstract number 2003AGUFM.B41D0927C.
- Dumbrell AJ, Nelson M, Helgason T, Dytham C, Fitter AH. (2010). Relative roles of niche and neutral processes in structuring a soil microbial community. *ISME J* **4**: 337–345.
- Fournier RO. (2005). Geochemistry and dynamics of the Yellowstone National Park Hydrothermal System. In: Inskeep WP, McDermott TR (eds). *Geothermal Biology and Geochemistry in Yellowstone National Park*. Montana State University: Bozeman, pp 143–162.
- Frey M. (2002). Hydrogenases: hydrogen-activating enzymes. *Chembiochem* **3**: 153–160.
- Gold T. (1992). The deep, hot biosphere. *Proc Natl Acad Sci USA* **89**: 6045–6049.
- Goslee SC, Urban DL. (2007). The ecodist package for dissimilarity-based analysis of ecological data. *J Stat Softw* **22**: 1–19.
- Harvey PH, Pagel MD. (1991). *The Comparative Method in Evolutionary Biology*. Oxford University Press: Oxford, UK.
- Hoehler TM, Bebout BM, Des Marais DJ. (2001). The role of microbial mats in the production of reduced gases on the early Earth. *Nature* **412**: 324–327.
- Hubbell SP. (2001). *The Unified Neutral Theory of Biodiversity and Biogeography*. Princeton University Press: Princeton, NJ.
- Huelsenbeck JP, Ronquist R. (2001). MRBAYES: Bayesian inference of phylogeny. *Bioinformatics* **17**: 754–755.
- Inskeep WP, McDermott TR. (2005). Geomicrobiology of acid-sulfate-chloride springs in Yellowstone National Park. In: Inskeep WP, McDermott TR (eds). *Geothermal Biology and Geochemistry in Yellowstone National Park*. Montana State University: Bozeman, pp 143–162.
- Jannasch HW, Mottl MJ. (1985). Geomicrobiology of deep-sea hydrothermal vents. *Science* **229**: 717–725.
- Johnson JB, Omeland KS. (2004). Model selection in ecology and evolution. *Trends Ecol Evol* **19**: 101–108.
- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H *et al.* (2007). Clustal W and Clustal X version 2.0. *Bioinformatics* **23**: 2947–2948.
- Laubert CL, Hamady M, Knight R, Fierer N. (2009). Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Appl Environ Microbiol* **75**: 5111–5120.
- Lehr CR, Frank SD, Norris TB, D'Imperio S, Kalinin AV, Toplin JA *et al.* (2007). Cyanidia (Cyanidiales) population diversity and dynamics in an acid-sulfate-chloride spring in Yellowstone National Park. *J Phycol* **43**: 3–14.
- Lin L-H, Slater GF, Sherwood Lollar B, Lacrampe-Couloume G, Onstott TC. (2005). The yield and isotopic composition of radiolytic H₂, a potential energy source for the deep subsurface biosphere. *Geochim Cosmochim Acta* **69**: 893–903.
- Lozupone CA, Knight R. (2007). Global patterns in bacterial diversity. *Proc Natl Acad Sci USA* **104**: 11436–11440.
- Martiny JBH, Bohannan BJM, Brown JH, Colwell RK, Fuhrman JA, Green JL *et al.* (2006). Microbial biogeography: putting microorganisms on the map. *Nat Rev Microbiol* **4**: 102–112.
- McCann KS. (2000). The diversity–stability debate. *Nature* **405**: 228–233.
- Morita RY. (2000). Is H₂ the universal energy source for long-term survival? *Microbiol Ecol* **38**: 307–320.
- Newton RJ, Jones SE, Helmus MR, McMahon KD. (2007). Phylogenetic ecology of the freshwater *Actinobacteria* acI lineage. *Appl Environ Microbiol* **73**: 7169–7176.
- Nordstrom DK, Ball JW, McCleskey RB. (2005). Ground water to surface water: chemistry of thermal outflows in Yellowstone National Park. In: Inskeep WP, McDermott TR (eds). *Geothermal Biology and Geochemistry in Yellowstone National Park*. Montana State University: Bozeman, pp 143–162.
- Oze C, Sharma M. (2005). Have olivine, will gas: serpentinization and the abiogenic production of methane on Mars. *Geophys Res Lett* **32**: L10203.
- Oze C, Sharma M. (2007). Serpentinization and the inorganic synthesis of H₂ in planetary surfaces. *Icarus* **186**: 557–561.
- Papke RT, Ramsing NB, Bateson MB, Ward DM. (2003). Geographical isolation in hot spring cyanobacteria. *Environ Microbiol* **5**: 650–659.
- R Development Core Team (2010). R Foundation for Statistical Computing: Vienna, Austria. <http://www.r-project.org/contributors.html>.
- Revsbech NP, Ward DM. (1984). Microelectrode studies of interstitial water chemistry and photosynthetic activity in a hot spring microbial mat. *Appl Environ Microbiol* **48**: 270–275.
- Sanderson MJ. (2002). Estimating absolute rates of molecular evolution and divergence times: a penalized likelihood approach. *Mol Biol Evol* **19**: 101–109.
- Schink B. (1997). Energetics of syntrophic cooperation in methanogenic degradation. *Microbiol Mol Biol Rev* **61**: 262–280.

- Schloss PD, Handelsman J. (2005). Introducing DOTUR, a computer program for defining operational taxonomic units and estimating species richness. *Appl Environ Microbiol* **71**: 1501–1506.
- Shock EL, Holland ME. (2007). Quantitative habitability. *Astrobiology* **7**: 839–851.
- Spear JR, Walker JJ, McCollom TM, Pace NR. (2005). Hydrogen and bioenergetics in the Yellowstone geothermal ecosystem. *Proc Natl Acad Sci USA* **102**: 2555–2560.
- Stevens TO, McKinley JP. (1995). Lithoautotrophic microbial ecosystems in deep basalt aquifers. *Science* **270**: 450–454.
- Stevens TO, McKinley JP. (2000). Abiotic controls on H₂ production from basalt-water reactions and implications for aquifer biogeochemistry. *Environ Sci Technol* **34**: 826–831.
- Takacs-Vesbach C, Mitchell K, Jackson-Weaver O, Reysenbach A-L. (2008). Volcanic calderas delineate biogeographic provinces among Yellowstone thermophiles. *Environ Microbiol* **10**: 1681–1689.
- Tilman D, Knops J, Wedin D, Reich P, Ritchie M, Siemann E. (1997). The influence of functional diversity and composition on ecosystem processes. *Science* **277**: 1300–1302.
- Vignais PM, Billoud B, Meyer J. (2001). Classification and phylogeny of hydrogenases. *FEMS Microbiol Rev* **25**: 455–501.
- Warnecke F, Luginbuhl P, Ivanova N, Ghassemian M, Richardson TH, Stege JT *et al*. (2007). Metagenomic and functional analysis of hindgut microbiota of a wood-feeding higher termite. *Nature* **450**: 560–565.
- Webb CO. (2000). Exploring the phylogenetic structure of ecological communities: an example for rain forest trees. *Am Naturalist* **156**: 145–155.
- Webb CO, Ackerly DD, Kembel SW. (2008). Phylocom: software for the analysis of phylogenetic community structure and character evolution. *Bioinformatics* **24**: 2098–2100.
- Whitaker RJ, Grogan DW, Taylor JW. (2003). Geographic barriers isolate endemic populations of hyperthermophilic Archaea. *Science* **301**: 976–978.
- Wiens JJ. (2004). Speciation and ecology revisited: phylogenetic niche conservatism and the origin of species. *Evolution* **58**: 193–197.
- Wolin MJ. (1982) In: Bull AT, Slater JH (eds). *Microbial Interactions and Communities*, Vol. 1. Academic Press: London, England, pp 323–356.
- Xing D, Ren N, Rittmann BE. (2008). Genetic diversity of hydrogen-producing bacteria in an acidophilic ethanol-H₂-coproducing system, analyzed using the [Fe]-hydrogenase gene. *Appl Environ Microbiol* **74**: 1232–1239.

Supplementary Information accompanies the paper on The ISME Journal website (<http://www.nature.com/ismej>)