

## ORIGINAL ARTICLE

## Functional responses of methanogenic archaea to syntrophic growth

Christopher B Walker<sup>1,2</sup>, Alyssa M Redding-Johanson<sup>3</sup>, Edward E Baidoo<sup>3</sup>, Lara Rajeev<sup>3</sup>, Zhili He<sup>4</sup>, Erik L Hendrickson<sup>5</sup>, Marcin P Joachimiak<sup>3</sup>, Sergey Stoliar<sup>1</sup>, Adam P Arkin<sup>3</sup>, John A Leigh<sup>5</sup>, Jizhong Zhou<sup>4</sup>, Jay D Keasling<sup>3</sup>, Aindrila Mukhopadhyay<sup>3</sup> and David A Stahl<sup>1</sup>

<sup>1</sup>Department of Civil and Environmental Engineering, University of Washington, Seattle, WA, USA;

<sup>2</sup>Geosyntec Consultants, Seattle, WA, USA; <sup>3</sup>Physical Biosciences Division, Lawrence Berkeley National

Laboratory, Berkeley, CA, USA; <sup>4</sup>Department of Botany and Microbiology, University of Oklahoma,

Norman, OK, USA and <sup>5</sup>Department of Microbiology, University of Washington, Seattle, WA, USA

***Methanococcus maripaludis* grown syntrophically with *Desulfovibrio vulgaris* was compared with *M. maripaludis* monocultures grown under hydrogen limitation using transcriptional, proteomic and metabolite analyses. These measurements indicate a decrease in transcript abundance for energy-consuming biosynthetic functions in syntrophically grown *M. maripaludis*, with an increase in transcript abundance for genes involved in the energy-generating central pathway for methanogenesis. Compared with growth in monoculture under hydrogen limitation, the response of paralogous genes, such as those coding for hydrogenases, often diverged, with transcripts of one variant increasing in relative abundance, whereas the other was little changed or significantly decreased in abundance. A common theme was an apparent increase in transcripts for functions using H<sub>2</sub> directly as reductant, versus those using the reduced deazaflavin (coenzyme F<sub>420</sub>). The greater importance of direct reduction by H<sub>2</sub> was supported by improved syntrophic growth of a deletion mutant in an F<sub>420</sub>-dependent dehydrogenase of *M. maripaludis*. These data suggest that paralogous genes enable the methanogen to adapt to changing substrate availability, sustaining it under environmental conditions that are often near the thermodynamic threshold for growth. Additionally, the discovery of interspecies alanine transfer adds another metabolic dimension to this environmentally relevant mutualism.**

The ISME Journal (2012) 6, 2045–2055; doi:10.1038/ismej.2012.60; published online 28 June 2012

**Subject Category:** microbe–microbe and microbe–host interactions

**Keywords:** methanogen; syntrophy; coculture; alanine utilization; sulfate-reducing bacteria

## Introduction

Virtually, all methane released to the biosphere, estimated to be approximately 1 billion tons per year, are produced by methanogenic archaea living in a close association with other anaerobic microorganisms (Thauer and Shima, 2008). When electron acceptors, such as sulfate, nitrate, Mn(IV) or Fe(III), are absent or present at low concentrations, organic carbon in anoxic environments is converted to CO<sub>2</sub> and methane via a microbial food web consisting of anaerobic bacteria, microeukaryotes, syntrophic bacteria and methanogens. Anaerobic bacteria are principally responsible for the hydrolysis of biopolymers and, in association with syntrophic bacteria, ferment organic carbon to acetic acid, CO<sub>2</sub>

and hydrogen. At the low hydrogen partial pressures, typical of most anoxic habitats (<10 Pa), these are the major fermentation products and the principle substrates for methanogens. Thus, methanogens rely on microorganisms that occupy the lower tiers of this anaerobic food web for their primary growth substrates. In turn, this food-web functions optimally only when the hydrogen concentrations are maintained at a low level by methanogenic activity. At higher hydrogen concentrations, many fermentation reactions do not yield sufficient energy to support growth. Transfer of the reduced metabolic byproducts, here mainly in the form of hydrogen, from lower trophic levels to the metabolically specialized methanogens is generically termed ‘interspecies electron/hydrogen transfer’. Thus, there is a close cooperativity and interdependency between organisms producing hydrogen (sometimes referred to as proton-reducing syntrophs) and hydrogenotrophic methanogens.

We have examined the genetic and metabolic basis of this common interdependency using a

Correspondence: A Mukhopadhyay, Physical Biosciences Division, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, MS 978, Berkeley, CA 94720, USA

E-mail: amukhopadhyay@lbl.gov

Received 21 November 2011; revised 9 May 2012; accepted 9 May 2012; published online 28 June 2012

model system composed of *Desulfovibrio vulgaris* growing in syntrophic association with the hydrogenotrophic methanogen *Methanococcus maripaludis* (Stolyar *et al.*, 2007; Walker *et al.*, 2009; Hillesland and Stahl, 2010). The pairing with *D. vulgaris*, a representative sulfate-reducing bacterium, provides a model system well suited for the study of this type of microbial mutualism, as both organisms have sequenced genomes (Heidelberg *et al.*, 2004; Hendrickson *et al.*, 2004) and established genetic systems (Moore and Leigh, 2005; Keller *et al.*, 2009). We previously used this pairing to identify an alternative electron-transfer system used by *D. vulgaris* when growing syntrophically (Walker *et al.*, 2009). In the current study, we sought to characterize *M. maripaludis* grown under both syntrophic and hydrogen-limited conditions.

*M. maripaludis* utilizes a very narrow range of substrates for carbon and energy (primarily CO<sub>2</sub>, hydrogen and formate). However, genomic and biochemical studies of this well-characterized methanogen reveal a large number of isofunctional genes, suggesting the potential for adaptive responses to conditions not found in the standard laboratory cultures. Previous studies (Hendrickson *et al.*, 2007) examined the transcriptional response of *M. maripaludis* grown under hydrogen limitation, a proxy condition for its natural environment, where hydrogen concentrations are generally <10 Pa (Thauer *et al.*, 2008). That study revealed hydrogen limitation greatly increased the expression of genes encoding enzymes of the methanogenic pathway that reduce or oxidize the electron-carrying deazaflavin coenzyme F<sub>420</sub>, primarily in response to hydrogen limitation, not growth rate. Although the low hydrogen concentration used in that study more closely reflected environmental conditions, it did not capture the syntrophic lifestyle that sustains most environmental populations of methanogens. In order to better approximate the natural condition, we characterized *M. maripaludis* growing in

syntrophic association with a *Desulfovibrio* in chemostats on a lactate-based medium. In the absence of sulfate, the two organisms are obligately coupled: the *Desulfovibrio* is dependent upon *M. maripaludis* to maintain the low H<sub>2</sub> concentration required for lactate fermentation and *M. maripaludis* is dependent upon the *Desulfovibrio* as a source of hydrogen. Here we present a system-level study of *M. maripaludis* physiology as compared between syntrophic growth versus growth under hydrogen limitation.

## Materials and methods

### Strains

Transcriptional and proteomic analyses were performed using *M. maripaludis* S2 and *D. vulgaris* Hildenborough. Additionally, seven mutant strains (all described in previous investigations) of *M. maripaludis* were used during phenotypic growth comparisons. Details for all strains are shown in Table 1.

### Biomass production

Three biological replicates of cocultures and hydrogen-limited *M. maripaludis* monocultures were grown in chemostats as previously described (Haydock *et al.*, 2004; Walker *et al.*, 2009). Briefly, cocultures were grown on 30 mM lactate in previously described coculture medium (CCM) (Walker *et al.*, 2009). *M. maripaludis* monocultures were grown in Bioflo 110 bioreactors (1.3 l vessel capacity; New Brunswick Scientific Co., Edison, NJ, USA) on modified CCM containing 30 mM acetate instead of lactate. Sodium sulfide was replaced in the medium with sparging (approximately 13 ml min<sup>-1</sup> of a 1% H<sub>2</sub>S:N<sub>2</sub> mixture) of hydrogen sulfide gas. Chemostat setup and the medium and gas delivery systems were identical to a previously described system (Haydock *et al.*, 2004).

**Table 1** Summary of strains used during investigation

Strain	Mutant	Gene(s)	Description	Reference
<i>Methanococcus maripaludis</i> strains				
S2	NA	NA	Wild-type strain	Jones <i>et al.</i> , 1983
Mm1145	<i>AfruA</i>	Mmp1382	Selenocysteine-containing F <sub>420</sub> -reducing hydrogenase deletion mutant	Hendrickson and Leigh, 2008
Mm1183	<i>AfrcA</i>	Mmp0820	Cysteine-containing F <sub>420</sub> -reducing hydrogenase deletion mutant	Hendrickson and Leigh, 2008
Mm1184	<i>AfrcAAfruA</i>	Mmp1382 Mmp0820	Double mutant of <i>fruA</i> and <i>frcA</i>	Hendrickson and Leigh, 2008
Mm1020	<i>AmtD</i>	Mmp0372	F <sub>420</sub> -dependent methylenetetrahydromethanopterin dehydrogenase deletion mutant	Hendrickson and Leigh, 2008
Mm1097	<i>Ahmd</i>	Mmp0127	Hydrogen-dependent methylenetetrahydromethanopterin dehydrogenase deletion mutant	Hendrickson and Leigh, 2008
Mm1002	<i>Aald</i>	Mmp1513	Alanine dehydrogenase deletion mutant	Moore and Leigh, 2005
Mm1018	<i>ΔagcS</i>	Mmp1511	Sodium:alanine symporter deletion mutant	Moore and Leigh, 2005
<i>Desulfovibrio vulgaris</i> strain				
Hildenborough (= ATCC 29 579)	NA	NA	Wild-type strain	ATCC

Abbreviations: NA, not applicable.

A 1-ml glycerol stock of previously grown coculture or monoculture was used to inoculate 100 ml of CCM (amended with sulfate for monocultures) in a 200 ml serum vial. Cultures were incubated in the dark at 37 °C with a shaking speed of 250 r.p.m. When the cultures reached an OD<sub>600</sub> of 0.25–0.30, they were transferred to a 3-l FairMenTec chemostat (FairMenTec GmbH, Göttingen, Germany) filled with 2 l of CCM. The chemostat was run in batch mode at 37 °C with a stirring speed of 250 r.p.m. N<sub>2</sub>:CO<sub>2</sub> (90:10) was flushed through a sterile cotton plug before entering the headspace of the reactor. Flow rate was maintained at 0.25 ml min<sup>-1</sup> using an Alicat Scientific mass controller (MC-20SCCM-D; Alicat Scientific, Tucson, AZ, USA). Headspace concentrations of CH<sub>4</sub>, CO<sub>2</sub>, H<sub>2</sub>, H<sub>2</sub>S, O<sub>2</sub> and N<sub>2</sub> were monitored at 30 min intervals using a Hiden Analytical QIC-20 mass spectrometer (Hiden Analytical, Warrington, UK). For coculture experiments, a culture medium flow rate of 1.3 ml min<sup>-1</sup> (25.6 h retention time) was initiated when the OD<sub>600</sub> reached 0.325–0.350. The chemostat was assumed to be at steady state when the variance of OD<sub>600</sub> readings was <10% over three retention periods. Cultures were harvested through ice-chilled sterile stainless steel tubing connected to the chemostat culture medium exhaust line. Falcon tubes (50 ml) were stored in an anoxic chamber and pre-chilled on ice prior to harvesting. The tubes were centrifuged at 4 °C for 15 min at 3220 g, after which the supernatant was poured off and the tubes were immediately frozen at -80 °C. Samples were shipped overnight on dry ice.

#### Transcriptional analysis

Whole-transcriptome microarrays containing 70-mer probes for each of the 1722 *M. maripaludis* S2 open reading frames were spotted on UltraGAPS glass slides (Corning Life Sciences, Corning, NY, USA) using a BioRobotics Microgrid II arrayer (Genomic Solutions, Ann Arbor, MI, USA). Each slide also contained 70-mer probes for 3531 *D. vulgaris* Hildenborough open reading frames. Each slide had duplicate spots for each open reading frame. Each biological replicate was hybridized to at least three slides. Thus, each log<sub>2</sub> expression level described here was obtained using triplicate biological replicates/slides, for each of which there were duplicate on-chip technical replicates.

RNA isolation, quantification and transcription were performed as previously described (Walker *et al.*, 2009) and fluorescently labeled using Cy5-dUTPs. Labeled RNA was compared against Cy3-dUTP-labeled genomic DNA and computational analyses were performed (Walker *et al.*, 2009). For log<sub>2</sub> R calculations, R = signal intensity ratio of coculture/monoculture. Z-values were calculated as described in Mukhopadhyay *et al.* (2006). In this study, genes with absolute Z-score values >1 were considered significantly changed. Gene-expression

data are available at Microbes Online ([www.microbesonline.org](http://www.microbesonline.org)) and under GEO reference GSE30831, GSM764979 and GSM764978.

#### Protein preparation, labeling and analysis

Cell pellets from biological triplicates were pooled into 1 ml of lysis buffer (500 mM triethylammonium bicarbonate with 4 M urea, pH = 7). The samples were lysed by sonication on ice for 3 min of active time with pulses of 5 s on and 10 s off. Because of high amounts of DNA associated with the *M. maripaludis* samples, 40 µl of RQ-1 DNase (1 U µl<sup>-1</sup>; Promega, Madison, WI, USA) was added to each sample and 1 M MgCl<sub>2</sub> was added to a final concentration of 10 mM. The samples were set at 37 °C for 1.5 h, after which the samples were clarified by centrifugation at 10 000 g for 30 min at 4 °C. Protein concentration was determined using the Biorad Assay (Biorad, Hercules, CA, USA). To provide a control for the coculture, equal amounts of protein were mixed from *M. maripaludis* and *D. vulgaris*, referred to hereafter as synthetic blend. The iTRAQ (isobaric tags for relative and absolute quantitation) labeling was carried out as previously described (Redding *et al.*, 2006) and samples were labeled as follows: tag114–coculture; tag115–synthetic blend; tag116–synthetic blend; and tag117–coculture. This strategy allowed two technical replicates for each of the samples, increasing confidence for proteins that show differential expression. All the collected data were processed using Protein Pilot (AB SCIEX, Framingham, MA, USA). The data is computed as ratios, so as to provide relative change. Because the actual ratio of *D. vulgaris* to *M. maripaludis* in the coculture was not 1:1, normalization of the fold change (coculture versus synthetic blend) was done based on the cellular ratio data obtained using DAPI (4'6-diamidino-2-phenylindole)-stained cell counts, which indicated that the true ratio was 80:20 (Walker *et al.*, 2009). For *D. vulgaris* proteins, the fold change was normalized by taking the log<sub>2</sub> ratio of the coculture to synthetic blend and subtracting the log<sub>2</sub> value of 80/50, whereas the *M. maripaludis* fold change was normalized by subtracting the log<sub>2</sub> value of 20/50. A detailed explanation of this normalization strategy and the complete proteomics data set are provided in the Supplementary Information and Supplementary Table S1. Normalized protein log<sub>2</sub> ratios >|2| were considered to be significantly changed.

#### Metabolite analysis

Cell pellets and supernatants from 50 ml cultures were collected (in triplicate) for cocultures and monocultures at OD<sub>600</sub> ~0.3. Metabolites were extracted from both pellets and supernatant using methanol extraction and lyophilized as described previously (Baidoo *et al.*, 2008). Briefly, 2 ml

pre-chilled methanol was used to extract metabolites from pellet, whereas 1 ml supernatant was extracted with 1 ml methanol. A total of 30 ml water (deionized high-performance liquid chromatography grade) was added to extracts and frozen in liquid N<sub>2</sub> and lyophilized. All reagents used were high-performance liquid chromatography grade. Capillary electrophoresis (CE) separation and mass spectrometry (MS) analysis were conducted as previously described (Baidoo *et al.*, 2008). The standard concentration curves using commercially available standards of pyruvate and alanine (Sigma, St Louis, MO, USA) were used to obtain absolute levels of target metabolites in pmol mg<sup>-1</sup> dry cell weight.

#### Phenotypic growth analyses

All the phenotypic growth assays were carried out in 17 ml Hungate tubes (BellCo Glass, Vineland, NJ, USA) equipped with rubber stoppers and crimp-tops. Cultures were incubated at 37 °C in the dark with a 300 r.p.m. shaking speed. Each tube contained 8 ml of CCM amended with 30 mM of electron donor (lactate or pyruvate). The headspace contained an overpressure of 18 kPa of N<sub>2</sub>:CO<sub>2</sub> (80:20). Dilution series out to 10<sup>-8</sup> were initiated from 1 ml glycerol stocks of each *Methanococcus* mutant and *D. vulgaris*. *Methanococcus* cultures were grown in CCM lacking lactate and amended with 5 mM acetate and 250 kPa overpressure of H<sub>2</sub>:CO<sub>2</sub> (80:20). *Desulfovibrio* cultures were grown in CCM amended with 30 mM sulfate. Cocultures were established by combining 0.5 ml of exponentially growing *Methanococcus* and 1 ml of exponentially growing *Desulfovibrio* from the highest dilutions. Cocultures were transferred (1% v/v) three times to ensure dilution of any residual sulfate/acetate or H<sub>2</sub> before inoculating triplicate tubes for growth experiments. Tubes were monitored for growth using OD<sub>600</sub> readings blanked against uninoculated medium.

## Results

The most general change associated with syntrophic growth was a decrease in transcripts for energy-consuming biosynthetic functions (for example, pyruvate oxidoreductase (Por), acetyl-CoA decarboxylase/synthase and energy-conserving hydrogenase B (Ehb), see Figure 1 for enzymatic reaction depictions) and an increase in transcripts in the energy-generating methanogenesis pathway (see Table 2 and Supplementary Tables S2 and S4; Supplementary Figure S1) plots transcriptional changes according to function as defined by clusters of orthologous groups). Figure 1 illustrates the differential expression of transcripts observed in coculture compared with hydrogen-limited *M. maripaludis* monocultures. However, compared with growth in monoculture under hydrogen


limitation, transcript levels for isofunctional genes often diverged, with transcripts of one variant significantly increasing, whereas the other was little changed or significantly decreased in their relative abundances. A common feature was an apparent increase in transcripts for functions using H<sub>2</sub> directly as reductant, versus those using the reduced deazaflavin (coenzyme F<sub>420</sub>). In several cases, the proteins were confidently identified (Table 3 and Figure 2) and the corresponding changes generally corroborated the observations at the transcript level. The greater importance of direct reduction by H<sub>2</sub> was supported by improved syntrophic growth of a deletion mutant in an F<sub>420</sub>-dependent dehydrogenase of *M. maripaludis* (described in more detail below). Metabolite, transcript and proteomic analyses also pointed to a unique, although undefined, role for alanine utilization within this syntrophic coupling.

#### Carbon assimilation and methanogenesis

A number of genes associated with carbon assimilation and biosynthesis displayed differences in transcript abundance during syntrophic growth. Most notably, lower transcript abundance was observed for many genes coding for subunits of Ehb (*ehbC*, *ehbD*, *ehbA*, *ehbN* and *ehbF*, Table 2), the energy-conserving hydrogenase proposed to be involved in CO<sub>2</sub> assimilation (Porat *et al.*, 2006), and genes coding for acetyl-CoA decarboxylase/synthase (Mmp0979-85, Supplementary Table S2) and Vor (Mmp1271-2, Supplementary Table S2), both associated with autotrophic growth. Fewer transcripts were also observed for the adenine diphosphate-forming acetyl-CoA synthetase (*acd*, Mmp0253, Supplementary Table S2), one of the two enzymes responsible for acetyl-CoA production via acetate.

Transcripts for genes encoding a formate dehydrogenase (*fdh*) (Mmp1297-8, Table 2), previously shown to be important in both monoculture (Wood *et al.*, 2003) and coculture (Stolyar *et al.*, 2007), were increased. Increased transcript abundance was accompanied by a significant increase in the protein abundance of the FdhC subunit (Figure 1, Table 3). Additionally, genes for an associated transporter and carbonic anhydrase (Mmp1299-301, Supplementary Table S2) displayed transcript increases. Despite the observed increases in both transcript and protein levels, formate was not detected in the culture medium (limit of detection, 0.1 mM).

Differential expression was observed for several of the genes coding for proteins involved either directly or indirectly in the seven steps in hydrogenotrophic methanogenesis. Transcripts of the *mch* gene (Mmp1191) and most of the *mtr* operon (Mmp1560-7, Table 2) increased slightly, as did transcripts for a single gene in the methyl coenzyme M reductase operon (*mcrC*, Mmp1556, Table 2). The small increase in the transcript levels for the *mtr* and



**Figure 1** Conceptual schematic of *M. maripaludis* and *D. vulgaris* syntrophic interaction, highlighting the central energy-generating and -consuming the pathways of the methanogen. Relative changes (Table 2) in transcript abundance during syntrophic growth are indicated by red (increase) and green (decrease) coloration. Blue coloration indicates no statistically significant change as specified in the Materials and methods section. Hydrogen-limited monocultures served as the control growth condition. Oxidation of formate to CO<sub>2</sub> and H<sub>2</sub> coupled with coenzyme F<sub>420</sub>-reduction (not depicted) is catalyzed by two alternative formate dehydrogenases, Fdh1 and Fdh2. One of two membrane-bound energy-conserving hydrogenases (Eha and Ehb) couple the chemiosmotic energy of ion gradients to H<sub>2</sub> oxidation and ferredoxin reduction. Of these, Ehb generates the low potential electron carrier used for anabolism, whereas Eha is hypothesized to function primarily in the energy-generating methanogenesis pathway, generating low potential-reducing equivalents for the reduction of CO<sub>2</sub> to formylmethanofuran (Major *et al.*, 2010). Two different formylmethanofuran dehydrogenases catalyze this first step in methanogenesis, tungsten (Fwd) and molybdenum (Fmd) forms. Transfer of the formyl group from methanofuran to methanopterin by Ftr and subsequent elimination of H<sub>2</sub>O by Mch yields methenyl-H<sub>4</sub>-methanopterin. Two different enzymes can then reduce methenyl-H<sub>4</sub>-MPT to methylene-H<sub>4</sub>MPT, one (Mtd) using H<sub>2</sub> as reductant and the other (Hmd) using reduced coenzyme F<sub>420</sub>. *M. maripaludis* has an Hmd paralog of unknown function (Mmp1716, Hmd<sub>II</sub>) that may also function in this step (Hendrickson *et al.*, 2004). Reduction of methylene-H<sub>4</sub>MPT by another F<sub>420</sub>-dependent reductase (Mer) yields methyl-H<sub>4</sub>MPT. The reduced coenzyme F<sub>420</sub> required for the formation of methyl-H<sub>4</sub>MPT by these two steps is generated by one of two alternative F<sub>420</sub>-reducing hydrogenase (Fru and Frc). The final steps to methane production are catalyzed by a methyl transferase (Mtr) and a reductase (Mcr) coupled to two forms of a F<sub>420</sub>-nonreducing hydrogenase (Vhu and Vhc). The mixed disulfide (CoM-CoB) produced by reduction of methyl coenzyme M is then reduced by one of two forms of the heterodisulfide reductase determined by the composition of the HdrA subunit (HdrA<sub>V</sub> or HdrA<sub>H</sub>). Fdh/ Hdr/Vhu/ Fwd are reported to have protein–protein interactions (Costa *et al.*, 2010). (Note: The interaction with Fwd could not be depicted here without compromising the clarity of the figure. Vhc is not part of this interaction). Other reactions include the transport of alanine (AlsT), and subsequent conversion to pyruvate via an alanine racemase (Alr) and dehydrogenase (Ald). Additional *M. maripaludis* proteins shown: pyruvate oxidoreductase, acetyl-CoA decarboxylase/synthase. The *D. vulgaris* metabolic pathway is based upon results as described in Walker *et al.*, 2009 and is updated to include an unspecified sodium/alanine transporter (Na<sup>+</sup>/ala sym). Other *D. vulgaris* proteins shown: lactate permease (lac per), lactate dehydrogenase (ldh), membrane-bound Coo hydrogenase (Coo), high-molecular weight cytochrome (Hmc), periplasmic hydrogenases (Hyd and Hyn), cytochrome c3 (Cyt c3) and oxidized and reduced ferredoxin (Fd).

mcr operons were not reflected in the proteomics data (Supplementary Table S3).

### Hydrogenases

The *M. maripaludis* genome encodes seven potential hydrogenases involved in the various pathways whose expression depends upon growth and nutrient conditions. Two of the hydrogenases (the Eha and Ehb complexes) are membrane-bound and function during ferredoxin reduction and energy conversion. In contrast to decreased transcripts observed for several Ehb subunits, a small number of the genes from the energy-conserving hydrogenase complex Eha showed increased transcript

levels (Mmp1447; *ehaA*, Mmp1448; *ehaC*, Mmp1450; and *ehaD*, Mmp1451, Table 2), although no statistically significant differences were observed for the remaining Eha subunits.

Of the two isofunctional cytoplasmic F<sub>420</sub>-reducing hydrogenases (Frc and Fru), transcripts for the selenocysteine-containing F<sub>420</sub>-reducing hydrogenase (Fru, Mmp1382-5, Table 2) increased during syntrophic growth, a phenomenon also observed under hydrogen limitation when selenium is present in the culture medium (Noll *et al.*, 1999; Hendrickson *et al.*, 2007; Baidoo *et al.*, 2008). A corresponding increase in the FruA (Table 3, Figure 2) protein was also observed. Transcripts for the cysteine-containing F<sub>420</sub>-reducing hydrogenase

**Table 2** Expression ratios of selected genes in the central energy-generating and -consuming pathways of *M. maripaludis*

Gene number	ID	Coculture versus H <sub>2</sub> -limited monoculture		H <sub>2</sub> -limited versus P <sub>1</sub> -limited <sup>a</sup>	
		log <sub>2</sub> R <sup>b</sup>	Z-score	log <sub>2</sub> R <sup>b</sup>	P-value
<i>F420</i> -interacting proteins					
Mmp0058	<i>mer</i>	<i>n.a.</i>	<i>n.a.</i>	2.51 (5.7)	4.80E-11
Mmp0372	<i>mtd</i>	0.55 (1.5)	0.79	4.14 (17.6)	2.30E-10
Mmp1298	<i>fdhAI</i>	1.70 (3.2)	2.22	2.26 (4.8)	1.50E-04
Mmp0138	<i>fdhA2</i>	1.54 (2.9)	0.77	0.41 (1.3)	7.60E-03
Mmp1385	<i>fruB</i>	3.29 (9.8)	2.49	1.83 (3.6)	3.00E-09
Mmp0818	<i>frcG</i>	-1.50 (0.35)	0.79	0.16 (1.1)	7.30E-03
<i>Methanogenesis pathway proteins (energy-generating)</i>					
Mmp0127	<i>hmd</i>	-1.15 (0.45)	0.75	-0.20 (0.87)	2.60E-01
Mmp1716	<i>hmdIII</i>	1.04 (2.1)	1.43	-0.09 (0.93)	7.40E-03
Mmp1155	<i>hdrB1</i>	0.55 (1.5)	0.94	0.22 (1.2)	1.50E-02
Mmp1053	<i>hdrB2</i>	1.10 (2.1)	1.5	0.46 (1.4)	9.60E-04
Mmp1697	<i>hdrAu</i>	0.81 (1.7)	1.12	0.40 (1.3)	3.30E-03
Mmp0825	<i>hdrAv</i>	-1.09 (0.47)	1.05	0.23 (1.2)	8.60E-04
Mmp1559	<i>mcrA</i>	1.31 <sup>c</sup> (2.5)	1.95	-0.30 (0.81)	6.70E-03
Mmp1609	<i>frt</i>	0.10 (1.1)	0.18	0.42 (1.3)	6.90E-07
Mmp1448	<i>ehaA</i>	0.91 <sup>d</sup> (1.9)	1.36	0.24 (1.2)	1.20E-02
Mmp1696	<i>vhuD</i>	1.61 (3.0)	2.31	<i>n.a.</i>	<i>n.a.</i>
Mmp0822	<i>vhcG</i>	-2.12 (0.23)	1.26	0.12 (1.1)	2.30E-02
Mmp1567	<i>mtrE</i>	1.05 (2.8)	1.43	-0.65 (0.64)	4.60E-01
Mmp1247	<i>fwDD</i>	0.93 (1.9)	1.15	0.12 (1.1)	3.00E-01
<i>Anabolic pathway proteins (energy-consuming)</i>					
Mmp0984	<i>cdh</i>	-1.39 (0.38)	1.94	0.43 (1.3)	4.30E-06
Mmp1504	<i>porB</i>	-0.86 (0.55)	1.33	0.38 (1.3)	7.50E-06
Mmp1622	<i>ehbM</i>	-1.09 <sup>e</sup> (0.47)	1.46	0.66 (1.6)	5.00E-08
Mmp1512	<i>alr</i>	2.74 (6.7)	2.85	-0.73 (0.60)	3.30E-10
Mmp1513	<i>ald</i>	3.85 (14.4)	2.63	-0.55 (0.68)	5.50E-09

<sup>a</sup>Data from Hendrickson *et al.* (2007).

<sup>b</sup>Values in parantheses are the R values.

<sup>c</sup>Data available only for *mcrD* (Mmp1556).

<sup>d</sup>Mean value for *ehaACD*. Data not available for *ehaBEFGHIJKLNO* (low Z-scores).

<sup>e</sup>Mean value for *ehbBCDNFG*. Data not available for *ehbOMLKJL* (low Z-scores).

**Table 3** Normalized relative ratios for significantly changed *M. maripaludis* proteins


Gene number	ID	Description	Log <sub>2</sub> (coculture/ synthetic coculture) <sup>a</sup>
MMP1382	FruA	Selenocystein-containing coenzyme F <sub>420</sub> -reducing hydrogenase, alpha subunit	2.93
MMP0802	FrcA	Coenzyme F <sub>420</sub> -reducing hydrogenase, alpha subunit	1.92
MMP0817	FrcB	Coenzyme F <sub>420</sub> -reducing hydrogenase, beta subunit	2.09
MMP1301	FdhC	Formate dehydrogenase, alpha subunit	2.03
MMP1513	ald	Alanine dehydrogenase	3.46
MMP1302		Hypothetical protein MMP1302	2.18
MMP1156		Hypothetical protein MMP1156	2.18
MMP1161		Hypothetical protein MMP1161	2.08

<sup>a</sup>ITRAQ proteomics data normalized as described in supplementary section based on *D. vulgaris*: *M. maripaludis* ratio of 4: 1. Data average of technical replicates.

(Frc, Mmp0817-20) did not show a statistically significant change, except for a reduced transcript level for a gene coding for FrcG (Table 2). However, increases in protein abundance were measured for two of the Frc subunits (FrcA and FrcB, Table 3, Figure 2).

A second set of isofunctional enzymes, the F<sub>420</sub> non reducing hydrogenases have been proposed to

transfer electrons from H<sub>2</sub> to the heterodisulfide complex formed in the last steps of methanogenesis (Afting *et al.*, 2000). Transcript levels for genes coding for the cysteine-containing coenzyme F<sub>420</sub>-nonreducing hydrogenase were decreased (Vhc, Mmp0821-3). The heterodisulfide reductase subunit HdrA<sub>v</sub>, predicted to be within the same operon, also showed a decrease at the transcript level (Mmp0825,




**Figure 2** Quantitative proteomic data. Analysis of coculture protein samples using iTRAQ labeling and shotgun liquid chromatography–mass spectrometry methods identified 82 proteins from *M. maripaludis* (out of 207 total coculture proteins) by at least two unique, high-confidence peptides in replicate runs. Among these 82 *M. maripaludis* proteins (Supplementary Table S1), 8 exhibited significant abundance increases (Table 3). Plot shows  $\log_2$  ratios of the iTRAQ ratios of proteins from coculture versus the synthetic blend. The synthetic blend contains a 50:50 mixture of *D. vulgaris* to *M. maripaludis*, whereas the coculture is a 80:20 mixture of the same. Normalized data was used. For complete data see Supplementary Table S1.

Table 2). In contrast, genes coding for the Vhu and the HdrA<sub>U</sub> subunit of Hdr showed an increase in transcript levels (Table 2). No significant changes were measured for VhcA, VhcD (Mmp0823, Mmp0821) and Vhu (Mmp1694) at the protein level (Supplementary Table S3).

Given the significant increases of both transcript and protein abundance, the roles of the Fru/c hydrogenases were examined using the corresponding gene deletion mutants in syntrophic growth with *D. vulgaris*. The  $\Delta fruA$  and  $\Delta frcA$  mutants each exhibited diminished growth rates and maximum cell densities during syntrophic growth on lactate (Figure 3). A deletion in both genes (a  $\Delta fruA\Delta frcA$  mutant) produced highly variable maximum cell densities, but syntrophic growth still occurred, albeit at generally slower rates than in wild-type cocultures. All the three mutants demonstrated an increase in lag time prior to initiating exponential growth. In contrast, when pyruvate was the substrate for syntrophic growth, there were no differences in growth rate or maximum cell densities between the wild-type and the  $\Delta fruA$  or  $\Delta frcA$  mutants, and there was only a slight decrease in maximum cell density for the  $\Delta frcA\Delta fruA$  double mutant (Supplementary Figure S2).

Although there were no statistically significant changes in transcript or protein, abundance for either the hydrogen-dependent (Hmd, Mmp0127) or the F<sub>420</sub>-dependent (Mtd, Mmp0372) methylene-H<sub>4</sub>MPT dehydrogenases, both the  $\Delta mtd$  and  $\Delta hmd$



**Figure 3** Growth curves for wild-type and mutant *M. maripaludis* cultures on lactate. The error bars indicate s.d. of triplicate cultures. Wild type (filled squares);  $\Delta mtd$  (open triangles);  $\Delta hmd$  (closed triangles);  $\Delta fruA$  (open diamonds);  $\Delta frcA$  (closed diamonds); and  $\Delta frcA\Delta fruA$  (closed circles).

mutant strains, affected syntrophic growth. Unexpectedly, the  $\Delta mtd$  strain growing in syntrophic association on lactate had a reduced lag time and achieved a greater maximum cell density than the wild type (Figure 3). Thus, restricting this step in methanogenesis to a hydrogen-dependent enzyme (Hmd) improved overall growth performance of the coculture, possibly by enhancing *Desulfovibrio* growth by maintaining a lower concentration of H<sub>2</sub>. Although the  $\Delta hmd$  strain showed a longer lag period, there was no difference in maximum cell density. However, deletion of *hmd* (Mmp0127) may be compensated for by its paralog *hmd<sub>II</sub>* (Mmp1716). These genes responded differentially to syntrophic growth, with a two-fold increase in Mmp1716 transcripts and a reduction in Mmp0127 transcripts (Table 2). As with the  $\Delta fruA$  or  $\Delta frcA$  mutants, no observable differences in growth rate or maximum cell density were noted for either the  $\Delta mtd$  or the  $\Delta hmd$  mutant strains compared with the wild type when grown in coculture using pyruvate as carbon source (Supplementary Figure S2).


#### Alanine utilization

Transcript levels for alanine dehydrogenase (*ald*) and alanine racemase (*alr*) genes in the methanogen (both involved in alanine utilization) were greatly elevated during syntrophic growth, with corresponding increases in protein abundances in the case of *ald* (Tables 2 and 3, Figure 2). The associated alanine transport gene, annotated as a sodium:alanine symporter (*alsT*, Mmp1511), exhibited a significant downregulation at the transcript level that may have resulted from repression by the *nprR* nitrogen-regulation gene (Xia et al., 2009). *AlsT* was not identified in the proteomics data set. Previous characterization of the *agcS* (*alsT*) and *ald* deletion mutants indicated that these genes are essential when alanine is the sole source of nitrogen (Moore and Leigh 2005). However, in an

ammonia-containing lactate medium, no significant differences in growth rate or cell density were observed for either deletion mutant in coculture as compared with wild type (Figure 4). Nonetheless, because the alanine dehydrogenase and racemase transcripts were among those most highly elevated in coculture, we further examined intra- and extracellular concentrations of alanine and its conversion product, pyruvate. Concentrations were determined for both cocultures and monocultures of the  $\Delta alsT$  mutant and wild-type *M. maripaludis*. Values for cocultures reflect aggregate contributions from both *D. vulgaris* and *M. maripaludis*. Notably for both co- and monocultures, the cultures containing the mutant strain had significantly higher intra- and extracellular concentrations of both alanine and pyruvate (Table 4). Additionally, higher internal concentrations of both metabolites were observed in wild-type monocultures of *M. maripaludis* when compared with equivalent amounts of coculture cells.

## Discussion

Syntrophic growth in the *Desulfovibrio*/*Methanococcus* model community requires that the two participating organisms share the free energy



**Figure 4** Growth curves for wild-type and alanine-related mutant *M. maripaludis* cultures on lactate. The error bars indicate s.d. of triplicate cultures.

available from the fermentation of lactate. Assuming that molecular hydrogen is the primary mediator of reduced metabolite exchange, the fraction of energy available to each organism is determined primarily by  $H_2$  concentration. A low  $H_2$  concentration favors the *Desulfovibrio*, but limits the methanogen. Conversely, a high  $H_2$  concentration favors the methanogen, but limits the *Desulfovibrio*. Although during steady-state growth the  $H_2$  concentration must be maintained at a concentration that satisfies the energetic needs of both organisms, the energy need not be divided equally (Worm et al., 2011). Using the average concentrations measured during the steady-state growth experiments (4 mM lactate, 26 mM acetate, 2.5 Pa  $H_2$ , 5,100 Pa  $CO_2$  and 61 Pa  $CH_4$ ; temperature 310 K), the total free energy available from lactate fermentation to methane, acetate and  $CO_2$  ( $-82.8 \text{ kJ mol}^{-1}$ ) is not equally shared between *Desulfovibrio* ( $\sim 60 \text{ kJ mol}^{-1}$ ) and *Methanococcus* ( $\sim 20 \text{ kJ mol}^{-1}$ ). Thus, the methanogen appears to be close to the thermodynamic threshold for growth defined by the minimum increment of energy ( $-15$  to  $-20 \text{ kJ mole}^{-1}$ ) required for adenosine triphosphate synthesis (Schink and Stams, 2006). In the current study, we examined the adaptive response of the methanogen to the energetic constraints of extreme hydrogen limitation imposed by syntrophic growth.

All the transcription data were related to *Methanococcus* growing under hydrogen limitation in chemostats in metal-replete medium at the same generation time as the coculture, as previously reported by Walker et al. (2009). In that study, we showed that an alternative electron-transfer system was required for lactate oxidation by the *Desulfovibrio* when growing in syntrophic association with *Methanococcus*, but not for respiratory growth on sulfate (Walker et al., 2009). Thus, it appeared that specific enzyme systems are dedicated to *Desulfovibrio* growth under conditions of syntrophy. In the current study, we observe changes in *Methanococcus* gene expression that also appear of greater relevance to syntrophic growth. Relative to growth under hydrogen limitation in monoculture, the syntrophically grown methanogen showed increased transcription of genes in the central

**Table 4** Measured intra- and extracellular alanine and pyruvate concentrations. Alanine and pyruvate concentrations (pmol/mg of dry cells) observed for cocultures (CC) and monocultures of *M. maripaludis* (MmS) and *D. vulgaris* (DvH). Wild type (WT) or alanine transporter deficient ( $\Delta alsT$ ) strains were used

	Average					s.d.				
	$CC_{WT}$	$CC_{\Delta alsT}$	$MmS_{WT}$	$MmS_{\Delta alsT}$	$DvH_{WT}$	$CC_{WT}$	$CC_{\Delta alsT}$	$MmS_{WT}$	$MmS_{\Delta alsT}$	$DvH_{WT}$
Internal alanine	0.13	1.60	0.58	2.46	7.16	0.04	0.08	n.a.	0.74	0.72
External alanine	82.27	442.61	382.8	772.14	1395.98	58.52	43.67	n.a.	135.32	109.36
Internal pyruvate	0.18	0.85	0.22	0.34	2.53	0.03	0.02	n.a.	0.09	0.8
External pyruvate	4.74	25.00	10.43	20.32	34.49	0.36	5.38	n.a.	0.93	3.42

Abbreviation: n.a., not applicable.

pathway for methanogenesis (*mtrH*, *mcrA* and *fwdD*) and of paralogs coding primarily for different steps in hydrogen uptake (*ehaA*, *hmd<sub>II</sub>*, *fruB*, *vhuD* and *hdrA<sub>U</sub>*). Notably, transcription of paralogs previously reported to be upregulated with hydrogen limitation in monoculture (Hendrickson *et al.*, 2007) was significantly reduced in coculture (*hmd*, *hdrA<sub>V</sub>*, *vhcG* and *frcG*). Thus, different paralogs presumably provide physiological advantages at different H<sub>2</sub> concentrations. An increase in *hmd<sub>II</sub>* levels in coculture, and a decrease in the levels of its paralog *hmd*, is particularly noteworthy as *Hmd<sub>II</sub>* was reported to not function as a hydrogenase/dehydrogenase based on *in vitro* characterization of the *Methanothermobacter marburgensis* paralogs (Afting *et al.*, 2000). Because the *Hmd* hydrogenase has >20-fold higher K<sub>m</sub> for H<sub>2</sub> (0.2 mM) than the F<sub>420</sub>-reducing (NiFe)-hydrogenase (0.01 mM), and because the paralog competitively binds a Fe-binding guanylylpyridinol cofactor required by *Hmd*, it was suggested that the paralog functions primarily to store the cofactor during hydrogen limitation (Goldman *et al.*, 2009; Thauer *et al.*, 2010). However, our observations of down-regulation of *hmd* and coincident upregulation of *hmd<sub>II</sub>* during hydrogen-limited syntrophic growth now point to a more direct function in hydrogen metabolism. Direct involvement of *Hmd<sub>II</sub>* in the reduction of methenyl-H<sub>4</sub>-MPT is also supported by the unexpected improvement in syntrophic growth of the *mtd* deletion mutant relative to the wild type (Figure 3), suggesting that *Hmd<sub>II</sub>* variant is functionally relevant in the reduction of methenyl-H<sub>4</sub>-MPT at extremely low H<sub>2</sub> concentrations (1 Pa).

The upregulation of *fruB* and the corresponding increase in Fru protein abundance points to the challenge of syntrophically grown *M. maripaludis* in maintaining reduced coenzyme F<sub>420</sub> at low H<sub>2</sub> concentrations (Nolling *et al.*, 1995; Morgan *et al.*, 1997; Hendrickson *et al.*, 2007). This is also consistent with the poor growth in coculture of the *ΔfrcA* and *ΔfruA* mutant strains. An impaired capacity of these mutants to link hydrogen oxidation with the reduction of F<sub>420</sub> is the likely basis for reduced growth rate and cell densities observed in coculture, but not in monoculture (Hendrickson *et al.*, 2008). Importantly, the growth of the *ΔfrcA ΔfruA* mutant in coculture indicates that the previously suggested reverse *Hmd*-*Mtd* pathway for coenzyme F<sub>420</sub>-reduction (Hendrickson *et al.*, 2008) may be sufficient to support syntrophy, although the other unknown pathways cannot be ruled out.

The increased transcription of genes coding for Vhu and HdrA<sub>U</sub>, paralogs of the heterodisulfide reductase complex, and decrease in Vhc and HdrA<sub>V</sub> is consistent with formation of a complex between Fdh, Fwd, Vhu and HdrA<sub>U</sub> in metal-replete medium (Berghofer and Klein 1995; Costa *et al.*, 2010). Within this complex, a flavin-mediated electron bifurcation at HdrA results in reduction of both the CoM-S-S-CoB heterodisulfide and ferredoxin

required by Fwd for the first step in methanogenesis. In the presence of formate, or during hydrogen limitation, expression of *fdh* increases and the protein is incorporated into the protein complex to facilitate electron transfer from formate to Hdr (Costa *et al.*, 2010). The coordinate increase in transcription of the genes coding for all members of this complex (*fwd*, *hdrA<sub>U</sub>*, *vhu* and *fdh*) also suggests a requirement for increased expression of the complex under conditions of hydrogen limitation imposed by syntrophic growth.

The prominent increase in alanine metabolism genes highlights a potentially unexplored mechanism of syntrophic coupling. *M. maripaludis* is capable of utilizing alanine as a nitrogen source, with the dehydrogenase and transporter being essential when alanine is the sole nitrogen source, but not in the presence of ammonia (Moore and Leigh, 2005). Production of alanine by *D. vulgaris* via reduction of pyruvate (presumably by an alanine dehydrogenase) could complement H<sub>2</sub> as a mediator of interspecies electron transfer. This form of reduced product exchange would also benefit the methanogen by providing fixed carbon and nitrogen. A variety of *Methanococcus* strains have capacity to assimilate alanine (Whitman *et al.*, 1987). However, no appreciable differences in growth rates or yields were observed relative to the wild type for cocultures established with either the *M. maripaludis* *Δald* or *ΔalsT* mutants. This might mean that the cost of alanine secretion to *Desulfovibrio* is compensated for by an energy advantage provided by the methanococcus, such that changes in growth rate or yield are not discernable by the aggregate measure of optical density. *Desulfovibrio* would forfeit one adenosine triphosphate for each pyruvate not further oxidized to acetate, whereas *Methanococcus* would benefit from a reduced energy investment in the autotrophic synthesis of pyruvate. However, another feature of alanine export via a Na<sup>+</sup>/alanine symporter would be the coincident export of Na<sup>+</sup>, contributing to a sodium motive force that might be used to drive energetically unfavorable reactions (Figure 1). Although it is also possible that *D. vulgaris* uses alanine primarily as a compatible osmolyte to alleviate salt stress imposed by the brackish CCM (He *et al.*, 2010) and *Methanococcus* benefits from alanine leakage, the data are most consistent with a flux of alanine between the two species. First, transcripts of genes for autotrophic growth (*por*, *ehb* and *cdh*) are all reduced in coculture relative to H<sub>2</sub>-limited monoculture. Second, when compared with monocultures and *ΔalsT* mutant strains, the lowest concentration of internal alanine was observed for the wild-type coculture sample. Finally, the introduction of the *ΔalsT* mutant to the coculture, preventing uptake of alanine by *M. maripaludis*, resulted in a greater concentration of both external and internal alanine.

Our study also adds to a number of previous theoretical and experimental studies of the

metabolic basis of syntrophic association. Those studies primarily examine alternative mechanisms for generating hydrogen or formate from the high-potential oxidation reactions mediated by the bacterial syntroph. It is now apparent that syntrophs use a variety of mechanisms to couple exergonic reactions with energetically unfavorable oxidation of substrates such as propionate, butyrate or lactate (McInerney *et al.*, 2009; Stams and Plugge 2009; Walker *et al.*, 2009; Müller *et al.*, 2010). Fewer studies have considered both the metabolism and gene expression of syntrophically associated methanogens (Luo *et al.*, 2002; Enoki *et al.*, 2011; Worm *et al.*, 2011). However, the use of phylogenetically divergent methanogens in these studies complicates direct comparison with our results. Worm and colleagues focused on changes in expression of alternative *fdh* and hydrogenases (*hyd*) in *Methanospirillum hungatei* growing in syntrophic association with *Syntrophobacter fumaroxidans*, showing that *fdh* and *hyd* genes were transcribed in both organisms in either monoculture or coculture. Notably, although expression levels did vary with culture conditions, the *fdh* and *hyd* genes in *M. hungatei* were both transcribed in monoculture on either formate or hydrogen (Worm *et al.*, 2011). Thus, the increased transcription of *fdh* genes we observed in *M. maripaludis* during syntrophic growth may not be in direct response to production of formate by *D. vulgaris*. In other studies, proteome and transcript analysis of *Methanothermobacter thermautotrophicus* in coculture with a fatty acid oxidizing syntroph showed primarily differential expression of two variants of the methyl coenzyme M reductase (*Mcr*), one variant (*MCR I*) was preferentially expressed in coculture and the other (*MCR II*) in monoculture. However, *M. maripaludis* contains only a single variant of *Mcr*.

These reports and our study point to a tremendous variety of strategies for the metabolic coupling of different syntrophic partners. The increased transcription of paralogs feeding electrons derived from hydrogen into the methanogenesis pathway and indications for novel exchange of alanine between species observed in our study seem to be of significance to the adaptive response and habitat preference (niche) of *Methanococcus* species. Thus, there is now added impetus for studies designed to further evaluate the biochemical properties and physiological significance of the adaptive changes observed in this syntrophic pair, and more generally how different syntrophic pairings influence anaerobic food-web structure and function.

## Acknowledgements

We thank Professor Michael J McInerney (University of Oklahoma) for valuable discussion and review of an earlier draft of this paper, Dr Christopher Petzold (LBNL) for help with iTRAQ data analysis on Protein Pilot and Jason Baumohl (LBNL) for help with submitting data to

the GEO database. This work is part of ENIGMA, a Scientific Focus Area Program supported by the US Department of Energy, Office of Science, Office of Biological and Environmental Research, Genomics: GTL Foundational Science through contract DE-AC02-05CH11231 between Lawrence Berkeley National Laboratory and the US Department of Energy.

## References

- Afting C, Kremmer E, Brucker C, Hochheimer A, Thauer RK. (2000). Regulation of the synthesis of H<sub>2</sub>-forming methylenetetrahydromethanopterin dehydrogenase (Hmd) and of Hmd<sub>II</sub> and Hmd<sub>III</sub> in *Methanothermobacter marburgensis*. *Arch Microbiol* **174**: 225–232.
- Baidoo EE, Benke PI, Neuss C, Pelzing M, Kruppa G, Leary JA *et al.* (2008). Capillary electrophoresis-fourier transform ion cyclotron resonance mass spectrometry for the identification of cationic metabolites via a pH-mediated stacking-transient isotachopheric method. *Anal Chem* **80**: 3112–3122.
- Berghofer Y, Klein A. (1995). Insertional Mutations in the Hydrogenase Vhc and Frc Operons Encoding Selenium-Free Hydrogenases in *Methanococcus voltae*. *Appl Environ Microbiol* **61**: 1770–1775.
- Costa KC, Wong PM, Wang T, Lie TJ, Dodsworth JA, Swanson I *et al.* (2010). Protein complexing in a methanogen suggests electron bifurcation and electron delivery from formate to heterodisulfide reductase. *Proc Natl Acad Sci USA* **107**: 11050–11055.
- Enoki M, Shinzato N, Sato H, Nakamura K, Kamagata Y. (2011). Comparative proteomic analysis of *Methanothermobacter thermautotrophicus* DeltaH in pure culture and in co-culture with a butyrate-oxidizing bacterium. *PLoS One* **6**: e24309.
- Goldman AD, Leigh JA, Samudrala R. (2009). Comprehensive computational analysis of Hmd enzymes and paralogs in methanogenic Archaea. *BMC Evol Biol* **9**: 199.
- Haydock AK, Porat I, Whitman WB, Leigh JA. (2004). Continuous culture of *Methanococcus maripaludis* under defined nutrient conditions. *Fems Microbiol Lett* **238**: 85–91.
- He Z, Zhou A, Baidoo E, He Q, Joachimiak MP, Benke P *et al.* (2010). Global transcriptional, physiological, and metabolite analyses of the responses of *Desulfovibrio vulgaris* Hildenborough to salt adaptation. *Appl Environ Microbiol* **76**: 1574–1586.
- Heidelberg JF, Seshadri R, Haveman SA, Hemme CL, Paulsen IT, Kolonay JF *et al.* (2004). The genome sequence of the anaerobic, sulfate-reducing bacterium *Desulfovibrio vulgaris* Hildenborough. *Nat Biotechnol* **22**: 554–559.
- Hendrickson EL, Kaul R, Zhou Y, Bovee D, Chapman P, Chung J *et al.* (2004). Complete genome sequence of the genetically tractable hydrogenotrophic methanogen *Methanococcus maripaludis*. *J Bacteriol* **186**: 6956–6969.
- Hendrickson EL, Haydock AK, Moore BC, Whitman WB, Leigh JA. (2007). Functionally distinct genes regulated by hydrogen limitation and growth rate in methanogenic Archaea. *Proc Natl Acad Sci USA* **104**: 8930–8934.

- Hendrickson EL, Liu Y, Rosas-Sandoval G, Porat I, Soll D, Whitman WB *et al.* (2008). Global responses of *Methanococcus maripaludis* to specific nutrient limitations and growth rate. *J Bacteriol* **190**: 2198–2205.
- Hillesland KL, Stahl DA. (2010). Rapid evolution of stability and productivity at the origin of a microbial mutualism. *Proc Natl Acad Sci USA* **107**: 2124–2129.
- Jones WJ, Whitman WB, Fields RD, Wolfe RS. (1983). Growth and plating efficiency of methanococci on agar media. *Appl Environ Microbiol* **46**: 220–226.
- Keller KL, Bender KS, Wall JD. (2009). Development of a markerless genetic exchange system for *Desulfovibrio vulgaris* Hildenborough and its use in generating a strain with increased transformation efficiency. *Appl Environ Microbiol* **75**: 7682–7691.
- Luo HW, Zhang H, Suzuki T, Hattori S, Kamagata Y. (2002). Differential expression of methanogenesis genes of *Methanothermobacter thermoautotrophicus* (formerly *Methanobacterium thermoautotrophicum*) in pure culture and in cocultures with fatty acid-oxidizing syntrophs. *Appl Environ Microbiol* **68**: 1173–1179.
- Major TA, Liu Y, Whitman WB. (2010). Characterization of energy-conserving hydrogenase B in *Methanococcus maripaludis*. *J Bacteriol* **192**: 4022–4030.
- McInerney MJ, Sieber JR, Gunsalus RP. (2009). Syntrophy in anaerobic global carbon cycles. *Curr Opin Biotechnol* **20**: 623–632.
- Moore BC, Leigh JA. (2005). Markerless mutagenesis in *Methanococcus maripaludis* demonstrates roles for alanine dehydrogenase, alanine racemase, and alanine permease. *J Bacteriol* **187**: 972–979.
- Morgan RM, Pihl TD, Nolling J, Reeve JN. (1997). Hydrogen regulation of growth, growth yields, and methane gene transcription in *Methanobacterium thermoautotrophicum* deltaH. *J Bacteriol* **179**: 889–898.
- Mukhopadhyay A, He ZL, Alm EJ, Arkin AP, Baidoo EE, Borglin SC *et al.* (2006). Salt stress in *Desulfovibrio vulgaris* Hildenborough: An integrated genomics approach. *J Bacteriol* **188**: 4068–4078.
- Müller N, Worm P, Schink B, Stams AJM, Plugge CM. (2010). Syntrophic butyrate and propionate oxidation processes: from genomes to reaction mechanisms. *Environ Microbiol Rep* **2**: 489–499.
- Noll I, Muller S, Klein A. (1999). Transcriptional regulation of genes encoding the selenium-free [NiFe]-hydrogenases in the archaeon *Methanococcus voltae* involves positive and negative control elements. *Genetics* **152**: 1335–1341.
- Nolling J, Pihl TD, Reeve JN. (1995). Cloning, Sequencing, and Growth Phase-Dependent Transcription of the Coenzyme F<sub>420</sub>-Dependent N-5,N-10-Methylenetetrahydromethanopterin Reductase-Encoding Genes from *Methanobacterium thermoautotrophicum* Delta-H and *Methanopyrus kandleri*. *J Bacteriol* **177**: 7238–7244.
- Porat I, Kim W, Hendrickson EL, Xia QW, Zhang Y, Wang TS *et al.* (2006). Disruption of the operon encoding Ehb hydrogenase limits anabolic CO<sub>2</sub> assimilation in the archaeon *Methanococcus maripaludis*. *J of Bacteriol* **188**: 1373–1380.
- Redding AM, Mukhopadhyay A, Joyner DC, Hazen TC, Keasling JD. (2006). Study of nitrate stress in *Desulfovibrio vulgaris* Hildenborough using iTRAQ proteomics. *Brief Funct Genomics Proteomics* **5**: 133–143.
- Schink B, Stams AJM. (2006). Syntrophism Among Prokaryotes. In: Dworkin M, Falkow S, Rosenberg E, Schliefer K-H, Stackebrandt E (eds). *The Prokaryotes: An Evolving Electronic Resource for the Microbiological Community*. Springer-Verlag: New York, pp 309–335.
- Stams AJ, Plugge CM. (2009). Electron transfer in syntrophic communities of anaerobic bacteria and archaea. *Nat Rev Microbiol* **7**: 568–577.
- Stolyar S, Van Dien S, Hillesland KL, Pinel N, Lie TJ, Leigh JA *et al.* (2007). Metabolic modeling of a mutualistic microbial community. *Mol Syst Biol* **3**: 92.
- Thauer RK, Kaster AK, Seedorf H, Buckel W, Hedderich R. (2008). Methanogenic archaea: ecologically relevant differences in energy conservation. *Nat Rev Microbiol* **6**: 579–591.
- Thauer RK, Shima S. (2008). Methane as fuel for anaerobic microorganisms. *Ann N Y Acad Sci* **1125**: 158–170.
- Thauer RK, Kaster AK, Goenrich M, Schick M, Hiromoto T, Shima S. (2010). Hydrogenases from methanogenic archaea, nickel, a novel cofactor, and H<sub>2</sub> storage. *Annu Rev Biochem* **79**: 507–536.
- Walker CB, He Z, Yang ZK, Ringbauer JA Jr., He Q, Zhou J *et al.* (2009). The electron transfer system of syntrophically grown *Desulfovibrio vulgaris*. *J Bacteriol* **191**: 5793–5801.
- Whitman WB, Sohn S, Kuk S, Xing R. (1987). Role of Amino Acids and Vitamins in Nutrition of Mesophilic *Methanococcus* spp. *Appl Environ Microbiol* **53**: 2373–2378.
- Wood GE, Haydock AK, Leigh JA. (2003). Function and regulation of the formate dehydrogenase genes of the methanogenic archaeon *Methanococcus maripaludis*. *J Bacteriol* **185**: 2548–2554.
- Worm P, Stams AJ, Cheng X, Plugge CM. (2011). Growth- and substrate-dependent transcription of formate dehydrogenase and hydrogenase coding genes in *Syntrophobacter fumaroxidans* and *Methanospirillum hungatei*. *Microbiology* **157**: 280–289.
- Xia Q, Wang T, Hendrickson EL, Lie TJ, Hackett M, Leigh JA. (2009). Quantitative proteomics of nutrient limitation in the hydrogenotrophic methanogen *Methanococcus maripaludis*. *BMC Microbiol* **9**: 149.



This work is licensed under the Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplementary Information accompanies the paper on The ISME Journal website (<http://www.nature.com/ismej>)