ORIGINAL ARTICLE

Aquatic metagenomes implicate *Thaumarchaeota* in global cobalamin productionAndrew C Doxey, Daniel A Kurtz, Michael DJ Lynch, Laura A Sauder and Josh D Neufeld
Department of Biology, University of Waterloo, Waterloo, Ontario, Canada

Cobalamin (vitamin B₁₂) is a complex metabolite and essential cofactor required by many branches of life, including most eukaryotic phytoplankton. Algae and other cobalamin auxotrophs rely on environmental cobalamin supplied from a relatively small set of cobalamin-producing prokaryotic taxa. Although several *Bacteria* have been implicated in cobalamin biosynthesis and associated with algal symbiosis, the involvement of *Archaea* in cobalamin production is poorly understood, especially with respect to the *Thaumarchaeota*. Based on the detection of cobalamin synthesis genes in available thaumarchaeotal genomes, we hypothesized that *Thaumarchaeota*, which are ubiquitous and abundant in aquatic environments, have an important role in cobalamin biosynthesis within global aquatic ecosystems. To test this hypothesis, we examined cobalamin synthesis genes across sequenced thaumarchaeotal genomes and 430 metagenomes from a diverse range of marine, freshwater and hypersaline environments. Our analysis demonstrates that all available thaumarchaeotal genomes possess cobalamin synthesis genes, predominantly from the anaerobic pathway, suggesting widespread genetic capacity for cobalamin synthesis. Furthermore, although bacterial cobalamin genes dominated most surface marine metagenomes, thaumarchaeotal cobalamin genes dominated metagenomes from polar marine environments, increased with depth in marine water columns, and displayed seasonality, with increased winter abundance observed in time-series datasets (e.g., L4 surface water in the English Channel). Our results also suggest niche partitioning between thaumarchaeotal and cyanobacterial ribosomal and cobalamin synthesis genes across all metagenomic datasets analyzed. These results provide strong evidence for specific biogeographical distributions of thaumarchaeotal cobalamin genes, expanding our understanding of the global biogeochemical roles played by *Thaumarchaeota* in aquatic environments.

The ISME Journal (2015) 9, 461–471; doi:10.1038/ismej.2014.142; published online 15 August 2014

Introduction

Cobalamin (vitamin B₁₂) is an enzyme cofactor that many prokaryotic and eukaryotic species use to catalyze rearrangement-reduction or methyl transfer reactions involved primarily in amino acid synthesis (i.e., cobalamin-dependent methionine synthase), return of carbon to central metabolism via the tricarboxylic acid cycle (i.e., adenosylcobalamin-dependent methylmalonyl-CoA mutase) and synthesis of DNA (ribonucleotide reductase). Other cobalamin-dependent enzymes include ethanolamine ammonia-lyase, methylaspartate ammonia-lyase and methylaspartate mutase (reviewed in Sañudo-Wilhelmy *et al.* (2014)). Cobalamin is related to heme, chlorophyll and the F₄₂₀ coenzyme of methanogens, and contains a tetrapyrrole corrin ring system surrounding a central cobalt atom. The fifth coordinated position of Co³⁺ is occupied by a

dimethyl-benzamidazole nucleotide loop and the sixth catalytic upper ligand position is either occupied by a methyl group (i.e., methylcobalamin) or with a deoxyadenosine (i.e., adenosylcobalamin).

Because of the absolute requirement of cobalamin coenzymes by many marine bacterioplankton and most cultured phytoplankton, the supply of B vitamins (including vitamins B₁, B₇ and B₁₂) have been implicated in controlling bloom distributions and species successions in the ocean water column (Sañudo-Wilhelmy *et al.*, 2006; Bertrand *et al.*, 2007; Gobler *et al.*, 2007). In particular, widespread nutritional auxotrophy results in a complete dependence by many marine organisms on an exogenous source of cobalamin. Auxotrophic phytoplankton are thought to establish symbiotic relationships with bacterial partners to obtain this vitamin (Croft *et al.*, 2005; Grant *et al.*, 2014), providing a source of organic carbon to the prokaryotic symbionts in return. Another strategy is to develop costly alternative pathways (e.g., cobalamin-independent methionine synthase) or to scavenge organic nutrients directly from the environment, which has been suggested for the ubiquitous *Pelagibacter ubique* of the SAR11 clade (Tripp *et al.*, 2008) and for a

Correspondence: AC Doxey or JD Neufeld, Department of Biology, University of Waterloo, 200 University Avenue West, Waterloo, Ontario, Canada N2L 3G1.

E-mail: acdoxey@uwaterloo.ca or jneufeld@uwaterloo.ca

Received 2 May 2014; revised 2 July 2014; accepted 3 July 2014; published online 15 August 2014

recently discovered class of uncultured marine *Euryarchaeota* (Iverson *et al.*, 2012).

Cobalamin biosynthesis represents one of the most complex and ancient metabolic pathways known (Raux *et al.*, 1999), and this vitamin has been described as ‘nature’s most beautiful cofactor’ (Stubbe, 1994). Owing to its elaborate structure, only a limited complement of *Bacteria* and *Archaea* are known to produce cobalamin. The ‘Black Queen Hypothesis’ (i.e., in the card game ‘Hearts’, players avoid holding the queen of spades; Morris *et al.*, 2012) has been invoked to explain why cobalamin producers maintain biosynthesis of cobalamin for potential community benefit, despite the high metabolic cost for those few taxa responsible for its production (Giovannoni, 2012). Based on genome analysis, known dominant marine bacterial taxa implicated in cobalamin production include select members of the *Cyanobacteria*, *Alphaproteobacteria*, *Gammaproteobacteria* and *Bacteroidetes* (Sañudo-Wilhelmy *et al.*, 2014). These characterized cobalamin producers synthesize cobalamin by either aerobic or anaerobic pathways that involve 20 enzymatic steps from a uroporphyrinogen III precursor, sharing several homologous enzymes, but with oxygen-requiring or oxygen-sensitive steps, depending on the pathway (Moore *et al.*, 2013). The aerobic pathway involves a late-insertion of the cobalt ion into the already contracted tetrapyrrole structure, whereas the early insertion pathway can operate under anoxic conditions and incorporates cobalt prior to ring contraction.

Although genomic analyses have implicated *Archaea* in cobalamin production (Rodionov *et al.*, 2003), examples have been limited to halophilic *Crenarchaeota* of hypersaline ponds and marine methanogens from the *Euryarchaeota*. To our knowledge, no study has reported a potential cobalamin synthesis role for the *Thaumarchaeota*. This is surprising because *Thaumarchaeota* are widespread in aquatic and terrestrial environments (Francis *et al.*, 2005; Leininger *et al.*, 2006; Prosser and Nicol, 2008) and dominate marine water columns (Wuchter *et al.*, 2006). Known *Thaumarchaeota* possess autotrophic and oligotrophic metabolisms that are fueled by ammonia oxidation, and may be stimulated by organic molecules such as pyruvate (Tournai *et al.*, 2011; Stahl and de la Torre, 2012). Implicating cobalamin synthesis in thaumarchaeotal metabolism has important global biogeochemical and ecological implications, especially for the marine environment, where *Thaumarchaeota* are estimated to account for ~20% of all prokaryotic cells (Karner *et al.*, 2001).

Based on our observation of cobalamin synthesis genes within available thaumarchaeotal genomes (Hallam *et al.*, 2006; Spang *et al.*, 2010; Walker *et al.*, 2010; Blainey *et al.*, 2011; Kim *et al.*, 2011; Lebedeva *et al.*, 2013; Lloyd *et al.*, 2013; Luo *et al.*, 2014), combined with the ubiquity and abundance of these organisms, we hypothesized that *Thaumarchaeota*

represent numerically relevant marine cobalamin producers. We also predicted that cobalamin synthesis gene distributions would demonstrate biogeographical patterns, with correlations to measured physical and chemical characteristics. Leveraging available marine metagenomic libraries, we demonstrate depth-specific, latitudinal and temporal gradients of thaumarchaeotal cobalamin synthesis genes. We also observed niche partitioning of phylum-specific cobalamin synthesis, which reinforces that vitamin synthesis is a coordinated and costly keystone ecosystem function provided by individual, rather than collective, community members. Our results implicate *Thaumarchaeota* among dominant global cobalamin producers and add vitamin B₁₂ production to the roles recognized for these abundant and enigmatic marine microbial community members.

Materials and methods

Genome pathway analysis

The Integrated Microbial Genomes system (Markowitz *et al.*, 2014) was used to detect matches to all enzymes within the Kyoto Encyclopedia of Genes and Genomes (KEGG; Kanehisa *et al.*, 2008) reference pathway map for ‘porphyrin and chlorophyll metabolism’ (map00860), containing both the aerobic and anaerobic cobalamin biosynthesis pathways. The presence of each enzyme was assessed within all available archaeal genomes, and these data were retrieved to generate a presence/absence matrix. The proportion of archaeal genomes possessing each enzyme within the pathway was used to compute an archaeal ‘conservation score’. This score was then mapped onto a pathway heatmap using the KEGG Mapper tool (<http://www.genome.jp/kegg/mapper.html>).

Metagenome datasets and processing

We retrieved 430 aquatic metagenomes from 28 studies from the Community Cyberinfrastructure for Advanced Microbial Ecology Research and Analysis (CAMERA) portal on 8 January 2014 (see Supplementary Table S1 for a full list of accession numbers). Included studies spanned all available marine and freshwater metagenomes, and excluded data from engineered environments or metagenomes related to host-associated, virus-, or bacteria-specific studies. Given that polar marine *Thaumarchaeota* were recently implicated in urea-fueled nitrification (Alonso-Sáez *et al.*, 2012), we also included the specific Arctic metagenome used in this previous study (NCBI SRA accession number ERP001178).

Pipeline for detection and taxonomic classification of cobalamin biosynthesis genes

Metagenome profile Hidden Markov Models (profile-HMMs) were used for genes of the cobalamin biosynthesis pathway that were retrieved from the TIGRfam (Haft *et al.*, 2003) and Pfam

(Bateman *et al.*, 2004) databases, as based on the KEGG pathway (map00860). A total of 11 genes (*cobA_cysG* C-terminal domain (TIGR01469), *cobI_cbiL* (TIGR01467), *cobJ_cbiH* (TIGR01466), *cobM_cbiF* (TIGR01465), *cbiT_cobL* (TIGR02469), *cbiE_cobL* (TIGR02467), *cbiC_cobH* (PF02570.10), *cbiA_cobB* (TIGR00379), *cbiB_cobD* (TIGR00380), *cbiP_cobQ* (TIGR00313), *cobS* (TIGR00317)) were selected as cobalamin pathway markers for further analysis due to their broad distribution throughout the cobalamin biosynthesis pathway. For each of these 11 genes, the corresponding HMM was used to scan for homologs in all sequence reads of each metagenome, which were initially processed into open reading frames using FragGeneScan (Rho *et al.*, 2010). The program *hmmsearch* within HMMER version 3.1b1 (hmm.org) was used with default parameters and an *E*-value threshold of 1×10^{-6} . The use of profile-HMMs based on protein family alignments eliminated bias that would have otherwise been introduced by homology searches based on single sequence queries. Tools and taxonomy indices from the Krona package (Ondov *et al.*, 2011) were used to assign taxonomy to recovered hits based on their top BLAST matches in the NCBI refseq database (version 60) with an *E*-value threshold of 1×10^{-6} .

Per-taxon cobalamin gene contribution

Taxonomic frequency profiles were generated by dividing the number of cobalamin synthesis hits per taxon by the total number of hits for a given gene category. Removing samples with low overall cobalamin synthesis gene frequencies (i.e., $n < 10$) reduced potential statistical bias resulting from low sample sizes. To generate an overall measure of per-taxon cobalamin gene contributions for each sample, we combined data for the 11 cobalamin marker genes as follows. For a given dataset, *D*, the per-taxon cobalamin (*cob/cbi*) gene contribution (*D_t*) was computed as:

$$D_t = \frac{\sum_{i=1}^{11} \# \text{cobalamin gene } i \text{ hits for taxon } t}{\sum_{i=1}^{11} \# \text{cobalamin gene } i \text{ hits}}$$

Metadata, correlation and cluster analysis

Correlation analyses of numerical metadata from each metagenome compared individual physical and chemical characteristics with per-taxon cobalamin gene contributions (i.e., *D_t* from above). We analyzed metadata correlations across all individual metagenome datasets and also for pooled sample datasets for larger studies, with many samples characterized by common study-specific metadata. Standard Pearson correlations were used to calculate correlation coefficients (*r*). Maps were generated with the maps package in R (<http://cran.r-project.org/web/packages/maps/index.html>), with metagenome samples plotted by their associated latitude and

longitude metadata. Hierarchical clustering was performed using the Ward method with Manhattan distances in R.

Phylogenetic marker gene analysis

The full analysis described above was repeated using the Pfam profile-HMMs of three universally conserved phylogenetic marker genes (Ribosomal_S12_S23 (PF00164), Ribosomal_L18e (PF00828), and Ribosomal_S9 (PF00380)) that were selected from a previous publication (Lang *et al.*, 2013).

Results

Thaumarchaeotal cobalamin synthesis

Genomic analysis of the ammonia-oxidizing archaeon *Nitrosopumilus maritimus* SCM1 and related thaumarchaeotal representatives indicates a genetic capacity for cobalamin biosynthesis. Using *N. maritimus* SCM1 as a reference, we identified a complete biosynthetic pathway and separate gene clusters encoding distinct stages of cobalamin biosynthesis (Figure 1, Supplementary Figures S1 and S2). Although both the aerobic and anaerobic pathways for cobalamin synthesis share a majority of their enzymes, several steps and associated enzymes are specific to each (Figure 1a). *N. maritimus* and related *Thaumarchaeota* (Figures 1b and c) lack genes encoding enzymes of the aerobic pathway (*CobG*, *CobF*, *CobNST*) and possess genes encoding enzymes specific to the anaerobic biosynthesis pathway (*CbiX*, *CbiG*, *CbiD*). This is supported by the presence and absence of marker genes as defined by the KEGG (Supplementary Figures S1 and S2). *N. maritimus* and other *Thaumarchaeota* were also found to encode cobalamin-dependent enzymes such as methionine synthase (Nmar_1267), ribonucleotide reductase (Nmar_1627), and methylmalonyl CoA mutase (B₁₂ binding domain, Nmar_0958), as well as probable cobalt transporters (Nmar_0878).

As further support for cobalamin biosynthetic potential, we identified dual *cob/cbi* gene clusters conserved across all known ammonia-oxidizing thaumarchaeotal genomes. The two relevant gene clusters are present in the genomes of *Candidatus* Nitrosoarchaeum limnia, *Candidatus* Nitrosopumilus salaria, *Candidatus* Nitrososphaera gargensis, and *Candidatus* Nitrosopumilus koreensis AR1 (Figures 1b and c), in addition to *Cenarchaeum symbiosum* and *Candidatus* Nitrosotenuis uzonensis (Supplementary Figure S2 and data not shown, respectively). With the possible exception of *Can. N. gargensis*, these gene clusters exhibit highly conserved synteny (Figures 1b and c, Supplementary Figure S2), indicative of selection for coordinated regulation. Furthermore, these gene clusters encode separate stages of cobalamin biosynthesis, with the first stage encoding six enzymes for the early pathway converting uroporphyrinogen III to

Figure 1 Identification of the cobalamin synthesis pathway and associated gene clusters in available thaumarchaeotal genomes. The pathway in (a) is adapted from the studies by Moore *et al.* (2013) and Raux *et al.* (1999). Horizontal arrows indicate homology between aerobic and anaerobic pathway enzymes. Grey enzyme names were not detected in thaumarchaeotal genomes. *CbiJ is also not present in other known archaeal cobalamin producers and therefore should not be considered an essential gene for this pathway in *Thaumarchaeota*. (b) A six-member cobalamin synthesis gene cluster with conserved synteny across several thaumarchaeotal genomes. The cluster encodes enzymes (highlighted by blue boxes in (a)) in the upper pathway, and includes three enzymes (bolded) specific to the anaerobic pathway. (c) A four-member gene cluster possessing enzymes (highlighted by green boxes in (a)) corresponding to the final steps of cobalamin synthesis. The apparent missing *cobD* gene in AR1 may be a genome annotation error given that it appears to be in the genome but may be truncated (data not shown).

cobyrinic acid a,c-diamide (Figures 1a and b) and the second stage encoding four enzymes for the later conversion of cobyrinic acid a,c-diamide to vitamin B₁₂ (Figures 1a and c).

Our results also show that many *Archaea* possess a similar complement of cobalamin biosynthetic genes, which cluster together and co-occur genomically (Supplementary Figures S2 and S3). This, along with the observed sequence and gene order

conservation, suggests widespread functional conservation of the cobalamin biosynthetic pathway among the *Archaea*.

Environmental distributions of cobalamin synthesis genes
We searched for cobalamin biosynthesis genes within 430 metagenomes derived from 28 studies, representing a diverse range of marine, freshwater

and hypersaline environments. Eleven cobalamin synthesis genes were selected based on their representation of both the aerobic and anaerobic pathway (Figure 1a), broad distribution throughout the cobalamin pathway and consistency with a core majority of genes used previously to identify cobalamin producers (Sañudo-Wilhelmy *et al.*, 2014).

Our screen identified 187 223 total cobalamin biosynthetic genes across all metagenomes, 95 911 of which matched the 11 marker genes (Figure 2a). Taxonomic assignments classified these 95 911 genes into three main phyla (Figure 2b): *Proteobacteria* (P), *Cyanobacteria* (C), and *Thaumarchaeota* (T). At the species level, *Thaumarchaeota* dominated among prevalent organisms (Table 1), comprising three of the four most frequent contributors of *cob/cbi* genes overall. *Thaumarchaeota* were also significantly overrepresented within the context of cobalamin synthesis (contributing 16.1% of *cob/cbi* genes) relative to their community abundance (contributing 2.5% of ribosomal marker genes). These results indicate that *Thaumarchaeota* are abundant global cobalamin producers through the anaerobic biosynthetic pathway, and may be a keystone species for cobalamin production.

In addition to evaluating all metagenome datasets together, we partitioned studies into their individual metagenome samples (430 in total), each of which was associated with distinct environmental parameters. Environment-specific metagenomes varied widely in taxonomic composition of cobalamin genes, within and between studies (Figure 2c). Hierarchical clustering of cobalamin taxonomic profiles of all metagenomes demonstrated visual ecotype partitioning (Figure 2d), including distinct T, P and C environments, or mixed environments (e.g., P + C or P + T). These recurring ecotypes show broad geographic distribution and non-uniform latitudinal gradients (Figure 2e).

Thaumarchaeota ranged from undetected to contributing as much as ~70–80% of the cobalamin genes in various samples from the Microbial Initiative in Low Oxygen areas off Concepcion and Oregon (MI_LOCO; <http://mi-loco.coas.oregonstate.edu>), Western Channel Observatory Microbial Metagenomic Study (http://camera.crbs.ucsd.edu/projects/details.php?id=CAM_PROJ_WesternChannelOMM; Gilbert *et al.*, 2010) and Antarctica metagenome (http://camera.crbs.ucsd.edu/projects/details.php?id=CAM_PROJ_AntarcticaAquatic) studies (Figure 2c, Supplementary Table S1). In addition, *Thaumarchaeota* were the dominant source of *cob/cbi* genes in several metagenomes included in this analysis (Supplementary Table S1), including the Arctic metagenome (66%, Figure 3b), deep-ocean bathypelagic habitat metagenome (50%), the station ALOHA (500 m depth) metagenome (67%) and individual sites from the Global Ocean Surveys dataset. Several metagenome studies included few samples (e.g., a single sample for the Arctic metagenome), and may therefore not be representative of their respective environments.

Thaumarchaeotal cobalamin genes correlate with environmental variables

The observed variation (Figure 2c) and clustering (Figures 2d and e) of distinct and recurring cobalamin ecotypes (Figures 2d and e) indicate that specific environmental variables drive the dynamics of cobalamin production. We performed correlation analyses across and within all metagenome datasets, and identified specific factors, including depth, nutrient levels, temperature and seasonality, that correlate with the taxonomic composition of cobalamin synthesis genes (Figure 3). The proportion of thaumarchaeotal cobalamin genes correlated most strongly ($r=0.55$) with sample depth (Figures 3a and b), consistent with observations of thaumarchaeotal dominance in deep water column metagenome studies (Figure 3 and Supplementary Figure S4). The correlation with depth was stronger across 63 samples from the MI_LOCO study ($r=0.73$, Figure 3d), which is likely attributed to higher within-study methodological consistency.

Further analysis of the MI_LOCO dataset revealed additional correlating variables that are consistent with known aspects of thaumarchaeotal ecology. Levels of thaumarchaeotal cobalamin synthesis genes exhibited strong positive correlations with numerous factors (Figure 3d), including nitrate ($r=0.75$), sample depth ($r=0.73$), phosphate ($r=0.61$) and dissolved organic nitrogen ($r=0.60$). The strongest negative correlations with thaumarchaeotal *cob/cbi* genes included bacterial cell number ($r=-0.53$) and ammonium concentration ($r=-0.40$). Low ammonium mixing ratios are consistent with a role for marine *Thaumarchaeota* as high affinity ammonia oxidizers (Martens-Habben *et al.*, 2009; Horak *et al.*, 2013; Nakagawa and Stahl, 2013).

Taxonomic cobalamin gene contributions also varied temporally. This was most apparent in a single metagenomic study from the Western English Channel time-series experiments sampled from station L4 (Figure 3b). The average proportion of thaumarchaeotal cobalamin synthesis genes was low (~3%) for the August and April time points, but increased up to 73% in January, suggesting that *Thaumarchaeota* are dominant contributors to cobalamin production in cold or winter-associated marine waters.

Additional phyla-vs-phyla correlations revealed a strong anticorrelation between thaumarchaeotal and cyanobacterial cobalamin synthesis genes (Figure 3a and see either-or effect in Figure 3c). This was evident in additional analyses that revealed no apparent T + C-rich ecotypes (Figure 2d). Similar patterns were observed for three phylogenetic marker genes (Supplementary Figure S5), suggesting a broader niche partitioning or exclusion between these two major marine clades. Unlike *Thaumarchaeota*, cyanobacterial cobalamin synthesis gene abundances correlated positively with temperature (Figure 3a), consistent with warm water

Figure 2 Survey of cobalamin synthesis genes across aquatic metagenomes. **(a)** The full set of *cob/cbi* genes and subset of eleven representative marker genes (black) used in the screen and their abundance (total number of detected genes). **(b)** Taxonomic breakdown of all detected *cob/cbi* genes across 430 aquatic metagenomes. **(c)** Distributions of thaumarchaeotal, cyanobacterial and proteobacterial cobalamin gene contributions (% of total *cob/cbi* genes) for 28 different studies. **(d)** Clustering and heatmap visualization of taxonomic composition of *cob/cbi* genes for the 430 metagenomes. The color scale reflects the proportion of *cob/cbi* genes in each sample affiliated with a particular taxonomic group. More information on each sample can be found in Supplementary Table S1. **(e)** Global distribution of cobalamin synthesis genes. To emphasize *Thaumarchaeota*-rich metagenomes, data points with higher thaumarchaeotal contributions have been slightly enlarged and overlaid preferentially in cases where there are overlapping data points. The *Thaumarchaeota*-rich Arctic metagenome described in the text is indicated by an asterisk.

Table 1 The 15 most abundant species contributing detected metagenomic *cob/cbi* genes across all datasets

Species	Phylum/class	<i>cob/cbi</i> genes
<i>Rhodobacterales</i> bacterium HTCC2255	Alphaproteobacteria	9091
<i>Candidatus Nitrosopumilus</i> sp. AR2	Thaumarchaeota	4394
<i>Candidatus Nitrosopumilus salaria</i>	Thaumarchaeota	4110
<i>Candidatus Nitrosoarchaeum limnia</i>	Thaumarchaeota	3656
SAR324 cluster bacterium SCGC AAA001 C10	Alphaproteobacteria	3283
<i>Nitrosopumilus maritimus</i> SCM1	Thaumarchaeota	2672
<i>Prochlorococcus marinus</i> str. MIT 9301	Cyanobacteria	2422
<i>Prochlorococcus marinus</i> str. AS9601	Cyanobacteria	2371
<i>Gamma proteobacterium</i> SCGC AAA007 O20	Gammaproteobacteria	2279
<i>Candidatus Nitrosopumilus koreensis</i> AR1	Thaumarchaeota	2039
<i>Chlorobium phaeovibrioides</i> DSM 265	Chlorobi	1942
<i>Candidatus Nitrosoarchaeum koreensis</i> MY1	Thaumarchaeota	1912
<i>Neptuniibacter caesariensis</i>	Gammaproteobacteria	1663
Endosymbiont of <i>Bathymodiolus</i> sp.	Gammaproteobacteria	1571
<i>Prochlorococcus marinus</i> subsp. <i>pastoris</i> str. CCMP1986	Cyanobacteria	1226

Taxonomic annotations were assigned based on the top BLAST match and are therefore dependent on the existing species diversity of the reference (NCBI) database. Thaumarchaeotal species are in bold.

cyanobacterial blooms, and negatively with depth (Figure 3a). We also identified a positive correlation between cobalamin gene proportions of *Thaumarchaeota* and *Nitrospinae*, a nitrite-oxidizing bacterial phylum. This likely reflects a functional association between these two groups, linked by nitrification.

Discussion

Available genomes and metagenomic studies provide strong support that *Thaumarchaeota* are globally important vitamin B₁₂ producers. Although cobalamin synthesis is distributed throughout both *Archaea* and *Bacteria*, only a restricted subset of microorganisms encodes this genetic capacity. Bacterial cobalamin synthesis is associated with select members of *Cyanobacteria*, *Alphaproteobacteria*, *Gammaproteobacteria*, and *Bacteroidetes* (Sañudo-Wilhelmy et al., 2014), which are consistent with the dominant phyla observed in the metagenomic datasets analyzed here (Figure 2b). Within the *Archaea*, only methanogenic *Euryarchaeota* and extremophilic crenarchaeal representatives were recognized as cobalamin producers prior to this study. For example, methanogens are known to produce and require cobalamin for acetoclastic and hydrogenotrophic methanogenesis (reviewed in DiMarco et al. (1990)). In addition, cobamides are produced and required by *Halobacterium* (Woodson et al., 2003) and known sulfur-metabolizing *Crenarchaeota* (Kräutler et al., 1988). Given that *Thaumarchaeota* are estimated to represent the most abundant *Archaea* on the planet, and are among the most abundant prokaryotes in the ocean (Karner et al., 2001), the capacity to synthesize vitamin cofactors within this phylum implies global biogeochemical, metabolic and ecological significance.

Genetic capacity for cobalamin synthesis is very tightly conserved. For example, several genes from the multi-step cobalamin synthesis pathway are

considered reliable indicators for the entire pathway. If a genome encodes *cbiA/cobB*, *cbiC/cobH* or *cobT*, then the entire cobalamin synthesis pathway is likely encoded and expressed in that microorganism (Bertrand et al., 2011). Indeed, all available thaumarchaeotal genomes encode this capacity, and demonstrate conserved clusters of cobalamin synthesis genes (Figure 1 and Supplementary Figures S2 and S3). Providing physiological support for cobalamin synthesis by the *Thaumarchaeota*, existing pure cultures of ammonia-oxidizing archaea (including, *N. maritimus* SCM1 and *Nitrososphaera viennensis*) are grown in medium prepared without vitamin supplementation (Martens-Habben and Stahl, 2011; Tourna et al., 2011). Given that the gene for the low efficiency B₁₂-independent methionine synthesis pathway (*metE*) is absent in the genome of *N. maritimus* SCM1, growth in the absence of vitamin supplementation provides strong supporting evidence for thaumarchaeotal cobalamin synthesis. Direct measurements of gene expression and cobalamin production by pure cultures of *N. maritimus* SCM1 and additional *Thaumarchaeota* would confirm thaumarchaeotal cobalamin synthesis and provide additional information on the structure, regulation and fate of this vitamin when produced by marine *Thaumarchaeota*. Although all known ammonia-oxidizing archaea possess the genetic capacity for cobalamin synthesis, the absence of homologous genes in *Calditerrivibrio subterranean* (Supplementary Figure S2), a deep branching archaeon related to the *Thaumarchaeota* (Nunoura et al., 2011), suggests that cobalamin synthesis is not universally distributed among related *Archaea*.

Cobalamin gene analysis of available aquatic metagenomic datasets provides compelling support for niche specificity and biogeochemical controls on cobalamin production. Taxonomic affiliations indicate that although proteobacterial cobalamin synthesis genes can co-occur with either *Cyanobacteria* or

Figure 3 Phylum and metadata correlations. (a) Correlation matrix for all species-metadata correlations with $r > 0.25$. Correlations were measured between numerical metadata from all 430 metagenome samples and the per-taxon proportion of cobalamin genes. Correlations between taxonomic proportions were also measured to identify potential species–species associations. (b) Left: proportion of cobalamin synthesis genes vs depth for three phyla (computed using all 430 metagenome samples). Middle: proportion of cobalamin synthesis genes vs different time points from the 2008–2009 L4 station time series. Right: per-gene taxonomic proportions of cobalamin genes for an Arctic metagenome. *Thaumarchaeota* (red), *Cyanobacteria* (green), *Proteobacteria* (blue), Unclassified (dark grey), *Bacteroidetes/Chlorobi* (purple), *Actinobacteria* (black), *Firmicutes* (orange), and Other (light gray). Bar widths are proportional to the square root of the sample sizes. (c) Phyla-vs-Phyla contributions of cobalamin synthesis genes across all 430 metagenome samples. All correlations exhibit a negative linear relationship, except for *Thaumarchaeota*-vs-*Cyanobacteria*, which exhibits an either-or relationship, suggestive of competitive or niche exclusion. (d) Correlation analysis for the MI_LOCO metagenome. The strongest correlating variables with thaumarchaeotal *cob/cbi* gene contribution are shown on the left, and plots for the top two positively (nitrate concentration and sample depth) and negatively correlated (ammonium and bacterial cell concentration) variables are shown on the right.

Thaumarchaeota, these two non-proteobacterial cobalamin producers display strong mutual exclusion (Figures 2 and 3). This trend was also observed when repeating the analysis using ribosomal phylogenetic markers (Supplementary Figure S5), suggesting broad niche partitioning between these major marine groups. In addition to light sensitivity of *Thaumarchaeota* (described below), a possibility is that high cyanobacterial photosynthetic production of molecular oxygen interferes with the anaerobic thaumarchaeotal cobalamin biosynthetic pathway, which possesses an oxygen-sensitive intermediate (cobalt-precorrin-5B; Moore *et al.*, 2013). In addition to niche partitioning of thaumarchaeotal and cyanobacterial cobalamin synthesis genes, metadata

correlations revealed a positive correlation with marine nitrite-oxidizing *Nitrospinae*, which is consistent with the ammonia-oxidizing role for most marine *Thaumarchaeota*. *Nitrospina*-like bacteria have also been implicated as the primary nitrite oxidizers in oxygen minimum upwelling zones (Levipan *et al.*, 2014), which is consistent with high thaumarchaeotal abundance in these marine environments (Supplementary Figure S4).

The metagenomic data are also consistent with observed geographical and temporal patterns of thaumarchaeotal abundances. In particular, the urea-fueled polar and halocline abundances of *Thaumarchaeota* (Alonso-Sáez *et al.*, 2012) were reflected here in polar cobalamin synthesis gene

abundances in the same datasets. Winter peak abundances of *Thaumarchaeota* (Wuchter *et al.*, 2006; Herfort *et al.*, 2007; Galand *et al.*, 2010; Pitcher *et al.*, 2011) are consistent with our observation of a winter thaumarchaeotal cobalamin gene maximum at station L4. In addition, previous research has reported high thaumarchaeotal abundance in oxygen minimum zones (Pitcher *et al.*, 2011; Peng *et al.*, 2013), which reflects our observation of abundant thaumarchaeotal cobalamin genes in the MI_LOCO metagenomes. The longstanding observation of abundant *Thaumarchaeota* (then ‘mesophilic *Crenarchaeota*’) in deep marine water columns (Karner *et al.*, 2001; Sintès *et al.*, 2013) is also supported strongly by the metagenomic data survey conducted in this study. Importantly, polar environments, aphotic water columns, oxygen minimum zones, and winter time points share low light conditions, which would be a likely explanation for the observed anticorrelation between thaumarchaeotal and cyanobacterial cobalamin synthesis and ribosomal genes. The biogeographical cobalamin synthesis gene distributions observed in this study are consistent with observations of light-dependent decreases in marine thaumarchaeotal abundance below sea ice (Alonso-Sáez *et al.*, 2012) and known photoinhibition of thaumarchaeotal ammonia oxidation (Merbt *et al.*, 2012). Other factors proposed to explain aquatic thaumarchaeotal distributions include nutrient availability, temperature, dissolved oxygen and organic substrates (Peng *et al.*, 2013).

Observations of vitamin B₁₂ maxima in ocean waters below the marine photic zone are consistent with an important role for *Thaumarchaeota* in marine cobalamin production, given their high proportional abundance with depth. Previous observations of increased cobalamin concentrations in ocean mesopelagic zones (Sañudo-Wilhelmy *et al.*, 2012) have led to suggestions that vitamin-producing microbial plankton below the photic zone are important contributors to marine biogeochemistry, in addition to essential suppliers of growth factors for phytoplankton and microbial communities in the photic zone (Sañudo-Wilhelmy *et al.*, 2014). Although ocean upwelling zones might be considered a mechanism for supplying cobalamin to surface waters, vitamin B₁₂ concentrations did not correlate with upwelling index off the coast of Baja California, Mexico (Panzeca *et al.*, 2009). With respect to high latitude environments, cobalt supplementation of North Atlantic marine samples led to increased vitamin B₁₂ production (Panzeca *et al.*, 2008). Our metagenomic analysis of high latitude samples implies that *Thaumarchaeota* were responsible for vitamin B₁₂ production in this experiment, and implies that thaumarchaeotal productivity may be limited by cobalt concentrations. Importantly, vitamin B₁₂ was shown to influence phytoplankton growth and community composition in samples taken from the subarctic Gulf of Alaska (Koch *et al.*, 2011), which is consistent with controls

on phytoplankton growth by vitamin B₁₂ in freshwater environments as well (Daisley, 1969; Cavari and Grossowicz, 1977). Together, these results provide early evidence that cobalamin production by *Thaumarchaeota* may directly influence phytoplanktonic communities in polar surface waters, freshwater and possibly in marine surface waters via mixing with ocean water column vitamin pools.

Conclusion

Thaumarchaeotal vitamin production in most of the global ocean would have enormous potential impacts on microbial communities and food web dynamics. Here we provide evidence that *Thaumarchaeota* are among the most abundant global cobalamin vitamin producers. All known ammonia-oxidizing members of the *Thaumarchaeota* possess the genetic capacity to produce cobalamin, and data from multiple metagenomic studies reveal niche partitioning of this keystone community function. Thaumarchaeotal cobalamin synthesis genes demonstrate spatial and temporal biogeographical distributions. Combining pure culture experimental work and field studies will help assess the proportion of marine cobalamin pools derived from archaeal synthesis, identify the role of archaeal cobalamin in marine biogeochemistry, and characterize links between global thaumarchaeotal cobalamin production and microbial community composition and succession.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

We thank Mirko Vucicevich for technical assistance. This research was supported by the National Science and Engineering Research Council of Canada (NSERC).

References

- Alonso-Sáez L, Waller AS, Mende DR, Bakker K, Farnelid H, Yager PL *et al.* (2012). Role for urea in nitrification by polar marine Archaea. *Proc Natl Acad Sci USA* **109**: 17989–17994.
- Bateman A, Coin L, Durbin R, Finn RD, Hollich V, Griffiths-Jones S *et al.* (2004). The Pfam protein families database. *Nucleic Acids Res* **32**: D138–D141.
- Bertrand EM, Saito MA, Rose JM, Riesselman CR, Lohan MC, Noble AE *et al.* (2007). Vitamin B₁₂ and iron colimitation of phytoplankton growth in the Ross Sea. *Limnol Oceanogr* **52**: 1079–1093.
- Bertrand EM, Saito MA, Jeon YJ, Neilan BA. (2011). Vitamin B₁₂ biosynthesis gene diversity in the Ross Sea: the identification of a new group of putative

- polar B₁₂ biosynthesizers. *Environ Microbiol* **13**: 1285–1298.
- Blainey PC, Mosier AC, Potanina A, Francis CA, Quake SR. (2011). Genome of a low-salinity ammonia-oxidizing archaeon determined by single-cell and metagenomic analysis. *PLoS One* **6**: e16626.
- Cavari B, Grossowicz N. (1977). Seasonal distribution of vitamin B₁₂ in Lake Kinneret. *Appl Environ Microbiol* **34**: 120–124.
- Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG. (2005). Algae acquire vitamin B₁₂ through a symbiotic relationship with bacteria. *Nature* **438**: 90–93.
- Daisley K. (1969). Monthly survey of vitamin B₁₂ concentrations in some waters of the English Lake District. *Limnol Oceanogr* **14**: 224–228.
- DiMarco AA, Bobik TA, Wolfe RS. (1990). Unusual coenzymes of methanogenesis. *Ann Rev Biochem* **59**: 355–394.
- Francis CA, Roberts KJ, Beman JM, Santoro AE, Oakley BB. (2005). Ubiquity and diversity of ammonia-oxidizing archaea in water columns and sediments of the ocean. *Proc Natl Acad Sci USA* **102**: 14683–14688.
- Galand PE, Gutiérrez-Provecho C, Massana R, Gasol JM, Casamayor EO. (2010). Inter-annual recurrence of archaeal assemblages in the coastal NW Mediterranean Sea (Blanes Bay Microbial Observatory). *Limnol Oceanogr* **55**: 2117–2125.
- Gilbert JA, Meyer F, Schriml L, Joint I, Mühling M, Field D. (2010). Metagenomes and metatranscriptomes from the L4 long-term coastal monitoring station in the Western English Channel. *Stand Genomic Sci* **3**: 183–193.
- Giovannoni SJ. (2012). Vitamins in the sea. *Proc Natl Acad Sci USA* **109**: 13888–13889.
- Gobler CJ, Norman C, Panzeca C, Taylor GT, Sañudo-Wilhelmy SA. (2007). Effect of B-vitamins (B₁, B₁₂) and inorganic nutrients on algal bloom dynamics in a coastal ecosystem. *Aquat Microb Ecol* **49**: 181–194.
- Grant MA, Kazamia E, Cicuta P, Smith AG. (2014). Direct exchange of vitamin B₁₂ is demonstrated by modelling the growth dynamics of algal–bacterial cocultures. *ISME J* **8**: 1418–1427.
- Haft DH, Selengut JD, White O. (2003). The TIGRFAMs database of protein families. *Nucleic Acids Res* **31**: 371–373.
- Hallam SJ, Mincer TJ, Schleper C, Preston CM, Roberts K, Richardson PM *et al.* (2006). Pathways of carbon assimilation and ammonia oxidation suggested by environmental genomic analyses of marine Crenarchaeota. *PLoS Biol* **4**: e95.
- Herfort L, Schouten S, Abbas B, Veldhuis MJ, Coolen MJ, Wuchter C *et al.* (2007). Variations in spatial and temporal distribution of Archaea in the North Sea in relation to environmental variables. *FEMS Microbiol Ecol* **62**: 242–257.
- Horak RE, Qin W, Schauer AJ, Armbrust EV, Ingalls AE, Moffett JW *et al.* (2013). Ammonia oxidation kinetics and temperature sensitivity of a natural marine community dominated by Archaea. *ISME J* **2013**: 2023–2033.
- Iverson V, Morris RM, Frazar CD, Berthiaume CT, Morales RL, Armbrust EV. (2012). Untangling genomes from metagenomes: revealing an uncultured class of marine Euryarchaeota. *Science* **335**: 587–590.
- Kanehisa M, Araki M, Goto S, Hattori M, Hirakawa M, Itoh M *et al.* (2008). KEGG for linking genomes to life and the environment. *Nucleic Acids Res* **36**: D480–D484.
- Karner MB, DeLong EF, Karl DM. (2001). Archaeal dominance in the mesopelagic zone of the Pacific Ocean. *Nature* **409**: 507–510.
- Kim BK, Jung MY, Yu DS, Park SJ, Oh TK, Rhee SK *et al.* (2011). Genome sequence of an ammonia-oxidizing soil archaeon, "*Candidatus Nitrosoarchaeum koreensis*" MY1. *J Bacteriol* **193**: 5539.
- Koch F, Alejandra Marcoval M, Panzeca C, Bruland KW, Sañudo-Wilhelmy SA, Gobler CJ. (2011). The effect of vitamin B₁₂ on phytoplankton growth and community structure in the Gulf of Alaska. *Limnol Oceanogr* **56**: 1023–1034.
- Kräutler B, Kohler HPE, Stupperich E. (1988). 5'-Methylbenzimidazolyl-cobamides are the corrinoids from some sulfate-reducing and sulfur-metabolizing bacteria. *Eur J Biochem* **176**: 461–469.
- Lang JM, Darling AE, Eisen JA. (2013). Phylogeny of bacterial and archaeal genomes using conserved genes: supertrees and supermatrices. *PLoS One* **8**: e62510.
- Lebedeva EV, Hatzenpichler R, Pelletier E, Schuster N, Hauzmayer S, Bulaev A *et al.* (2013). Enrichment and genome sequence of the group I.1a ammonia-oxidizing archaeon "*Ca. Nitrosotenuis uzonensis*" representing a clade globally distributed in thermal habitats. *PLoS One* **8**: e80835.
- Leininger S, Urich T, Schlöter M, Schwark L, Qi J, Nicol G *et al.* (2006). Archaea predominate among ammonia-oxidizing prokaryotes in soils. *Nature* **442**: 806–809.
- Levipan HA, Molina V, Fernandez C. (2014). *Nitrospina*-like bacteria are the main drivers of nitrite oxidation in the seasonal upwelling area of the Eastern South Pacific (Central Chile ~36°S). *Environ Microbiol Rep*; doi:10.1111/1758-2229.12158.
- Lloyd KG, Schreiber L, Petersen DG, Kjeldsen KU, Lever MA, Steen AD *et al.* (2013). Predominant archaea in marine sediments degrade detrital proteins. *Nature* **496**: 215–218.
- Luo H, Tolar BB, Swan BK, Zhang CL, Stepanauskas R, Moran MA *et al.* (2014). Single-cell genomics shedding light on marine Thaumarchaeota diversification. *ISME J* **8**: 732–736.
- Markowitz VM, Chen I-MA, Palaniappan K, Chu K, Szeto E, Pillay M *et al.* (2014). IMG 4 version of the integrated microbial genomes comparative analysis system. *Nucleic Acids Res* **42**: D560–D567.
- Martens-Habbena W, Berube PM, Urakawa H, de la Torre JR, Stahl DA. (2009). Ammonia oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. *Nature* **461**: 976–979.
- Martens-Habbena W, Stahl DA. (2011). Nitrogen metabolism and kinetics of ammonia-oxidizing archaea. *Methods Enzymol* **496**: 465–487.
- Merbt SN, Stahl DA, Casamayor EO, Martí E, Nicol GW, Prosser JL. (2012). Differential photoinhibition of bacterial and archaeal ammonia oxidation. *FEMS Microbiol Lett* **327**: 41–46.
- Moore SJ, Lawrence AD, Biedendieck R, Deery E, Frank S, Howard MJ *et al.* (2013). Elucidation of the anaerobic pathway for the corrin component of cobalamin (vitamin B₁₂). *Proc Natl Acad Sci USA* **110**: 14906–14911.

- Morris JJ, Lenski RE, Zinser ER. (2012). The Black Queen Hypothesis: evolution of dependencies through adaptive gene loss. *mBio* **3**: e00036–00012.
- Nakagawa T, Stahl DA. (2013). Transcriptional response of the archaeal ammonia oxidizer *Nitrosopumilus maritimus* to low and environmentally relevant ammonia concentrations. *Appl Environ Microbiol* **79**: 6911–6916.
- Nunoura T, Takaki Y, Kakuta J, Nishi S, Sugahara J, Kazama H *et al.* (2011). Insights into the evolution of Archaea and eukaryotic protein modifier systems revealed by the genome of a novel archaeal group. *Nucleic Acids Res* **39**: 3204–3223.
- Ondov BD, Bergman NH, Phillippy AM. (2011). Interactive metagenomic visualization in a Web browser. *BMC Bioinformatics* **12**: 385.
- Panzeca C, Beck AJ, Leblanc K, Taylor GT, Hutchins DA, Sañudo-Wilhelmy SA. (2008). Potential cobalt limitation of vitamin B₁₂ synthesis in the North Atlantic Ocean. *Glob Biogeochem Cycles* **22**: GB2029.
- Panzeca C, Beck AJ, Tovar-Sanchez A, Segovia-Zavala J, Taylor GT, Gobler CJ *et al.* (2009). Distributions of dissolved vitamin B₁₂ and Co in coastal and open-ocean environments. *Estuar Coast Shelf Sci* **85**: 223–230.
- Peng X, Jayakumar A, Ward BB. (2013). Community composition of ammonia-oxidizing archaea from surface and anoxic depths of oceanic oxygen minimum zones. *Front Microbiol* **4**: 177.
- Pitcher A, Villanueva L, Hopmans EC, Schouten S, Reichart G-J, Damsté JSS. (2011). Niche segregation of ammonia-oxidizing archaea and anammox bacteria in the Arabian Sea oxygen minimum zone. *ISME J* **5**: 1896–1904.
- Prosser JI, Nicol GW. (2008). Relative contributions of archaea and bacteria to aerobic ammonia oxidation in the environment. *Environ Microbiol* **10**: 2931–2941.
- Raux E, Schubert HL, Roper JM, Wilson KS, Warren MJ. (1999). Vitamin B₁₂: insights into biosynthesis's mount improbable. *Bioorg Chem* **27**: 100–118.
- Rho M, Tang H, Ye Y. (2010). FragGeneScan: predicting genes in short and error-prone reads. *Nucleic Acids Res* **38**: e191–e191.
- Rodionov DA, Vitreschak AG, Mironov AA, Gelfand MS. (2003). Comparative genomics of the vitamin B₁₂ metabolism and regulation in prokaryotes. *J Biol Chem* **278**: 41148–41159.
- Sañudo-Wilhelmy SA, Cutter LS, Durazo R, Smail EA, Gómez-Consarnau L, Webb EA *et al.* (2012). Multiple B-vitamin depletion in large areas of the coastal ocean. *Proc Natl Acad Sci USA* **109**: 14041–14045.
- Sañudo-Wilhelmy SA, Gómez-Consarnau L, Suffridge C, Webb EA. (2014). The role of B vitamins in marine biogeochemistry. *Ann Rev Mar Sci* **6**: 339–367.
- Sañudo-Wilhelmy S, Gobler C, Okbamichael M, Taylor G. (2006). Regulation of phytoplankton dynamics by vitamin B₁₂. *Geophys Res Lett* **33**: L04604.
- Sintes E, Bergauer K, De Corte D, Yokokawa T, Herndl GJ. (2013). Archaeal *amoA* gene diversity points to distinct biogeography of ammonia-oxidizing Crenarchaeota in the ocean. *Environ Microbiol* **15**: 1647–1658.
- Spang A, Hatzepichler R, Brochier-Armanet C, Rattei T, Tischler P, Spieck E *et al.* (2010). Distinct gene set in two different lineages of ammonia-oxidizing archaea supports the phylum Thaumarchaeota. *Trends Microbiol* **18**: 331–340.
- Stahl DA, de la Torre JR. (2012). Physiology and diversity of ammonia-oxidizing archaea. *Ann Rev Microbiol* **66**: 83–101.
- Stubbe J. (1994). Binding site revealed of nature's most beautiful cofactor. *Science* **266**: 1663–1664.
- Tourna M, Stieglmeiera M, Spanga A, Könneke M, Schintlmeister A, Urich T *et al.* (2011). *Nitrososphaera viennensis*, an ammonia oxidizing archaeon from soil. *Proc Natl Acad Sci USA* **108**: 8420–8425.
- Tripp HJ, Kitner JB, Schwalbach MS, Dacey JW, Wilhelm LJ, Giovannoni SJ. (2008). SAR11 marine bacteria require exogenous reduced sulphur for growth. *Nature* **452**: 741–744.
- Walker CB, de la Torre JR, Klotz MG, Urakawa H, Pinel N, Arp DJ *et al.* (2010). *Nitrosopumilus maritimus* genome reveals unique mechanisms for nitrification and autotrophy in globally distributed marine crenarchaea. *Proc Natl Acad Sci USA* **107**: 8818–8823.
- Woodson JD, Peck RF, Krebs MP, Escalante-Semerena JC. (2003). The *cobY* gene of the archaeon *Halobacterium* sp. strain NRC-1 is required for de novo cobamide synthesis. *J Bacteriol* **185**: 311–316.
- Wuchter C, Abbas B, Coolen MJ, Herfort L, van Bleijswijk J, Timmers P *et al.* (2006). Archaeal nitrification in the ocean. *Proc Natl Acad Sci USA* **103**: 12317–12322.

This work is licensed under a Creative Commons Attribution 3.0 Unported License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/3.0/>

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)