

ORIGINAL ARTICLE

The ecology of pelagic freshwater methylotrophs assessed by a high-resolution monitoring and isolation campaign

Michaela M Salcher^{1,2}, Stefan M Neuenschwander¹, Thomas Posch¹ and Jakob Pernthaler¹

¹Limnological Station, Institute of Plant Biology, University of Zurich, Kilchberg, Switzerland and ²Institute of Hydrobiology, Biology Centre CAS, Ceske Budejovice, Czech Republic

Methylotrophic planktonic bacteria fulfill a particular role in the carbon cycle of lakes via the turnover of single-carbon compounds. We studied two planktonic freshwater lineages (LD28 and PRD01a001B) affiliated with *Methylophilaceae* (Betaproteobacteria) in Lake Zurich, Switzerland, by a combination of molecular and cultivation-based approaches. Their spatio-temporal distribution was monitored at high resolution ($n=992$ samples) for 4 consecutive years. LD28 methylotrophs constituted up to 11×10^7 cells l^{-1} with pronounced peaks in spring and autumn–winter, concomitant with blooms of primary producers. They were rare in the warm water layers during summer but abundant in the cold hypolimnion, hinting at psychrophilic growth. Members of the PRD01a001B lineage were generally less abundant but also had maxima in spring. More than 120 axenic strains from these so far uncultivated lineages were isolated from the pelagic zone by dilution to extinction. Phylogenetic analysis separated isolates into two distinct genotypes. Isolates grew slowly ($\mu_{\max} = 0.4 d^{-1}$), were of conspicuously small size, and were indeed psychrophilic, with higher growth yield at low temperatures. Growth was enhanced upon addition of methanol and methylamine to sterile lake water. Genomic analyses of two strains confirmed a methylotrophic lifestyle with a reduced set of genes involved in C1 metabolism. The very small and streamlined genomes (1.36 and 1.75 Mb) shared several pathways with the marine OM43 lineage. As the closest described taxa (*Methylotenera* sp.) are only distantly related to either set of isolates, we propose a new genus with two species, that is, ‘*Candidatus Methylopumilus planktonicus*’ (LD28) and ‘*Candidatus Methylopumilus turicensis*’ (PRD01a001B).

The ISME Journal (2015) 9, 2442–2453; doi:10.1038/ismej.2015.55; published online 5 May 2015

Introduction

Pelagic freshwater bacteria are key factors in the biogeochemical processes of lakes. One particular role in the carbon cycle is fulfilled by methylotrophic bacteria that use single-carbon (C1) compounds such as methanol as sole sources of carbon and energy. Methanol is a main degradation product of pectin and lignin and is therefore mainly associated with plant growth and decay (Heikes *et al.*, 2002; Vorholt, 2012). In aquatic systems, methanol and other C1 compounds (for example, methylated sulfur species and methylated amines) may also be linked to primary producers. C1 units can be either directly released by phytoplankton species or can result of bacterial breakdown of algal carbohydrates (Sieburth and Keller, 1989; Milne *et al.*, 1995; Heikes *et al.*, 2002). Other potential C1 sources include atmospheric

deposition, allochthonous import (Heikes *et al.*, 2002), or photochemical degradation of dissolved organic carbon (Dixon *et al.*, 2011a; Mopper *et al.*, 1991). Methanol is chemically stable and highly soluble in water because of its polarity (Dixon *et al.*, 2011b; Heikes *et al.*, 2002). So far, reports of *in situ* concentrations of methanol have been restricted to marine habitats with concentrations ranging in the nm scale (70–429 nmol l^{-1}) and turnover rates as low as 1 day (range: 1–83 days) in very oligotrophic sites (Dixon *et al.*, 2011a, b). An indication for high methylotrophic activity and an important role in the *in situ* biogeochemical cycling of C1 compounds was provided by the overrepresentation of methanol dehydrogenases (MDHs) affiliated with *Methylophilaceae* (Betaproteobacteria) in metatranscriptomic and metaproteomic surveys in freshwater and marine habitats (Sowell *et al.*, 2011; Gifford *et al.*, 2013; Georges *et al.*, 2014; Hanson *et al.*, 2014).

Aerobic methylotrophs can be found in various families of Proteobacteria, Verrucomicrobia, NC10, and Gram-positive microbes (Chistoserdova 2011). The family *Methylophilaceae* includes four formally described genera (*Methylophilus*, *Methylovorus*, *Methylobacillus* and *Methylotenera*) isolated from

Correspondence: MM Salcher. Current address: Biology Centre CAS, Institute of Hydrobiology, Na Sádkách 7, 370 05, České Budějovice, Czech Republic.

E-mail: michaelasalcher@gmail.com

Received 22 December 2014; revised 19 February 2015; accepted 6 March 2015; published online 5 May 2015

terrestrial environments or freshwater sediments (Yordy and Weaver, 1977; Jenkins *et al.*, 1987; Govorukhina and Trotsenko, 1991; Kalyuzhnaya *et al.*, 2006). Several representatives of these genera have been studied in detail and genome sequenced (Lapidus *et al.*, 2011; Beck *et al.*, 2014). Moreover, members of the planktonic marine lineage OM43 have been isolated via dilution to extinction methods and genome sequenced (Connon and Giovannoni, 2002; Giovannoni *et al.*, 2008; Song *et al.*, 2009; Huggett *et al.*, 2012). OM43 have very small genomes (1.3 Mb) and a reduced set of genes encoding methylotrophic pathways (Giovannoni *et al.*, 2008; Chistoserdova 2011; Halsey *et al.*, 2011; Huggett *et al.*, 2012). Members of this lineage are abundant in coastal oceans, especially during phytoplankton blooms (Sekar *et al.*, 2004; Morris *et al.*, 2006). Less is known about planktonic freshwater *Methylophilaceae* (Newton *et al.*, 2011). Two ubiquitous species-like lineages (LD28 and PRD01a001B) are widely distributed and especially LD28 are highly abundant and seasonally persistent in the pelagic zone of lakes (Newton *et al.*, 2011; Parveen *et al.*, 2011; Salcher *et al.*, 2011a; Eiler *et al.*, 2012).

Here, we present an in-depth ecological assessment of members of the freshwater LD28 and PRD01a001B lineages in prealpine Lake Zurich, Switzerland. We used a multiphasic approach combining (i) high-resolution monitoring of their distribution, (ii) isolation of 127 pure cultures via dilution to extinction, (iii) genome sequencing of two representatives and (iv) laboratory experiments about their methylotrophic lifestyle and adaptation to cold water temperatures. Based on these analyses, we propose the establishment of a new genus with two species, that is, '*Candidatus* *Methylopumilus* planktonicus' and '*Candidatus* *Methylopumilus* turicensis'.

Materials and methods

Study site and sampling

Lake Zurich is a large, deep (136 m), prealpine, oligomesotrophic lake, characterized by persistent annual blooms of the toxic cyanobacterium *Planktothrix rubescens* (Posch *et al.*, 2012). Water samples were taken every second week from 17 January 2008 to 14 December 2011 at the deepest part of the lake. Vertical profiles of temperature, conductivity, turbidity, oxygen and chlorophyll *a* content were recorded using a YSI multiprobe (Yellow Springs Instruments, model 6600, Yellow Springs, OH, USA) and a bbe fluoroprobe (TS-16-12, bbe Moldaenke GmbH, Schwentinental, Germany). The fluoroprobe was calibrated to distinguish different phytoplankton groups and *P. rubescens* according to the fluorescent spectra of their pigments (Beutler *et al.*, 2002). Water samples were taken from 0, 5, 10, 20, 30, 40, 60, 80, 100 m depth, and additionally from the depth of recorded chlorophyll *a* maxima. Forty milliliters of water were fixed with formaldehyde (2% final

concentration) for flow cytometric counts of bacteria, and 5–10 ml were fixed with paraformaldehyde (pH 7.4, 2% final concentration) for CARD-FISH (fluorescence *in situ* hybridization followed by catalyzed reporter deposition) analyses. Unfixed water samples for the isolation of bacteria and the preparation of media were transported to the laboratory within 15 min. Chemical parameters were determined in monthly intervals with standard techniques by the Zurich Water Supply Company (Zurich, Switzerland).

Abundances of microbes affiliated with LD28 and PRD01a001B

Total microbial abundances were determined by an inFlux V-GS cell sorter (Becton Dickinson, Franklin Lakes, NJ, USA) equipped with an ultraviolet (355 nm) laser. All samples were stained with 4',6-diamidino-2-phenylindole (DAPI, 1 µg ml⁻¹ final concentration), and weighted before and after measurements for sample volume quantification. Scatter plots of DAPI fluorescence vs 90° light scatter were analyzed with an in-house software (J Villiger, unpublished).

CARD-FISH with fluorescein-labeled tyramides was carried out as previously described (Sekar *et al.*, 2003) with probes specifically detecting microbes affiliated with LD28 (probe LD28-1017; Salcher *et al.*, 2011a) and a newly designed probe targeting the PRD01a001B lineage. Probe PRD-732 (5'-TCAGTA TTAGTCCAGGGGGCTG-3') and its competitor (5'-TC AGTATTAGSCCAGGGGGCTG-3') was designed in ARB and tested following the workflow described in Salcher *et al.* (2011b). Highest stringency was obtained with 50% formamide in the hybridization buffer. CARD-FISH-stained samples were analyzed by fully automated high-throughput microscopy (Zeder and Pernthaler, 2009). Images were analyzed with the freely available image analysis software ACMEtool (technobiology.ch), and interfering autofluorescent cyanobacteria or debris particle were individually excluded from hybridized cells. At least 10 high-quality images and >1000 DAPI-stained bacteria were analyzed per sample. A lower limit for the quantification for FISH-stained cells (0.14% of all DAPI-stained particles) was applied as determined previously (Neuenschwander *et al.*, 2015). Cell size estimations of LD28 bacteria were done on 12 samples (*n* = 1225 cells) with the software LUCIA (Laboratory Imaging Prague, Prague, Czech Republic) following the workflow described in Posch *et al.* (2009). Samples corresponded to different LD28 population dynamics, that is, at times of population increase in spring (5 m depth, 09–24 April 2008), population decline in spring (5 m depth, 31 March–11 May 2009), population increase in autumn (5 m depth, 18 October–14 December 2011) and in the hypolimnion (80 m depth, 11 May 2009).

Isolation of planktonic bacteria affiliated with *Methylophilaceae*

A total of 46 isolation campaigns were carried out during 2010–2013. Unfixed water samples from 5 m

depth were used for isolation of planktonic bacteria. Medium was mostly prepared from filtered and autoclaved lake water (0.2 µm pore sized polysulfone filters, Millipore, Billerica, MA, USA), amended with different types of carbon and nutrient sources (for details, see Supplementary Table S1). Two milliliter aliquots of medium were dispersed to 24-well plates (Thermo Fisher Scientific, Waltham, MA, USA) and inoculated with prefiltered (0.45 µm pore sized polycarbonate filters, Millipore) and diluted water samples. Total bacterial cell counts of inocula were determined by microscopy and samples were diluted to final inoculum sizes of approximately 1–10 cells well⁻¹ (Supplementary Table S1). Twenty-four-well plates were incubated at close to *in situ* temperatures in a dark–light cycle (16:8 h) for 2–3 months. Growth of bacteria in individual wells was examined by pipetting 50 µl aliquots onto diagnostic glass slides (Thermo Fisher Scientific), which were air-dried at 45 °C, embedded into DAPI mix and inspected with a fluorescence microscope (Zeiss AxioImagerM.1, Carl Zeiss, Jena, Germany). Wells containing dense bacterial cultures were further split in 500 µl aliquots that were (i) propagated to fresh medium, (ii) amended with glycerol (30% final concentration) and stored at –80 °C, and (iii) pelleted by centrifugation (30 min, 16 000 × *g*) and used for DNA isolation, PCR and subsequent sequencing of the 16S ribosomal RNA genes. PCR was conducted with general bacterial primers GM3f and GM4r (Muyzer *et al.*, 1995) and 1 µl template of either extracted DNA (GenElute Bacterial Genomic DNA Kit, Sigma-Aldrich, St Louis, MO, USA) or pelleted bacterial biomass subjected to 2–3 cycles of freeze thawing. PCR products were purified and sequenced with the ABI BigDye chemistry on an ABI 3130x Genetic Analyzer (Applied Biosystems, Foster City, CA, USA) with primers GM3f, GM4r and GM1f (Muyzer *et al.*, 1995). Partial sequences were assembled with the software DNA baser v3.5.0 (Heracle BioSoft, Lilienthal, Germany), aligned with the SINA web aligner (www.arb.silva.de) and imported into ARB (Ludwig *et al.*, 2004) using the SILVA database SSU Ref 111 (Pruesse *et al.*, 2007). A bootstrapped maximum likelihood tree (GTR-GAMMA model) of sequences of isolates affiliated with *Methylophilaceae*, their closest relatives and other described genera of *Methylophilaceae* was constructed on a dedicated web server (Stamatakis *et al.*, 2005).

Growth characteristics and cell size of strain MMS-2-53

Strain MMS-2-53 was grown in triplicated setups of 40 ml sterile lake water amended with a mix of vitamins and amino acids (Supplementary Table S1) and different concentrations of methanol (0, 10 µM, 100 µM and 1 mM methanol) in Tissue Culture Flasks (25 cm², TPP, Trasadingen, Switzerland) at 22 °C in the dark for 69 days. One milliliter subsamples were taken twice a week, fixed with formaldehyde and counted by flow cytometry (see above). Sizing of

cells was conducted with a Zeiss AxioImagerM.1 microscope after DAPI staining and filtration (0.2 µm filters, Osmonics, Penang, Malaysia) with the software LUCIA (Posch *et al.*, 2009). At least 100 individual cells per sample were subjected to image analysis (total *n* = 4384).

Growth on methanol and methylamine and temperature preferences of strain MMS-2-53 were tested in triplicated setups of 40 ml sterile lake water amended with amino acids, vitamins and 1 mM methanol (variant A), or 1 mM methanol and 100 µM methylamine (variant B) in Tissue Culture Flasks (TPP) incubated in the dark at 6 °C, 13 °C, 18 °C and 22 °C for 6 weeks.

Genome sequencing and annotation of strains MMS-2-53 and MMS-10A-171

Whole-genome sequencing of strains MMS-2-53 and MMS-10A-171 was done after an initial multiple displacement amplification as cell yields and DNA concentrations were too low for direct sequencing. Unfixed cells of strain MMS-2-53 were sorted on a flow cytometer, lysed and amplified with either the Repli-G Single Cell Kit (Qiagen, Hilden, Germany; three parallel samples with 20 000–40 000 sorted cells) following the manufacturer's instructions or with a RepliPHI (epicenter, 14 samples with 5000 or 10 000 sorted cells) according to Dean *et al.* (2002) and Woyke *et al.* (2011). DNA extracts of strain MMS-10A-171 were amplified with the RepliPHI (11 samples). Multiple displacement amplification products were pooled for three and one individual libraries for strains MMS-2-53 and MMS-10A-171, respectively. 550-bp libraries were produced according to the manufacturer's instructions (TruSeq DNA PCR-Free LT Sample Preparation Kit, Illumina, San Diego, CA, USA) and 2 × 150-bp paired end sequences were generated on an Illumina MiSeq instrument (MCS versions 2.3.0.3 and 2.4.1.3, Illumina) with a 300-cycle MiSeq Reagent v2 kit (Illumina). Draft assemblies with A5-miseq (Coil *et al.*, 2014) resulted in 6 and 38 contigs of strain MMS-2-53 and MMS-10A-171, respectively, whereof the largest 5 and 8 contigs (>5000 bp) were used for primer design. A total of 53 and 71 primers targeting contig ends and regions with a coverage of <10× were designed with the software Geneious 7.1.7 (www.geneious.com) and used for additional PCRs. Totals of 83 and 121 additional sanger sequences finally closed all gaps and poorly covered regions in the two genomes.

Gene calling and draft annotations were done with the Rapid Annotation using Subsystem Technology (RAST) server version 2.0 (Aziz *et al.*, 2008) and the MicroScope web platform (Vallenet *et al.*, 2013). Inconsistencies between the two methods and overlapping CDS were manually curated in ARTEMIS (Rutherford *et al.*, 2000) and short CDS (<150 bp) without BLAST hits were discarded as artefacts. Homologies between genes of the two strains and

their closest relatives, that is, methylophilales bacterium HTCC2181 (Giovannoni *et al.*, 2008), *Methylothena mobilis* JLW8, *Methylothena versatilis* 301, *Methylovorus glucosetrophus* SIP3-4 (Lapidus *et al.*, 2011), *Methylophilus methylotrophus* ATCC 53528 (NZ_ARJW000000000), *Methylobacillus flagellatus* KT (Chistoserdova *et al.*, 2007) and a typical freshwater representative (*Polynucleobacter necessarius* subsp. *asymbioticus* QLW-P1DMWA-1; Meincke *et al.*, 2012) were identified with the MicroScope web platform using protein identity cutoffs of $\geq 30\%$ and $\geq 70\%$ (minLrap ≥ 0.8 ; maxLrap ≥ 0). Reconstructions of methylotrophy pathways were modeled according to their closest relatives using comparative genomics tools of the MicroScope platform (Vallet *et al.*, 2013). Digital DNA–DNA hybridization of the two genomes and closely related strains was done via the Genome-to-Genome Distance Calculator using the GGDC 2.0 BLAST+ alignment method (Meier-Kolthoff *et al.*, 2013). Annotated genome sequences have been deposited at EMBL (sequence accessions LN827929 and LN794158).

Statistical analyses

Before statistical analyses, data of relative abundances (% of DAPI) were arcsine transformed and cell numbers and physicochemical variables were $\log(x+1)$ transformed to better approximate a normal distribution. Redundancy analysis was used for determining the effects of environmental data on microbes affiliated with LD28 and PRD01a001B in Lake Zurich. The significance of added variables was tested by a Monte Carlo permutation test (500 permutations), and only variables with a highly significant correlation ($P < 0.0001$) were included in the analysis. Significant differences in the growth of isolate MMS-2-53 according to different treatments were tested by one-way analysis of variances (Tukey *post-hoc* tests). All analyses were performed with the Microsoft EXCEL add-in program XLSTAT (www.xlstat.com).




Figure 1 Collapsed maximum likelihood tree of the 16S rDNA of *Methylophilaceae*. Lineages LD28 ('*Ca. Methylophilum planktonicus*') and PRD01a001B ('*Ca. Methylophilum turicensis*') are shown in gray. The scale bar at the bottom applies to 1% sequence divergence.

Results

Spatio-temporal distribution of *Methylophilaceae* in Lake Zurich in relation to environmental parameters

Microbes affiliated with LD28 (see Figure 1 for phylogenetic positioning) were present in all analyzed samples ($n=992$) in varying numbers. Two seasonal maxima (up to 11.3×10^7 cells l^{-1} or 4.1% of DAPI) in spring and autumn–winter were discernable in all four analyzed years (Figure 2a, Supplementary Figure S1). Bacteria of the PRD01a001B lineage were below the limit of detection ($<0.14\%$ of DAPI-stained cells) in most of our samples ($n=642$). However, a distinct maximum of up to 4% of DAPI-stained cells (3.2×10^7 cells l^{-1}) was found in the hypolimnion in summer 2011 (Figure 2b, Supplementary Figure S1). This abundance peak lasted for several weeks between April and August 2011, but was absent in the other years, where maxima of 1% or 2.4×10^7 cells l^{-1} were restricted to the epilimnion in spring.

Redundancy analysis resulted in a significant explanation of variability in both lineages (Figure 3). Axis 1 correlated mostly with chlorophyll *a* concentrations (total chlorophyll *a* and chlorophyll *a* associated with the cyanobacterium *Planktothrix rubescens* and diatoms), nitrate, phosphate and ammonium, whereas the second axis separated chlorophyll *a* of *P. rubescens* from that of diatoms, as well as nitrate from phosphate and ammonium (see Supplementary Figure S2 for details of all environmental parameters). Population sizes of both lineages were clearly separated from each other; LD28 methylotrophs were more related to total and *P. rubescens* chlorophyll *a* (Pearson's correlation coefficient $R=0.729$ and $R=0.611$, respectively, $P < 0.0001$), whereas PRD01a001B bacteria were more related to ammonium and diatom chlorophyll *a* ($R=0.353$ and $R=0.309$, respectively, $P < 0.0001$). Nitrate and phosphate was negatively related to both methylotrophs ($R=-0.618$ and $R=-0.233$ for nitrate and $R=-0.655$ and $R=-0.160$ for phosphate, correlation to LD28 and PRD01a001B, respectively, $P < 0.0001$).

Isolation of *Methylophilaceae* from Lake Zurich

2077 of a total of 6507 inspected wells contained dense cultures, 788 of these appeared to be monocultures, and 115 and 12 of these were pure cultures affiliated with LD28 and PRD01a001B, respectively (Figure 1, Supplementary Figures S3, S4, Table S1). All isolates within each of the two lineages were very closely related to each other according to the 16S ribosomal RNA genes (98.6–100% and 99.7–100% sequence similarity for LD28 and PRD01a001B, respectively), but they were distantly related to the strains of the other genotype (95.2–96.5%) and to so far described taxa of *Methylophilaceae* ($<95.6\%$; Table 1). Both clades and the marine OM43 formed a monophyletic sister lineage to *Methylothena*, with

PRD01a001B branching more deeply from LD28 and OM43 (Figure 1, Supplementary Figures S3 and S4). One isolate of LD28 (strain MMS-2-53) and PRD01a001B (strain MMS-10A-171) were subjected to genome sequencing (Table 2). Digital whole-genome DNA–DNA hybridization of our isolates and genome sequenced *Methylophilaceae* resulted in very low similarities (<26.2%; Table 1). Thus, these isolates, together with the marine OM43, likely represent a new genus within *Methylophilaceae*. We therefore propose to establish a new ‘*Candidatus*’ genus with two new species, that is, ‘*Candidatus* *Methylopumilus planktonicus*’ for isolates affiliated with LD28 (strain MMS-2-53) and ‘*Candidatus* *Methylopumilus turicensis*’ for isolates affiliated with PRD01a001B (strain MMS-10A-171).

Growth of strain MMS-2-53 with different concentrations of methanol

Cultivation of strain MMS-2-53 was only successful in sterile lake water, whereas no growth was visible on agar plates (neither prepared with lake water nor with

conventional R2A, NSY or K-medium agar plates amended with methanol) or in conventional liquid media (R2, NSY, K-medium, inorganic basal medium




Figure 3 Redundancy analysis of environmental parameters explaining the variability in cell numbers of microbes affiliated with LD28 and PRD01a001B in Lake Zurich. LD28, abundances of LD28; PRD, abundances of PRD01a001B; total chla, total chlorophyll *a*; *P.rub. chla*, chlorophyll *a* associated with *Planktothrix rubescens*; diat. chla, chlorophyll *a* associated with diatoms; O₂, oxygen concentrations; NH₄, ammonium concentrations; NO₃, nitrate concentrations; PO₄, phosphate concentrations.




Figure 2 Seasonality of abundances (10⁷ cells l⁻¹) of microbes affiliated with LD28 (‘*Ca. Methylopumilus planktonicus*’. (a) and PRD01a001B (‘*Ca. Methylopumilus turicensis*’. (b) in Lake Zurich, Switzerland, during the investigation period 2008–2011. Sampling dates and depths are displayed as white circles.

Table 1 16S rDNA similarities, digital DNA–DNA hybridization, and proportions of shared protein-coding genes ($\geq 70\%$ protein similarity) of the two strains and different members of *Methylophilaceae*

	16S rDNA		DNA–DNA hyb.		Shared genes (%)	
	MMS-2-53	MMS-10A-171	MMS-2-53	MMS-10A-171	MMS-2-53	MMS-10A-171
‘ <i>Ca. M. planktonicus</i> ’ MMS-2-53		96.5		22.5		12.8
‘ <i>Ca. M. turicensis</i> ’ MMS-10A-171	96.5		22.5		12.4	
HTCC2181 (marine OM43)	95.3	96.3	17.1	23.4	10.9	8.0
<i>Methylothermobacter mobilis</i> JLW8	95.2	95.4	24.5	21.7	10.0	20.0
<i>Methylothermobacter versatilis</i> 301	94.7	95.6	24.4	20.1	8.6	17.9
<i>Methylophilus methylotrophus</i> ATCC 53528	93.7	93.8	21.3	19.9	8.4	14.2
<i>Methylobacillus flagellatus</i> KT	94.2	94.2	26.2	18.5	8.0	17.3
<i>Methylovorus glucosetrophus</i> SIP3-4	93.9	94.4	25.1	20.2	8.4	19.8

Abbreviation: rDNA, ribosomal DNA.

Table 2 Genome statistics of strains MMS-2-53 and MMS-10A-171

	‘ <i>Ca. M. planktonicus</i> ’ MMS-2-53	‘ <i>Ca. M. turicensis</i> ’ MMS-10A-171
Genome size (bp)	1 356 428	1 754 988
Sequencing coverage	481.6x	191.6x
Organization	1 chromosome	1 chromosome
GC content (%)	36.97	44.52
Total genes	1434	1806
RNA genes	45	49
rRNA operons	1	2
Protein-coding genes	1389	1757
Mean-coding sequence length (bp)	917.2	923.9
Mean intergenic length (bp)	69.8	83.2
Protein-coding density (%)	93.56	92.29
EMBL sequence accession number	LN827929	LN794158

Abbreviation: rRNA, ribosomal RNA.

amended with methanol). Strain MMS-2-53 reached maximal cell densities of 1.8×10^9 cells l^{-1} in sterile lake water after 36 days, corresponding to a maximal growth rate of $\mu_{\max} = 0.16 \text{ d}^{-1}$ (Figure 4a). Significantly higher growth yields (4.9×10^9 cells l^{-1}) and growth rates ($\mu_{\max} = 0.37 \text{ d}^{-1}$) were achieved by the addition of methanol at different concentrations. Substrate saturation was apparently reached at 100 μM methanol, which yielded identical growth rates as the addition of 1 mM methanol. A precise calculation of K_m or K_s was hampered by the unknown concentration of methanol in the lake water. The cell sizes of strain MMS-2-53 were very small (median = $0.041 \mu\text{m}^3$, $n = 3552$) during stationary growth phase (Figure 4b), however, volumes significantly increased during exponential phase ($0.075 \mu\text{m}^3$, $n = 832$). No significant differences in cell volume were recorded at different concentrations of methanol. Interestingly, cell volumes similar to those of stationary phase cultures were also obtained from CARD-FISH-stained preparations of LD28 microbes in Lake Zurich during population decline in the epilimnion in spring ($0.048 \mu\text{m}^3$, $n = 416$) and in the hypolimnion ($0.055 \mu\text{m}^3$, $n = 83$), whereas cells were even significantly smaller in autumn samples ($0.03 \mu\text{m}^3$, $n = 327$). LD28 growing to population maxima in the epilimnion in spring were larger ($0.061 \mu\text{m}^3$, $n = 399$) and in between the values obtained from cultures (Figure 4c).


Effects of water temperature and additional C1 substrates

The relative abundances of LD28 microbes in Lake Zurich showed recurrent maxima in winter and were negatively related to water temperature in the upper water layers (epilimnion, 0–10 m depth, Figures 5a and b). To test for direct effects of water temperature and additional C1 substrates, we incubated strain MMS-2-53 with 1 mM methanol at different temperatures and in the presence or absence of 100 μM of methylamine (Figure 5c). There was only a slight effect of incubation temperatures on abundances when cells were grown with 1 mM methanol only ($4.5\text{--}6.7 \times 10^9$ cells l^{-1}). The addition of methylamine increased the cell yield by 1.2–2-fold and led to significantly higher cell numbers at 6 °C (12.8×10^9 cells l^{-1}) when methylamine served as second C1 substrate.

Insight into the genomes of planktonic freshwater *Methylophilaceae*

The sizes of the two newly sequenced complete genomes of strains MMS-2-53 and MMS-10A-171 were 1.36 and 1.75 Mb with a GC content of 37% and 44%, respectively (Table 2, Supplementary Figures S5 and S6). Only very low proportions of the 1389 and 1757 protein-coding genes were shared with other *Methylophilaceae* ($< 20\%$) at a protein similarity cutoff of $\geq 70\%$ (Figure 6a, Table 1).

The ISME Journal



Discussion

Our high-resolution sampling campaign in Lake Zurich ($n=992$ individual samples, Figure 2,

Supplementary Figures S1 and S2) helped to identify potential environmental drivers for planktonic freshwater *Methylophilaceae*. The striking annually recurrent maxima of LD28 microbes in spring and autumn–winter seem to be mainly determined by the availability of C1 substrates released by phytoplankton. Seasonal maxima of these *Methylophilaceae* in late spring and autumn–winter of up to 3.5% of pyrosequencing reads were also reported from Lake Erken, Sweden (Eiler *et al.*, 2012). However, the latter study found negative correlations to chlorophyll *a*, whereas we observed the opposite, which may be a consequence of the higher resolution of our data set (Figures 2 and 3). The same authors also reported an association of LD28 microbes with cyanobacterial blooms in several Swedish lakes (Eiler and Bertilsson, 2004), which is in accordance to our results from a single system (Figures 2 and 3). A positive correlation of LD28 to the chlorophyll *a* of the dominant primary producer of Lake Zurich, the

cyanobacterium *Planktothrix rubescens*, has been reported before (Salcher *et al.*, 2011a). Moreover, an operational taxonomic unit affiliated with LD28 was exclusively present in the depth layer of maximal *P. rubescens* densities in this lake (Van den Wyngaert *et al.*, 2011). *Methylophilaceae* were also highly abundant in mesocosms containing the cyanobacterial toxin microcystin (Mou *et al.*, 2013). Marine *Methylophilaceae* were enriched in seawater chemostats amended with diatom and cyanobacterial dissolved organic matter only after six generation times, but not after three, pointing at slow growth and a potential cross-feeding of C1 compounds that were released by other microbes (Landa *et al.*, 2014). Slow growth is also in accordance with our experimental data from isolate MMS-2-53 (Figure 4). Interestingly, strain HTCC2181 of the marine OM43 lineage grows much faster than strain MMS-2-53 (Giovannoni *et al.*, 2008; Halsey *et al.*, 2011), although it shares several metabolic and genomic features with LD28 (for example, comparable cell yields, synergistic effects of a second C1 substrate, genome size, GC content and methylotrophic pathways).

Nutrient acquisition by LD28 and PRD01a001B

Both genotypes showed a negative correlation to nitrate and phosphate concentrations (Figure 3, Supplementary Figure S2), which is partly in agreement with earlier studies (Salcher *et al.*, 2011a; Eiler *et al.*, 2012). High-affinity phosphate transporters (Pst) for scavenging phosphorus at low concentrations were encoded in both genomes. Therefore, these microbes seem to be well adapted to phosphorus limitation. Some freshwater *Methylophilaceae* showed chemotaxis toward ammonium and phosphate (Dennis *et al.*, 2013), although the taxonomic resolution in that study was not sufficient to discriminate between different lineages. We found a positive correlation to ammonium (Figure 3), as well as several genes encoding ammonium transporters in both genotypes. Thus, it seems likely that nitrogen demand is covered by ammonium uptake.

It is unclear what triggered the high abundances of microbes affiliated with PRD01a001B in summer 2011 (Figure 2, Supplementary Figure S1), but our observation nevertheless illustrates the importance of long-term time series for understanding the ecology and occurrence patterns of these slowly growing methylotrophs in lakes. Generally, the PRD01a001B lineage of *Methylophilaceae* is less explored than LD28, however, they were detected in a wide range of freshwater and oligosaline lakes (Wu *et al.*, 2006).

Psychrophilic methylotrophs?

Negative correlations of planktonic freshwater *Methylophilaceae* and water temperature have been reported previously (Salcher *et al.*, 2008, 2011a). However, so far it was not clear if these patterns were

triggered by water temperature *per se* or by other co-varying factors such as higher nutrient availability in deeper water layers and during mixis events. Here, we experimentally demonstrate a relationship between incubation temperature and cell yield of strain MMS-2-53 (Figure 5). Although growth was also observed at higher temperatures (22 °C), bacteria reached significantly lower densities compared with 6 °C. Thus, we propose that these microbes are psychrophilic, that is, adapted to low temperatures. Interestingly, this psychrophilic aspect of growth seems to be enhanced by the availability of an additional C1 source, methylamine. Most of our strains were isolated during autumn–spring (65%), while in summer, when water temperatures reached >20 °C, we were less successful (Supplementary Table S1). However, further studies are needed to test for strain-specific temperature optima, as even closely related isolates can profoundly differ in their thermal adaptation (Hahn *et al.*, 2014; Yung *et al.*, 2014).

Genomic similarities and differences within *Methylophilaceae*

The very small genome size and low GC content of strain MMS-2-53 (Table 2, Supplementary Figure S5) is comparable to the streamlined genomes of marine methylotrophs HTCC2181 and HIMB624 of the OM43 clade (Giovannoni *et al.*, 2008; Huggett *et al.*, 2012). Strain MMS-10A-171 has a slightly larger genome, a higher GC content and shares more protein-coding genes with *Methylotenera*, *Methylobacillus* and *Methylovorus* (Table 2, Figure 6, Supplementary Figure S6). Both strains and members of the marine OM43 lineage encode similar types of rhodopsins, light driven proton pumps that are tuned to green light and thus, may be beneficial in productive conditions in lakes (Giovannoni *et al.*, 2008). Moreover, our isolates were obligate methylotrophs, as already suggested earlier because LD28 bacteria did not incorporate amino acids or sugars in microautoradiography assays (Salcher *et al.*, 2008, 2013). Both isolates possess genes encoding a special variant of MDH (*xoxF*) that most likely require the rare earth metal lanthanum as cofactor (Hibi *et al.*, 2011; Beck *et al.*, 2014). Methylotrophy is a very modular metabolic system; formaldehyde in particular can be oxidized via several different pathways, either by a formaldehyde dehydrogenase, via the H₄F pathway or the H₄MPT pathway (Chistoserdova 2011). Both isolates encode genes for the H₄F pathway. Strain MMS-10A-171 also encodes genes for the H₄MPT pathway, whereas it is absent in MMS-2-53 and in OM43 microbes (Giovannoni *et al.*, 2008; Huggett *et al.*, 2012; Figure 6b). Isolate MMS-2-53 grew to higher densities in the presence of methylamine as second C1 substrate, although we could not detect genes encoding methylamine oxidation (Figures 5 and 6). However, alternative routes for C1 substrate entry to the oxidative H₄F

pathway have been proposed for the marine OM43 (Halsey *et al.*, 2011). One gene (*metF*) that may facilitate methylamine oxidation together with a so far unknown methyltransferase is present in both analyzed genomes (Figure 6, Supplementary Table S2); thus, these microbes may be able to oxidize also other C1 compounds than methanol. Methanol is mainly used by microbes for obtaining energy (up to 97% in oligotrophic regions of oceans) but can also contribute up to 50% to bacterial carbon demand (Dixon *et al.*, 2011a, 2013). In the presence of other C1 compounds, methanol may thus be directly channeled to the ribulose monophosphate cycle for biomass production, whereas other C1 compounds may fuel energy demands. However, gene expression analyses are needed to verify this hypothesis and identify genes for methylamine oxidation.

Putative description of a new genus of Methylophilaceae

The low degree of sequence similarities on the level of 16S ribosomal RNA genes (<95.6%), digital whole-genome DNA–DNA hybridization (<26.2%) and shared protein-coding genes (<20%, Table 1) suggests that the here described isolates as well as the marine OM43 clade are not affiliated with any described genus of *Methylophilaceae*. Therefore, we propose a new provisional genus with two new species, that is, ‘*Candidatus* Methylopumilus planktonicus’ for LD28 with MMS-2-53 as type strain and ‘*Candidatus* Methylopumilus turicensis’ for PRD01a001B with MMS-10A-171 as type strain. Currently, additional analyses required for a formal species description, such as detailed metabolic analyses and cellular phospholipid fatty acids profiling, are hampered by the high amounts of biomass required for such assays. Moreover, we still face difficulties in composing a completely synthetic growth medium; so far, both isolates only grow well in sterile lake water.

Conclusion

We provide first insight in the ecology and unique lifestyle of an important group of so far uncultivated ubiquitous freshwater bacteria that occupy a distinct niche in the pelagic zone of lakes. This was achieved by combining high-resolution environmental monitoring, isolation of pure cultures, metabolic tests and genome sequencing of isolates. The two investigated strains remarkably differed in genome size (1.36 vs 1.76 Mb) and methylotrophic pathways (Figure 6, Table 2, Supplementary Table S2). It is conceivable that a common ancestor of our freshwater strains and the marine OM43 lineage already underwent gene streamlining (that is, losing several genes encoding methylotrophic pathways), but also gained new genes via horizontal gene transfer (encoding, for example, rhodopsins). As genome reduction is more

pronounced in strain MMS-2-53 than in MMS-10A-171, we believe that these microbes are ideal model organisms for future studies regarding gene loss and genome streamlining.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

We thank captain E Loher for help during sampling and the Zurich Water Supply (WVZ) for the provision of limnochemical data. Illumina library preparation and sequencing was done at the Genetic Diversity Centre (GDC), ETH Zurich. This study was supported by the Swiss National Science Foundation (SNF, Project 31EE30-132771) as part of the European Science Foundation EUROCORES Programme EuroEEFG.

References

- Aziz R, Bartels D, Best A, DeJongh M, Disz T, Edwards R *et al.* (2008). The RAST Server: rapid annotations using subsystems technology. *BMC Genomics* **9**: 75.
- Beck DAC, McTaggart TL, Setboonsarng U, Vorobev A, Kalyuzhnaya MG, Ivanova N *et al.* (2014). The expanded diversity of *Methylophilaceae* from Lake Washington through cultivation and genomic sequencing of novel ecotypes. *PLoS One* **9**: e102458.
- Beutler M, Wiltshire K, Meyer B, Moldaenke C, Lüring C, Meyerhöfer M *et al.* (2002). A fluorometric method for the differentiation of algal populations *in vivo* and *in situ*. *Photosynthesis Res* **72**: 39–53.
- Chistoserdova L, Lapidus A, Han C, Goodwin L, Saunders L, Brettin T *et al.* (2007). Genome of *Methylobacillus flagellatus*, molecular basis for obligate methylotrophy, and polyphyletic origin of methylotrophy. *J Bacteriol* **189**: 4020–4027.
- Chistoserdova L. (2011). Modularity of methylotrophy, revisited. *Environ Microbiol* **13**: 2603–2622.
- Coil D, Jospin G, Darling AE. (2014). A5-miseq: an updated pipeline to assemble microbial genomes from illumina miseq data. *arXiv preprint* **1401.5130**.
- Connon SA, Giovannoni SJ. (2002). High-throughput methods for culturing microorganisms in very-low-nutrient media yield diverse new marine isolates. *Appl Environ Microbiol* **68**: 3878–3885.
- Dean FB, Hosono S, Fang LH, Wu XH, Faruqi AF, Bray-Ward P *et al.* (2002). Comprehensive human genome amplification using multiple displacement amplification. *Proc Natl Acad Sci USA* **99**: 5261–5266.
- Dennis PG, Seymour J, Kumbun K, Tyson GW. (2013). Diverse populations of lake water bacteria exhibit chemotaxis towards inorganic nutrients. *ISME J* **7**: 1661–1664.
- Dixon JL, Beale R, Nightingale PD. (2011a). Rapid biological oxidation of methanol in the tropical Atlantic: significance as a microbial carbon source. *Biogeosci Discuss* **8**: 3899–3921.
- Dixon JL, Beale R, Nightingale PD. (2011b). Microbial methanol uptake in northeast Atlantic waters. *ISME J* **5**: 704–716.
- Dixon JL, Sargeant S, Nightingale PD, Colin Murrell J. (2013). Gradients in microbial methanol uptake:

- productive coastal upwelling waters to oligotrophic gyres in the Atlantic Ocean. *ISME J* 7: 568–580.
- Eiler A, Bertilsson S. (2004). Comparison of freshwater bacterial communities associated with cyanobacterial blooms in four swedish lakes. *Environ Microbiol* 6: 1228–1243.
- Eiler A, Heinrich F, Bertilsson S. (2012). Coherent dynamics and association networks among lake bacterioplankton taxa. *ISME J* 6: 330–342.
- Georges AA, El-Swais H, Craig SE, Li WKW, Walsh DA. (2014). Metaproteomic analysis of a winter to spring succession in coastal northwest Atlantic Ocean microbial plankton. *ISME J* 8: 1301–1313.
- Gifford SM, Sharma S, Booth M, Moran MA. (2013). Expression patterns reveal niche diversification in a marine microbial assemblage. *ISME J* 7: 281–298.
- Giovannoni SJ, Hayakawa DH, Tripp HJ, Stingl U, Givan SA, Cho JC et al. (2008). The small genome of an abundant coastal ocean methylotroph. *Environ Microbiol* 10: 1771–1782.
- Govorukhina NI, Trotsenko YA. (1991). Methylovorus, a new genus of restricted facultatively methylotrophic bacteria. *Int J Syst Bacteriol* 41: 158–162.
- Hahn MW, Koll U, Jezberová J, Camacho A. (2014). Global phylogeography of pelagic Polynucleobacter bacteria: restricted geographic distribution of subgroups, isolation by distance, and influence of climate. *Environ Microbiol* 17: 829–840.
- Halsey KH, Carter AE, Giovannoni SJ. (2011). Synergistic metabolism of a broad range of C₁ compounds in the marine methylotrophic bacterium HTCC2181. *Environ Microbiol* 14: 630–640.
- Hanson B, Hewson I, Madsen E. (2014). Metaproteomic survey of six aquatic habitats: discovering the identities of microbial populations active in biogeochemical cycling. *Microb Ecol* 67: 520–539.
- Heikes BG, Chang W, Pilson MEQ, Swift E, Singh HB, Guenther A et al. (2002). Atmospheric methanol budget and ocean implication. *Global Biogeochem Cycles* 16: 1133.
- Hibi Y, Asai K, Arafuka H, Hamajima M, Iwama T, Kawai K. (2011). Molecular structure of La3+-induced methanol dehydrogenase-like protein in Methylobacterium radiotolerans. *J Biosci Bioengineer* 111: 547–549.
- Huggett M, Hayakawa D, Rappe M. (2012). Genome sequence of strain HIMB624, a cultured representative from the OM43 clade of marine Betaproteobacteria. *Stand Genomic Sci* 6: 11–20.
- Jenkins O, Byrom D, Jones D. (1987). *Methylophilus* - a new genus of methanol-utilizing bacteria. *Int J Syst Bacteriol* 37: 446–448.
- Kalyuzhnaya MG, Bowerman S, Lara JC, Lidstrom ME, Chistoserdova L. (2006). *Methylotenera mobilis* gen. nov., sp. nov., an obligately methylamine-utilizing bacterium within the family Methylophilaceae. *Int J Syst Evol Microbiol* 56: 2819–2823.
- Landa M, Cottrell MT, Kirchman DL, Kaiser K, Medeiros PM, Tremblay L et al. (2014). Phylogenetic and structural response of heterotrophic bacteria to dissolved organic matter of different chemical composition in a continuous culture study. *Environ Microbiol* 16: 1668–1681.
- Lapidus A, Clum A, LaButti K, Kaluzhnaya MG, Lim S, Beck DAC et al. (2011). Genomes of three methylotrophs from a single niche reveal the genetic and metabolic divergence of the Methylophilaceae. *J Bacteriol* 193: 3757–3764.
- Ludwig W, Strunk O, Westram R, Richter L, Meier H, Yadhukumar et al. (2004). ARB: a software environment for sequence data. *Nucl Acid Res* 32: 1363–1371.
- Meier-Kolthoff J, Auch A, Klenk H-P, Goker M. (2013). Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinform* 14: 60.
- Meincke L, Copeland A, Lapidus A, Lucas S, Berry K, Glavina Del Rio T et al. (2012). Complete genome sequence of Polynucleobacter necessarius subsp. asymbioticus type strain (QLW-P1DMWA-1T). *Stand Genomics* 6: 74–83.
- Milne PJ, Riemer DD, Zika RG, Brand LE. (1995). Measurement of vertical distribution of isoprene in surface seawater, its chemical fate, and its emission from several phytoplankton monocultures. *Marine Chem* 48: 237–244.
- Mopper K, Zhou X, Kieber RJ, Kieber DJ, Sikorski RJ, Jones RD. (1991). Photochemical degradation of dissolved organic carbon and its impact on the oceanic carbon cycle. *Nature* 353: 60–62.
- Morris RM, Longnecker K, Giovannoni SJ. (2006). Pirellula and OM43 are among the dominant lineages identified in an Oregon coast diatom bloom. *Environ Microbiol* 8: 1361–1370.
- Mou X, Lu X, Jacob J, Sun S, Heath R. (2013). Metagenomic identification of bacterioplankton taxa and pathways involved in microcystin degradation in Lake Erie. *PLoS One* 8: e61890.
- Muyzer G, Teske A, Wirsén C, Jannasch H. (1995). Phylogenetic relationship of *Thiomicrospira* species and their identification in deep-sea hydrothermal vent samples by denaturing gel electrophoresis of 16S rDNA fragments. *Arch Microbiol* 164: 165–172.
- Neuenschwander SM, Pernthaler J, Posch T, Salcher MM. (2015). Seasonal growth potential of rare lake water bacteria suggest their disproportional contribution to carbon fluxes. *Environ Microbiol* 17: 781–795.
- Newton RJ, Jones SE, Eiler A, McMahon KD, Bertilsson S. (2011). A guide to the natural history of freshwater lake bacteria. *Microbiol Mol Biol R* 75: 14–49.
- Parveen B, Reveilliez J-P, Mary I, Ravet V, Bronner G, Mangot J-F et al. (2011). Diversity and dynamics of free-living and particle-associated Betaproteobacteria and Actinobacteria in relation to phytoplankton and zooplankton communities. *FEMS Microbiol Ecol* 77: 461–476.
- Posch T, Franzoi J, Prader M, Salcher MM. (2009). New image analysis tool to study biomass and morphotypes of three major bacterioplankton groups in an alpine lake. *Aquat Microb Ecol* 54: 113–126.
- Posch T, Koster O, Salcher MM, Pernthaler J. (2012). Harmful filamentous cyanobacteria favoured by reduced water turnover with lake warming. *Nat Climate Change* 2: 809–813.
- Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J et al. (2007). SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucl Acid Res* 35: 7188–7196.
- Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream M-A et al. (2000). Artemis: sequence visualization and annotation. *Bioinformatics* 16: 944–945.
- Salcher MM, Pernthaler J, Zeder M, Psenner R, Posch T. (2008). Spatio-temporal niche separation of planktonic Betaproteobacteria in an oligo-mesotrophic lake. *Environ Microbiol* 10: 2074–2086.

- Salcher MM, Pernthaler J, Frater N, Posch T. (2011a). Vertical and longitudinal distribution patterns of different bacterioplankton populations in a canyon-shaped, deep prealpine lake. *Limnol Oceanogr* **56**: 2027–2039.
- Salcher MM, Pernthaler J, Posch T. (2011b). Seasonal bloom dynamics and ecophysiology of the freshwater sister clade of SAR11 bacteria 'that rule the waves' (LD12). *ISME J* **5**: 1242–1252.
- Salcher MM, Posch T, Pernthaler J. (2013). *In situ* substrate preferences of abundant bacterioplankton populations in a prealpine freshwater lake. *ISME J* **7**: 896–907.
- Sekar R, Pernthaler A, Pernthaler J, Warnecke F, Posch T, Amann R. (2003). An improved protocol for quantification of freshwater *Actinobacteria* by fluorescence *in situ* hybridization. *Appl Environ Microbiol* **69**: 2928–2935.
- Sekar R, Fuchs BM, Amann R, Pernthaler J. (2004). Flow sorting of marine bacterioplankton after fluorescence *in situ* hybridization. *Appl Environ Microbiol* **70**: 6210–6219.
- Sieburth JM, Keller MD. (1989). Methylaminotrophic bacteria in xenic nanoalgal cultures: incidence, significance, and role of methylated algal osmoprotectants. *Biol Oceanograph* **6**: 383–395.
- Song J, Oh H-M, Jang-Cheon C. (2009). Improved culturability of SAR11 strains in dilution-to-extinction culturing from the East Sea, West Pacific Ocean. *FEMS Microbiol Lett* **295**: 141–147.
- Sowell SM, Abraham PE, Shah M, Verberkmoes NC, Smith DP, Barofsky DF et al. (2011). Environmental proteomics of microbial plankton in a highly productive coastal upwelling system. *ISME J* **5**: 856–865.
- Stamatakis A, Ludwig T, Meier H. (2005). RAXML-II: a program for sequential, parallel and distributed inference of large phylogenetic. *Concurr Comput-Pract Exp* **17**: 1705–1723.
- Vallenet D, Belda E, Calteau A, Cruveiller S, Engelen S, Lajus A et al. (2013). MicroScope - an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. *Nucl Acid Res* **41**: D636–D647.
- Van den Wyngaert S, Salcher MM, Pernthaler J, Zeder M, Posch T. (2011). Quantitative dominance of seasonally persistent filamentous cyanobacteria (*Planktothrix rubescens*) in the microbial assemblages of a temperate lake. *Limnol Oceanogr* **56**: 97–109.
- Vorholt JA. (2012). Microbial life in the phyllosphere. *Nat Rev Microbiol* **10**: 828–840.
- Woyke T, Sczyrba A, Lee J, Rinke C, Tighe D, Clingenpeel S et al. (2011). Decontamination of MDA reagents for single cell whole genome amplification. *PLoS One* **6**: e26161.
- Wu Q, Zwart G, Schauer M, Kamst-van AMP, Hahn M. (2006). Bacterioplankton community composition along a salinity gradient of sixteen high-mountain lakes located on the Tibetan plateau, China. *Appl Environ Microbiol* **72**: 5478–5485.
- Yordy JR, Weaver TL. (1977). Methylobacillus: a new genus of obligately methylotrophic bacteria. *Int J Syst Bacteriol* **27**: 247–255.
- Yung C-M, Vereen MK, Herbert A, Davis KM, Yang J, Kantorowska A et al. (2014). Thermally adaptive tradeoffs in closely-related marine bacterial strains. *Environ Microbiol*; doi:10.1111/1462-2920.12714.
- Zeder M, Pernthaler J. (2009). Multispot live-image autofocus for high-throughput microscopy of fluorescently stained bacteria. *Cytometry Part A* **75A**: 781–788.

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)