ORIGINAL ARTICLE

The methanogenic redox cofactor F_{420} is widely synthesized by aerobic soil bacteria

Blair Ney^{1,2,5}, F Hafna Ahmed^{1,5}, Carlo R Carere³, Ambarish Biswas⁴, Andrew C Warden², Sergio E Morales⁴, Gunjan Pandey², Stephen J Watt¹, John G Oakeshott², Matthew C Taylor², Matthew B Stott³, Colin J Jackson¹ and Chris Greening^{2,6}

¹Research School of Chemistry, Australian National University, Acton, Australian Capital Territory, Australia;

²The Commonwealth Scientific and Industrial Research Organisation, Land and Water, Acton, Australian Capital Territory, Australia; ³GNS Science, Wairakei Research Centre, Taupō, New Zealand and ⁴Department of Microbiology and Immunology, University of Otago, Dunedin, New Zealand

F_{420} is a low-potential redox cofactor that mediates the transformations of a wide range of complex organic compounds. Considered one of the rarest cofactors in biology, F_{420} is best known for its role in methanogenesis and has only been chemically identified in two phyla to date, the Euryarchaeota and Actinobacteria. In this work, we show that this cofactor is more widely distributed than previously reported. We detected the genes encoding all five known F_{420} biosynthesis enzymes (*cofC*, *cofD*, *cofE*, *cofG* and *cofH*) in at least 653 bacterial and 173 archaeal species, including members of the dominant soil phyla Proteobacteria, Chloroflexi and Firmicutes. Metagenome datamining validated that these genes were disproportionately abundant in aerated soils compared with other ecosystems. We confirmed through high-performance liquid chromatography analysis that aerobically grown stationary-phase cultures of three bacterial species, *Paracoccus denitrificans*, *Oligotropha carboxidovorans* and *Thermomicrobium roseum*, synthesized F_{420} , with oligoglutamate sidechains of different lengths. To understand the evolution of F_{420} biosynthesis, we also analyzed the distribution, phylogeny and genetic organization of the *cof* genes. Our data suggest that although the F_0 precursor to F_{420} originated in methanogens, F_{420} itself was first synthesized in an ancestral actinobacterium. F_{420} biosynthesis genes were then disseminated horizontally to archaea and other bacteria. Together, our findings suggest that the cofactor is more significant in aerobic bacterial metabolism and soil ecosystem composition than previously thought. The cofactor may confer several competitive advantages for aerobic soil bacteria by mediating their central metabolic processes and broadening the range of organic compounds they can synthesize, detoxify and mineralize.

The ISME Journal (2017) 11, 125–137; doi:10.1038/ismej.2016.100; published online 9 August 2016

Introduction

F_{420} serves as the cofactor in the catalysis of some of the most chemically demanding redox reactions in biology. Among them are the one-carbon reactions of methanogenesis (Thauer, 1998; Shima *et al.*, 2000; Hagemeyer *et al.*, 2003), the biosynthesis pathways of tetracycline antibiotics (Wang *et al.*, 2013) and the biodegradation of picrate and aflatoxins (Ebert *et al.*,

2001; Taylor *et al.*, 2010; Lapalika *et al.*, 2012). The cofactor appears to have been selected for these roles because of its unique electrochemical properties compared with the ubiquitous flavin and nicotinamide cofactors FMN (flavin mononucleotide), FAD (flavin adenine dinucleotide) and NAD(P) (nicotinamide adenine dinucleotide (phosphate)) (Walsh, 1986; Greening *et al.*, 2016a). As a 5-deazaflavin, F_{420} is structurally and biosynthetically related to FMN and FAD, but exhibits distinct electrochemical properties because of several key substitutions. It has a relatively low redox potential of -340 mV under standard conditions and -380 mV under certain physiological conditions (de Poorter *et al.*, 2005). This enables reduced F_{420} ($F_{420}H_2$) to reduce a wide range of organic compounds otherwise recalcitrant to activation (Jacobson and Walsh, 1984; Greening *et al.*, 2016a). As an obligate two-electron carrier, the cofactor can transform alkene, alkyne, alcohol and imine groups through hydride transfer reactions (Shima *et al.*, 2000; Hagemeyer *et al.*, 2003;

Correspondence: CJ Jackson, Research School of Chemistry, Australian National University, Sullivans Creek Road, Acton, Australian Capital Territory 2601, Australia.

E-mail: colin.jackson@anu.edu.au

or C Greening, The Commonwealth Scientific and Industrial Research Organisation, Land and Water Flagship, Clunies Ross Street, Acton, Australian Capital Territory 2601, Australia.

E-mail: chris.greening@csiro.au

⁵These two authors contributed equally to this work.

⁶Current address: Monash University, School of Biological Sciences, Clayton, Victoria, Australia.

Received 11 March 2016; revised 7 June 2016; accepted 13 June 2016; published online 9 August 2016

Aufhammer *et al.*, 2004; Shen *et al.*, 2009; Wang *et al.*, 2013).

Half a century since its discovery (Cheeseman *et al.*, 1972), F₄₂₀ is still perceived to be a rare cofactor synonymous with methanogenesis (Greening *et al.*, 2016a). It has been confirmed to be synthesized in just two phyla to date: the Euryarchaeota (Eirich *et al.*, 1979) and Actinobacteria (Eker *et al.*, 1980; Daniels *et al.*, 1985). It is thought to serve as the primary catabolic electron carrier in multiple lineages of Euryarchaeota, including representatives of the methanogenic (Eirich *et al.*, 1979), methanotrophic (Michaelis *et al.*, 2002; Knittel *et al.*, 2005) and sulfate-reducing orders (Lin and White, 1986). Genomic and spectroscopic evidence suggests that the cofactor is also synthesized in the aerobic ammonia-oxidizing phylum Thaumarchaeota (Spang *et al.*, 2012). Among bacteria, the cofactor has been chemically identified only within the Actinobacteria, where its physiological roles remain under investigation. In these organisms, F₄₂₀ reduction is coupled to either glucose 6-phosphate or NADPH oxidation and hence is dependent on the pentose phosphate pathway (Greening *et al.*, 2016a). The reduced cofactor (F₄₂₀H₂) is reported to enhance the metabolic flexibility of mycobacteria by facilitating the catalysis of a wide range of reductions of endogenous and exogenous organic compounds (Ahmed *et al.*, 2015; Greening *et al.*, 2016a). F₄₂₀ also has roles in antibiotic synthesis and xenobiotic degradation in species of *Streptomyces* (Wang *et al.*, 2013), *Rhodococcus* (Heiss *et al.*, 2002) and *Nocardioides* (Ebert *et al.*, 1999).

F₄₂₀ is synthesized in three major steps in bacteria and archaea (Figure 1). In the first, a riboflavin precursor (5-amino-6-(*D*-ribitylamino)uracil) is condensed with tyrosine to form 8-hydroxy-5-deazaflavin, also known as F_o; this step is catalyzed by the radical *S*-adenosylmethionine enzymes CofG and CofH that are fused into a single protein in some bacteria (known as CofGH or FbiC) (Choi *et al.*, 2002; Philmus *et al.*, 2015). Subsequently, LPPG (L-lactyl-2-diphospho-5'-guanosine) is proposed to be synthesized from 2-phospho-L-lactate by CofC (Grochowski *et al.*, 2008) and transferred to F_o by CofD (also known as FbiA) (Choi *et al.*, 2001; Graupner and White, 2001; Graupner *et al.*, 2002). The resulting LPPG sidechain is finally elongated with glutamate residues by the F₄₂₀: γ -L-glutamyl ligase CofE (also known as FbiB) (Choi *et al.*, 2001; Li *et al.*, 2003; Nocek *et al.*, 2007) that is fused with an FMN-dependent oxidoreductase in Actinobacteria (Bashiri *et al.*, 2016). For reasons still not understood, the number of glutamate residues added varies between organisms, ranging from two to three in most methanogens (Gorris and van der Drift, 1994), four to five in *Methanosarcina* (Gorris and van der Drift, 1994) and five to seven in *Mycobacterium* (Bair *et al.*, 2001). In addition to being an obligate intermediate in the F₄₂₀ biosynthesis pathway, F_o is also

synthesized independently by bacteria, archaea and eukaryotes for use as an antennal chromophore in DNA repair photolyases (Eker *et al.*, 1988; Yasui *et al.*, 1988; Kiener *et al.*, 1989; Epple and Carell, 1998).

We recently proposed that F₄₂₀ could be synthesized in a wider range of bacteria than currently described in the literature (Greening *et al.*, 2016a). Our analysis of a protein superfamily, the flavin/deazaflavin oxidoreductases, identified putative F₄₂₀-utilizing oxidoreductases in microorganisms other than the Euryarchaeota and Actinobacteria, including Chloroflexi, Proteobacteria and Firmicutes (Ahmed *et al.*, 2015). In this work, we explored the distribution of the genes encoding the F₄₂₀ biosynthesis enzymes CofC, CofD, CofE, CofG and CofH in public genomes and metagenomes. This revealed that the genes required to synthesize F₄₂₀ are encoded in a broad range of aerobic bacteria and are widespread in soil and aquatic ecosystems. Using this information, we validated through pure culture studies that three representatives of the dominant soil phyla Proteobacteria and Chloroflexi synthesize F₄₂₀. We propose that F₄₂₀ is much more widely distributed in microorganisms than previously reported and present a model of the evolution of the F_o and F₄₂₀ biosynthesis pathways to explain the origin and dispersal of this cofactor.

Materials and methods

Gene sequence retrieval

The amino acid sequences of all known F₄₂₀ biosynthesis enzymes (CofC, CofD, CofE, CofG and CofH) represented in the NCBI (National Center for Biotechnology Information) Reference Sequence (RefSeq) database (Pruitt *et al.*, 2007) were retrieved by Protein BLAST and PSI-BLAST (Altschul *et al.*, 1990). The homologous proteins in Aigarchaeota, Bathyarchaeota, Geoarchaeota, Lokiarchaeota and Tectomicrobia were retrieved from the Joint Genome Institute's Integrated Microbial Genomes database (Markowitz *et al.*, 2012). Taxonomic annotations for the sequences were obtained from the NCBI Taxonomy database and sequences duplicated at the species level were deleted. Clustering on sequence similarity networks (Atkinson *et al.*, 2009) generated using the Enzyme Function-Initiative Enzyme Similarity Tool (Gerlt *et al.*, 2015) were used to identify homologs of characterized proteins from nonspecific hits. In this analysis, nodes represent individual proteins and edges represent the all-versus-all BLAST *E*-values (Altschul *et al.*, 1990) between them. Closely related proteins form visual clusters, allowing the identification of sequences belonging to a protein family from those belonging to related families (Atkinson *et al.*, 2009). Final sequence sets were obtained by decreasing log*E*-value cutoffs until no major changes in clustering were observed with large increases in cutoff value. Final log*E*-value

Figure 1 Biosynthesis pathways for F₀ and F₄₂₀.

cutoffs used to identify sequences were –20 for CofC, –52 for CofE and –60 for CofD, CofG and CofH.

Evolutionary analysis

For robust phylogenetic tree construction, representative smaller sequence sets were generated by removing any sequences with >90% sequence identity using CD-HIT (Fu *et al.*, 2012). This was first done for CofH and the same taxa as in the resulting sequence set were used for the phylogenetic analysis of the other proteins for consistency. RogueNaRok (Aberer *et al.*, 2013) was used to remove fast-evolving sequences (all belonging to Firmicutes, Tectomicrobia, Thermoleophilia and Rubrobacteria) that otherwise caused long-branch attractions and hence unreliable evolutionary inferences (Anderson and Swofford, 2004). Sequences were aligned using MAFFT (Katoh and Standley, 2013) or MUSCLE (Edgar, 2004), with gaps and poorly aligned variable regions in the alignment removed using Gblocks (Castresana, 2000). Approximate maximum-likelihood trees were generated using Fasttree 2 (Price *et al.*, 2010), with the JTT +CAT evolutionary model for 100 bootstrap replicates that were generated using SeqBoot in the Phylib package (Felsenstein, 2005). To analyze co-

evolution of the F₄₂₀ biosynthesis proteins, phylogenetic tree topologies were compared and correlated using MirrorTree (Kann *et al.*, 2009). Genetic organization was compared using the Microbial Genomic Context Viewer (Overmars *et al.*, 2013) and the Integrated Microbial Genomes database (Markowitz *et al.*, 2012).

Motif analysis and homology modeling

BLAST and PSI-BLAST (Altschul *et al.*, 1990) were used to retrieve sequences encoding probable F₄₂₀-dependent oxidoreductases from the NCBI reference genomes of *Oligotropha carboxidovorans*, *Paracoccus denitrificans* and *Thermomicrobium roseum*. Characterized representatives of the 20 previously described F₄₂₀-dependent oxidoreductase enzyme families were used as seed sequences (Greening *et al.*, 2016a). Phyre2 (Kelley *et al.*, 2015) was used to model protein structures based on these sequences based on solved protein structures with highest percentage amino acid sequence identity. The quality of the models at the global (full protein) and local (F₄₂₀-binding site) scales were assessed using ProQ2 (Ray *et al.*, 2012). F₄₂₀-binding motifs were identified based on previous studies (Eguchi *et al.*, 1984; Purwantini and Daniels, 1998; Aufhammer *et al.*, 2004; Ahmed *et al.*, 2015).

Metagenome analysis

Metagenomes were screened for the presence of F₄₂₀ biosynthesis genes via a translated BLAST screen against the reference database as previously described (Greening *et al.*, 2016b). In all, 19 publicly available metagenomes (11 ecosystem types) were randomly subsampled to an equal depth of 4 million reads with minimum read length >140 nucleotides. To remove false positives, hits within the initial screen were further sieved by removing any result with a minimum percentage identity <60 or minimum query coverage <40 amino acids.

Bacterial culturing and harvesting

O. carboxidovorans OM5 (Meyer and Schlegel, 1978) was grown on carbon monoxide oxidizer media supplemented with 36 mM sodium acetate (Meyer and Schlegel, 1978) and maintained on solid carbon monoxide oxidizer containing 1.2% (w/v) agar. *P. denitrificans* strain PD1222 (de Vries *et al.*, 1989) was grown and maintained in lysogeny broth liquid media and agar. *Mycobacterium smegmatis* mc²155 (Snapper *et al.*, 1990) was grown and maintained in lysogeny broth liquid media and agar supplemented with 0.05% (v/v) Tween 80. *T. roseum* (Jackson *et al.*, 1973) cultures were grown in Castenholz salts solution supplemented with 5 g l⁻¹ peptone and 2.5 g l⁻¹ sucrose (Houghton *et al.*, 2015). Liquid cultures (500 ml) of *O. carboxidovorans*, *P. denitrificans* and *M. smegmatis* were grown to early stationary-phase in 2-l Erlenmeyer flasks in a rotary incubator (200 r.p.m., 37 °C). *T. roseum* was grown in an equivalent manner in 1-l cultures at 130 r.p.m. and 68 °C. Cells from stationary-phase liquid cultures were harvested by centrifugation (4 °C, 10 000 g, 20 min) and the supernatant was discarded. Pellets were resuspended in 1 to 3 ml 50 mM sodium phosphate buffer (pH 7.0) and lysed by boiling. Lysates were centrifuged at 11 000 g for 10 min, and the supernatant was removed for analysis.

HPLC and liquid chromatography/mass spectrometry analysis

F₄₂₀ was detected by ion-pair reversed-phase high-performance liquid chromatography (HPLC) using an Agilent (Santa Clara, CA, USA) 1200 series HPLC system equipped with an autosampler, fluorescence detector and a Poroshell 120 EC-C18 2.1 × 100 mm, 2.7 µm column. Gradients of two HPLC buffers were used, A (20 mM ammonium phosphate, 10 mM tetrabutylammonium phosphate, pH 7.0) and B (100% acetonitrile), to separate F₄₂₀ species at high resolution by the length of their oligoglutamate tails. The applied gradient was 0–1 min 25% B; 1–10 min from 25% to 35% B; 10–13 min 35% B; 13–16 min from 35% to 40% B; 16–19 min from 40% to 25% B. Columns were extensively washed between sample runs, and test runs validated there was no carryover of F₄₂₀. For fluorescence detection, the samples were excited at 420 nm and emission spectra were recorded between 470 and 600 nm. To verify the

relative abundance of F₄₂₀ species, concentrated F₄₂₀ standard was purified from a recombinant F₄₂₀ overexpression strain of *M. smegmatis* mc²4517 (Bashiri *et al.*, 2010) as previously described (Isabelle *et al.*, 2002). The standard was lyophilized for storage, resuspended in 50 µl ultrapure water and serially diluted. The sample was analyzed on a Waters (Milford, MA, USA) LCT Premier OA-TOF (orthogonal acceleration time-of-flight) mass spectrometer. Samples were ionized by electrospray in negative ion mode at a rate of 5 µl per min with a capillary voltage of 2500 V, desolvation temperature of 150 °C and source temperature of 100 °C. Deconvolution of the mass data was facilitated by MaxEnt software (Princeton, NJ, USA).

Results

The genetic determinants of F₄₂₀ biosynthesis are widely encoded by aerobic soil bacteria

We initially retrieved the sequences of all F₄₂₀ biosynthesis genes in publicly available genomes in order to understand the distribution and evolution of F₄₂₀ in microorganisms (Supplementary Table S1). A total of 653 bacterial and 173 archaeal species named in the NCBI Reference Sequence database encoded all five proteins specifically required to synthesize F₄₂₀ (CofC, CofD, CofE, CofG and CofH) (Supplementary Table S2).

Among archaea, the genes encoding F₄₂₀ biosynthesis enzymes were unsurprisingly widespread in the Euryarchaeota, specifically in all six validated methanogenic orders, Halobacteria and Archaeoglobi. Biosynthesis genes were also detected in full suites in the Thaumarchaeota and Geoarchaeota, and in partial repertoires in the metagenome-derived incomplete genomes of Aigarchaeota, Bathyarchaeota and Lokiarchaeota representatives (Figure 2a and Supplementary Table S3). Among bacteria, F₄₂₀ biosynthesis genes were present in the majority of sequenced Actinobacteria, as well as multiple species within the phyla Proteobacteria (classes: Alpha, Beta and Gamma) and Chloroflexi (classes: Thermomicrobia, Ktedonobacteria and Ardenticatenia) (Figure 2a and Supplementary Table S3). Within these phyla often predominating in soil environments, species harboring *cof* genes were obligate aerobes and facultative anaerobes rather than obligate anaerobes. For example, these genes were widespread in the aerobic lineages of Chloroflexi, but not anaerobic lineages such as the Dehalococcoidia. In addition, *cof* genes were disproportionately abundant in known soil- and marine-dwelling orders of Proteobacteria (for example, Rhizobiales and Alteromonadales) compared with host-associated orders (for example, Rickettsiales and Enterobacteriales) (Supplementary Table S1). Finally, F₄₂₀ biosynthesis genes were also detected in two Firmicutes species and within the Tectomicrobia (Supplementary Table S3), a recently discovered

Figure 2 Genomic and metagenomic distribution of the *cof* genes that encode F₄₂₀ biosynthesis enzymes. The genes encoding the five known proteins specifically required for F₄₂₀ synthesis are shown, namely CofC, CofD, CofE, CofG and CofH, where CofGH represents a fusion protein. (a) Distribution of *cof* genes by phyla in the NCBI Reference Sequence database. (b) Distribution of *cof* genes in 19 publicly available metagenomes by enzyme. (c) Distribution of *cof* genes in 19 publicly available metagenomes by phylum of the closest BLAST hit (>60% identity).

candidate phylum with wide biosynthetic capacity (Wilson *et al.*, 2014). Putative F₄₂₀-dependent oxidoreductases from multiple families were identified in the genomes of the phyla predicted to synthesize F₄₂₀ (Supplementary Table S4).

We also analyzed the distribution of the two genes (*cofG* and *cofH*) required for synthesis of F_o, both an independent chromophore and an F₄₂₀ precursor (Figure 2a and Supplementary Table S3). It has been postulated that F_o is universally distributed given it is used in DNA photolyases across the three domains of life (Kiener *et al.*, 1989; Mees *et al.*, 2004; Petersen and Ronan, 2010). Our analysis revealed, however, that *cofG* and *cofH* genes were almost exclusively encoded by predicted F₄₂₀ producers, that is, those organisms that harbor all five F₄₂₀ biosynthesis genes. The only exceptions were Cyanobacteria, Chlorophyta and Streptophyta (Supplementary Table S1) that encoded the genetic determinants of F_o biosynthesis (*cofG* and *cofH*) but not F₄₂₀ synthesis (*cofC*, *cofD* and *cofE*). These three lineages comprise oxygenic phototrophs known to encode F_o-utilizing photolyases (Mees *et al.*, 2004; Petersen and Ronan, 2010). As previously observed (Graham *et al.*, 2003), the *cofG* and *cofH* genes are fused into a single open reading frame (referred hereafter as *cofGH*) in most Actinobacteria, some Proteobacteria and phototrophic eukaryotes.

To explore the ecological role of F₄₂₀, we surveyed publicly available metagenomes for the genetic determinants of F₄₂₀ synthesis (Supplementary Table S5). In total, a full complement of F₄₂₀ biosynthesis genes were identified in all 17 soil,

water and sediment metagenomes analyzed (Supplementary Table S5). These genes were especially abundant in the aerated forest and agricultural soil samples, accounting for ~0.02% of total sequence reads (Figure 2b). Consistent with the microbial community composition of soil ecosystems (Janssen, 2006), the majority of the identified reads matched those from Actinobacteria, Alphaproteobacteria and Chloroflexi (Figure 2c). Based on metagenomes, F₄₂₀ biosynthesis genes were also abundant in eight aquatic environments. In these ecosystems, genes encoding F_o synthases (*cofG* and *cofH*) were 1.4 to 1.8 times more abundant relative to those encoding F₄₂₀-specific biosynthesis enzymes (*cofC*, *cofD* and *cofE*) (Figure 2b), suggesting that these environments harbor large communities of both F_o-utilizing phototrophs and F₄₂₀-synthesizing chemotrophs (Figure 2c). F₄₂₀ biosynthesis genes closely related to those encoded in Euryarchaeota were unsurprisingly common in marine sediments known to harbor high concentrations of methane-cycling archaea (Figure 2c). Consistent with their community composition, F₄₂₀ biosynthesis genes were detected in very low abundance within gut ecosystems (Supplementary Table S5).

Representatives of the Proteobacteria and Chloroflexi synthesize F₄₂₀

To validate these genome-based predictions, we investigated whether isolates of the phyla Proteobacteria and Chloroflexi synthesized F₄₂₀. We cultured three bacteria to test for F₄₂₀ production

under conditions that would promote aerobic, heterotrophic growth: *O. carboxidovorans*, an obligate aerobe and facultative lithotroph of the alphaproteobacterial order Rhizobiales (Meyer and Schlegel, 1978); *P. denitrificans*, a facultative anaerobe and facultative lithotroph of the alphaproteobacterial order Rhodobacterales (de Vries *et al.*, 1989); and *T. roseum*, a thermophilic obligate aerobe and obligate heterotroph of the phylum Chloroflexi (Jackson *et al.*, 1973). F_{420} was detected in whole-cell lysates of all three bacteria as well as within the positive control *M. smegmatis* (Figure 3). In HPLC analysis, F_{420} species from all four organisms eluted with retention times (Figure 3a) identical to those from a purified F_{420} standard validated by mass spectrometry (Figure 3b). When excited at 420 nm, all samples emitted fluorescence ($\lambda_{\text{max}} = 480$ nm) (Figure 3c) with a spectrum characteristic of F_{420} (Eirich *et al.*, 1978).

The quantity of F_{420} produced and distribution of oligoglutamate chain lengths differed between the strains analyzed (Table 1). Whereas *T. roseum* and *M. smegmatis* produced large quantities of F_{420} , normalized production was ~ 100 -fold lower for the proteobacterial strains under comparable conditions (Figure 3a); F_{420} was nevertheless unambiguously and reproducibly detected in such organisms. Ion-pair HPLC was used to resolve the F_{420} species based on the length of their oligoglutamate side chains (Figure 3a). Mass spectrometry of the purified standard confirmed the lengths of the dominant HPLC peaks (Figure 3b). The number of glutamate residues attached differed between species, with 4 to 6 glutamates detected in *P. denitrificans*, 5 to 7 detected in *O. carboxidovorans* and 6 to 8 dominating in the *T. roseum* and *M. smegmatis* samples (Table 1).

Having established that species from the phyla Proteobacteria and Chloroflexi synthesize F_{420} ,

Figure 3 Chemical detection of F_{420} in dominant soil phyla. (a) HPLC traces showing F_{420} from cell lysates of different species relative to a purified F_{420} standard. The traces show the intensity of the fluorescence emitted ($\lambda_{\text{excitation}} = 420$ nm, $\lambda_{\text{emission}} = 480$ nm). The dotted lines show the times at which each species started to elute. (b) Mass spectra confirming the molecular weight and oligoglutamate side chain length of the purified F_{420} standard. (c) Fluorescence emission spectra of F_{420} from cell lysates of five different species against a purified F_{420} standard.

Table 1 Length of oligoglutarate chains of F₄₂₀ in the cultured organisms

F ₄₂₀ species (%)	F ₄₂₀ -2	F ₄₂₀ -3	F ₄₂₀ -4	F ₄₂₀ -5	F ₄₂₀ -6	F ₄₂₀ -7	F ₄₂₀ -8	F ₄₂₀ -9
F ₄₂₀ standard	<1	<1	1	1	9	45	36	7
<i>Oligotropha carboxidovorans</i>	—	2	7	14	32	37	8	—
<i>Paracoccus denitrificans</i>	—	—	13	82	5	—	—	—
<i>Thermomicrobium roseum</i>	—	<1	<1	3	20	39	34	3
<i>Mycobacterium smegmatis</i>	—	1	2	5	35	50	7	<1

The percentage amount of each F₄₂₀ species compared with the total F₄₂₀ present is shown.

we investigated whether any F₄₂₀-dependent enzymes that have been characterized in Archaea and Actinobacteria are conserved in the genomes of *P. denitrificans*, *O. carboxidovorans* and *T. roseum*. We identified close homologs of Fgd (F₄₂₀-reducing glucose 6-phosphate dehydrogenase) in the Chloroflexi and Fno (F₄₂₀-reducing NADPH dehydrogenase) in the two Proteobacteria. Putative F₄₂₀H₂-dependent reductases from two superfamilies, the luciferase-like hydride transferases (Greening *et al.*, 2016a) and flavin/deazaflavin oxidoreductases (Ahmed *et al.*, 2015), were also encoded in the three organisms (Supplementary Table S6). Multiple sequence alignments and protein homology models confirmed the residues involved in F₄₂₀ binding are almost entirely conserved in these sequences, including residues that interact with the isoalloxazine ring and oligoglutarate chain, that are not found in the flavins FAD and FMN (Supplementary Figure S1). The presence of these sequences and the high conservation of the F₄₂₀-binding sequence motifs provide support that these organisms encode F₄₂₀-dependent oxidoreductases.

F₄₂₀ biosynthesis pathways evolved through bacterial-to-archaeal horizontal gene transfers

Finally, we analyzed the evolution of the F₄₂₀ biosynthesis enzymes (CofC, CofD, CofE, CofG and CofH) to understand the origin and distribution of the cofactor. With respect to all five proteins, phylogenetic tree topologies (Figure 4a) and sequence similarity networks (Supplementary Figure S2) were similar. In all cases, two large clades corresponding to the euryarchaeotal and actinobacterial enzymes flanked the trees and networks, and there were smaller middle clades corresponding to homologous enzymes in Proteobacteria, Chloroflexi, Tectomicrobia, Firmicutes, Thaumarchaeota and deep-branching actinobacterial lineages such as Acidimicrobiia. A MirrorTree analysis comparing the topologies of the phylogenetic trees indicated that the five Cof enzymes co-evolved (with Pearson's correlation coefficients between 0.71 and 0.93) (Supplementary Figure S3). This is consistent with selection for all five subunits being required to produce functional F₄₂₀. The F₀ biosynthesis proteins CofG and CofH in Cyanobacteria have diverged further from their homologs in F₄₂₀-synthesizing organisms (Supplementary Figures S2 and S4),

consistent with their lack of co-evolution with the rest of the pathway.

To better understand the evolutionary origin of these proteins, phylogenetic trees were rooted with related families containing the same protein fold (Figure 4a), namely biotin synthase (a radical S-adenosylmethionine enzyme) for CofG and CofH, adenosylcobinamide guanosine transferase (MobA) for CofC and YvcK (hypothetical protein family) for CofD. CofE adopts a novel protein fold and does not share any significant sequence homology with any other known proteins (Nocek *et al.*, 2007) and thus is presented as an unrooted tree with no evolutionary origin inferred. Our analysis indicates that CofG is likely to have originated in an ancestral archaeon. Subsequently, the functionally and structurally related CofH evolved from an ancient duplication of the *cofG* gene within an early methanogenic euryarchaeon. In contrast, the roots of the CofC and CofD trees indicate these enzymes originated in an early actinobacterium. These findings are consistent with the recent work suggesting *cofC* and *cofD* were among the metabolic genes acquired from bacteria during the diversification of archaea (Nelson-Sathi *et al.*, 2015). Our observations suggest that, in contrast to the F₀ synthases, enzymes specific for F₄₂₀ biosynthesis evolved in bacteria, most probably an early actinobacterium, and were subsequently horizontally transferred into archaea.

The genetic organization of the *cof* genes is highly variable (Figure 4b). The five opening reading frames are genomically separated in two methanogenic orders (Methanococcales and Methanomicrobiales) and are partially clustered in Actinobacteria, Chloroflexi, Tectomicrobia and all other archaea. In contrast, *cof* genes are organized as complete operons within Proteobacteria; consistent with the presence of two distinct clusters of proteobacterial sequences in all phylogenetic trees, these operons were organized in two distinct configurations (designated Proteobacteria I and Proteobacteria II). Our analysis suggests that the *cof* genes initially evolved separately, but became increasingly syntenic because of selective pressures to inherit all five enzymes for F₄₂₀ production. This analysis shows that the deepest-rooted branches of the CofGH fusion protein are between the classes Actinobacteria and Acidimicrobiia. This suggests the CofGH fusion occurred early in actinobacterial evolution, presumably to enhance the catalytic efficiency of the

Figure 4 Evolution of the determinants of F_{420} biosynthesis. (a) Phylogenetic trees of the F_{420} biosynthesis proteins (CofC, CofD, CofE, CofG and CofH). Trees were rooted with a related protein family characterized by the same protein fold, except for CofE (novel protein fold), presented as an unrooted tree. Clades are labeled according to phyla, except for Actinobacteria and Euryarchaeota that are labeled by class. The Firmicutes, Tectomicrobia, Thermoleophilia and Rubrobacteria lineages have been omitted as their inclusion compromises phylogenetic inferences because of long-branch attractions. Gray-shaded regions represent the CofGH fusion proteins. CofG and CofH trees incorporating cyanobacterial sequences are presented in Supplementary Figure S4 as these sequences caused low bootstrap values at key nodes. (b) Generalized schematic of the genetic organization of the genes encoding the five enzymes specifically required for F_{420} biosynthesis (CofC, CofD, CofE, CofG and CofH) from five bacterial phyla (Actinobacteria, Proteobacteria, Chloroflexi, Tectomicrobia and Firmicutes) and three archaeal phyla (Euryarchaeota, Thaumarchaeota and Geoarchaeota). Fgd, F_{420} -reducing glucose 6-phosphate dehydrogenase; Fno, F_{420} -reducing NADPH dehydrogenase; HFDR, predicted $F_{420}H_2$ -dependent reductase; LLHT, predicted $F_{420}H_2$ -dependent luciferase-like oxidoreductase; Mer, methylenetetrahydromethanopterin reductase; NR, hypothetical nitroreductase. (c) Schematic representation of the proposed evolutionary origin of F_{420} biosynthesis genes and their acquisition by different phyla. Solid lines/circles indicate vertical acquisition, whereas dashed lines/circles indicate horizontal acquisition. The capacity for F_{420} biosynthesis appears to have evolved on at least two separate occasions in both the Actinobacteria (Actinobacteria I = Actinobacteria, Acidimicrobiia; Actinobacteria II = Thermoleophilia, Rubrobacteria) and Proteobacteria (Proteobacteria I = Rhizobiales, Sphingomonadales, Betaproteobacteria; Proteobacteria II = Rhodobacterales, Gammaproteobacteria).

coordinated radical *S*-adenosylmethionine reactions required for F_0 synthesis (Philmus *et al.*, 2015). In some organisms, F_{420} -reducing dehydrogenases (Fgd, Fno and Mer) and putative $F_{420}H_2$ -dependent reductases also appear to be operonic with the *cof* genes (Figure 4b).

Discussion

Until now, the vast majority of research into F_{420} has focused on the roles of the cofactor in methane

cycling and tuberculosis infection (Greening *et al.*, 2016a). However, our findings demonstrate that the cofactor is widely distributed among diverse taxa and ecosystems, and is likely to play a role in a broader array of metabolic and ecological phenomena than is currently recognized. We have demonstrated, using a combination of comparative genomics and analytical chemistry approaches, that F_{420} is widely distributed in three of the five most dominant soil phyla (Actinobacteria, Proteobacteria and Chloroflexi) and provided genomic evidence that the cofactor is encoded in other phyla

(Tectomicrobia, Firmicutes, Thaumarchaeota and Geoarchaeota). The finding that F₄₂₀ biosynthesis genes are widely distributed within aerobic bacterial and archaeal taxa indicates that F₄₂₀ is far from a niche methanogenic cofactor. Furthermore, the abundance of these genes in soil ecosystems suggests that F₄₂₀ influences the biological and chemical composition of soils.

There are three feasible explanations for the observed distribution of F₄₂₀ in biological systems: (1) F₄₂₀ biosynthesis evolved pre-LUCA (last universal common ancestor); (2) F₄₂₀ biosynthesis evolved in an archaeon and was acquired by bacteria; or (3) F₄₂₀ biosynthesis evolved in a bacterium and was acquired by archaea. Of these, the first hypothesis is unsupported given that *cof* gene evolution does not parallel 16S rRNA gene phylogenies and the cofactor appears to be completely absent from proposed deep-branching bacteria (for example, Aquificae and Thermotogae) and archaea (for example, Crenarchaeota and Thermococci). The widely assumed archaea-to-bacteria hypothesis remains plausible, but is challenged by phylogenetic trees suggesting a bacterial origin for CofC and CofD. The bacteria-to-archaea hypothesis for F₄₂₀ acquisition is more robustly supported by our phylogenetic trees and is also consistent with the findings of the large-scale Martin analysis on interdomain lateral gene transfers (Nelson-Sathi *et al.*, 2015).

In Figure 4c, we present an evolutionary model of the origin and distribution of F₄₂₀. Our studies strongly support that the genes required for F_o synthesis (*cofG* and *cofH*) evolved independently and probably earlier than the genes required for F₄₂₀ synthesis from F_o (*cofC*, *cofD* and *cofE*). We hypothesize that an early methanogen synthesized F_o but not F₄₂₀ using the products of ancient *cofG* and *cofH* genes. This cofactor may have initially sustained many of the redox functions now supported by F₄₂₀, including hydrogenotrophic methanogenesis. Our evolutionary analysis suggests that an early actinobacterium subsequently acquired *cofG* and *cofH* (later fused into *cofGH*) and evolved the *cofC*, *cofD* and likely *cofE* genes required to produce F₄₂₀, the lactyloligoglutamyl derivative of F_o. Multiple horizontal gene transfer events thereafter led to the acquisition of these genes in the archaea and other bacteria. The euryarchaeotal branches of the CofC, CofD, and CofE phylogenetic trees mirror those of 16S rRNA gene trees (Hedderich and Whitman, 2013). This is consistent with their genes being acquired from Actinobacteria early in the evolution of Euryarchaeota (Figure 4), resulting in the capacity for F₄₂₀ biosynthesis being vertically inherited by the six methanogenic orders, Archaeoglobi, Halobacteria and likely anaerobic methanotrophs (ANME). Unicellular eukaryotes (Chlorophyta and Streptophyta) appear to have acquired the chromophore F_o by laterally acquiring the fused *cofGH* gene from Proteobacteria. The origin of the cyanobacterial *cofG* and *cofH* remains unclear because of their

evolutionary distance from the rest of the sequences; scenarios involving acquisition from methanogens or Proteobacteria are both plausible.

We propose that the evolutionary driving force resulting in F_o synthesis was the need for redox cofactors with distinct electrochemical properties to flavins and nicotinamides to drive the unique reactions of hydrogenotrophic methanogenesis. By hijacking the riboflavin biosynthesis pathway with CofG and CofH, an early euryarchaeon would have produced a cofactor with the ideal electrochemistry to mediate the central reactions of methanogenesis: H₂ oxidation, methenyl and methylene reduction and NADP reduction. F_o could have functioned as the primordial methanogenic redox cofactor as it has near-identical electrochemical properties to F₄₂₀, including a standard redox potential of −340 mV and obligate two-electron reactivity (Walsh, 1986; Greening *et al.*, 2016a). Furthermore, it has been demonstrated that F_o can functionally substitute for F₄₂₀ in various enzymatic processes *in vitro*, including in the crucial methanogenesis enzymes Frh (F₄₂₀-reducing hydrogenase) and Fno (F₄₂₀-dependent NADP reductase) (Yamazaki and Tsai, 1980; Muth *et al.*, 1987). The current distribution of *cof* genes emphasizes that deazaflavins have primarily been selected for their electrochemical and not photochemical properties. With the exception of cyanobacterial and eukaryotic phototrophs, all organisms appear to have retained *cofG* and *cofH* genes primarily to generate F₄₂₀ as a redox cofactor rather than F_o as an independent chromophore. We propose that the need to produce a charged derivative of F_o, F₄₂₀, served as the main selection pressure for the evolution and dispersal of CofC, CofD and CofE. The feature that distinguishes F₄₂₀ from its precursor F_o is the presence of a negatively charged sidechain containing phosphate and glutamate groups (Eirich *et al.*, 1978). Thus, whereas F_o diffuses through membranes, F₄₂₀ is retained in the cytosol (Glas *et al.*, 2009). The electrostatic properties of F₄₂₀ may also facilitate higher affinity enzyme–cofactor interactions according to recent structural studies (Greening *et al.*, 2016a). We also show here that the length of the oligoglutamate sidechain of F₄₂₀ varies between phyla and progressively increases between the Euryarchaeota (Gorris and van der Drift, 1994), Proteobacteria, Actinobacteria and Chloroflexi (Table 1). We are presently investigating the reasons and mechanisms behind these variations.

F₄₂₀ may confer a number of competitive advantages to these bacteria by mediating both endogenous and exogenous metabolic processes. F₄₂₀ is known to serve as a redox cofactor in an increasing number of endogenous processes in mycobacteria, including central carbon metabolism (Bashiri *et al.*, 2008), cell wall modification (Purwantini and Mukhopadhyay, 2013), antioxidant production (Ahmed *et al.*, 2015) and possibly quinone reduction (Gurumurthy *et al.*, 2013). Although the cofactor is synthesized in high

levels under oxic conditions (Figure 2), phenotypic evidence suggests it is particularly important for survival under hypoxia (Gurumurthy *et al.*, 2013). In this condition, mycobacteria may increasingly depend on low-potential cofactors such as F₄₂₀ to maintain redox homeostasis (Berney *et al.*, 2014; Cook *et al.*, 2014). Many of the F₄₂₀H₂-dependent oxidoreductases involved in mycobacterial metabolism are also conserved in other taxa, where they may have similar roles (Supplementary Table S5). F₄₂₀ is likely to be particularly important in the redox metabolism of Chloroflexi and other Actinobacteria given representatives of such phyla produce large amounts of the compound even under optimal growth conditions (Figure 3a). A central role for F₄₂₀ in the metabolism of Proteobacteria seems less likely, given the levels detected of the cofactor in *O. carboxidovorans* (Proteobacteria I lineage) and *P. denitrificans* (Proteobacteria II lineage) were low (Figure 3a). It is likely that transcription of the *cof* genes is constitutively repressed in such organisms, but is activated in response to environmental stresses or other signals. Consistently, TetR and MarR family transcriptional repressors were identified immediately downstream of the *cof* operons in most Proteobacteria, including *P. denitrificans* and *O. carboxidivorans*.

The richest literature on the physiological roles of F₄₂₀ in bacteria concerns the metabolism of complex organic compounds. F₄₂₀H₂-dependent reductases mediate the biodegradation of nitroaromatic explosives (Ebert *et al.*, 2001), triphenyl dyes (Guerra-Lopez *et al.*, 2007) and furanocoumarins (Taylor *et al.*, 2010), as well as the biosynthesis of tetracycline and pyrrolobenzodiazepine antibiotics (Li *et al.*, 2009; Wang *et al.*, 2013). A role for F₄₂₀ in equivalent processes seems particularly likely for the Chloroflexi and Tectomicrobia; both phyla contain an abundance of F₄₂₀H₂-dependent reductases (Supplementary Tables S3 and S6) and are reputed for their biosynthetic versatility and biodegradative capacities (Björnsson *et al.*, 2002; Wilson *et al.*, 2014). In turn, these F₄₂₀-dependent processes are likely to affect the biological composition of soil ecosystems, for example, by shaping the antibiotics arms race. They may also influence the chemical composition of soil environments by controlling the levels of complex organic compounds produced and consumed. Better understanding of the endogenous and exogenous roles of F₄₂₀ depends on defining the functions of the numerous putative F₄₂₀-dependent oxidoreductases encoded in aerobic bacteria and archaea (Supplementary Table S3). In turn, further understanding the physiological and ecological significance of F₄₂₀ may open opportunities in the fields of pharmaceuticals, biocatalysis and bioremediation.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

We thank Dr Ghader Bashiri (University of Auckland) for providing the *fbiABC* overexpression plasmid, Professor William Jacobs Jr (Albert Einstein College of Medicine) for supplying *Mycobacterium smegmatis* mc²4517 and the three anonymous reviewers for their helpful suggestions. This work was supported by Australian Research Council research Grants (DE120102673, DP130102144) awarded to CJJ; Australian National University Summer and PhD scholarships awarded respectively to BN and FHA; and a CSIRO Office of the Chief Executive Postdoctoral Fellowship awarded to CG.

Author contributions

CG, CJJ, BN, FHA, CRC, MBS, MCT, ACW, JGO, SJW and GP designed experiments. BN, FHA, CG, CRC, AB, SJW, and CJJ performed experiments. CG, CJJ, BN and FHA analyzed data. CJJ, CG, MCT, ACW, JGO and SEM supervised students and researchers. CG, CJJ, FHA, BN and JGO wrote the paper.

References

- Aberer AJ, Krompass D, Stamatakis A. (2013). Pruning rogue taxa improves phylogenetic accuracy: an efficient algorithm and webservice. *Syst Biol* **62**: 162–166.
- Ahmed FH, Carr PD, Lee BM, Afriat-Jurnou L, Mohamed AE, Hong N-S *et al.* (2015). Sequence-structure-function classification of a catalytically diverse oxidoreductase superfamily in mycobacteria. *J Mol Biol* **427**: 3554–3571.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. (1990). Basic local alignment search tool. *J Mol Biol* **215**: 403–410.
- Anderson FE, Swofford DL. (2004). Should we be worried about long-branch attraction in real data sets? Investigations using metazoan 18S rDNA. *Mol Phylogenet Evol* **33**: 440–451.
- Atkinson HJ, Morris JH, Ferrin TE, Babbitt PC. (2009). Using sequence similarity networks for visualization of relationships across diverse protein superfamilies. *PLoS One* **4**: e4345.
- Aufhammer SW, Warkentin E, Berk H, Shima S, Thauer RK, Ermmler U. (2004). Coenzyme binding in F₄₂₀-dependent secondary alcohol dehydrogenase, a member of the bacterial luciferase family. *Structure* **12**: 361–370.
- Bair TB, Isabelle DW, Daniels L. (2001). Structures of coenzyme F₄₂₀ in *Mycobacterium* species. *Arch Microbiol* **176**: 37–43.
- Bashiri G, Rehan AM, Greenwood DR, Dickson MJ, Baker EN. (2010). Metabolic engineering of cofactor F₄₂₀ production in *Mycobacterium smegmatis*. *PLoS One* **5**: e15803.
- Bashiri G, Rehan AM, Sreebhavan S, Baker HM, Baker EN, Squire CJ. (2016). Elongation of the poly-gamma-glutamate tail of F₄₂₀ requires both domains of the F₄₂₀:gamma-glutamyl ligase (FbiB) of *Mycobacterium tuberculosis*. *J Biol Chem* **291**: 6882–6894.
- Bashiri G, Squire CJ, Moreland NJ, Baker EN. (2008). Crystal structures of F₄₂₀-dependent glucose-6-

- phosphate dehydrogenase FGD1 involved in the activation of the anti-tuberculosis drug candidate PA-824 reveal the basis of coenzyme and substrate binding. *J Biol Chem* **283**: 17531–17541.
- Berney M, Greening C, Conrad R, Jacobs WR, Cook GM. (2014). An obligately aerobic soil bacterium activates fermentative hydrogen production to survive reductive stress during hypoxia. *Proc Natl Acad Sci USA* **111**: 11479–11484.
- Björnsson L, Hugenholtz P, Tyson GW, Blackall LL. (2002). Filamentous *Chloroflexi* (green non-sulfur bacteria) are abundant in wastewater treatment processes with biological nutrient removal. *Microbiology* **148**: 2309–2318.
- Castresana J. (2000). Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* **17**: 540–552.
- Cheeseman P, Toms-Wood A, Wolfe RS. (1972). Isolation and properties of a fluorescent compound, Factor420, from *Methanobacterium* strain M.o.H. *J Bacteriol* **112**: 527–531.
- Choi K-P, Bair TB, Bae Y-M, Daniels L. (2001). Use of transposon Tn5367 mutagenesis and a nitroimidazopyran-based selection system to demonstrate a requirement for *fbtA* and *fbtB* in coenzyme F₄₂₀ biosynthesis by *Mycobacterium bovis* BCG. *J Bacteriol* **183**: 7058–7066.
- Choi K-P, Kendrick N, Daniels L. (2002). Demonstration that *fbtC* is required by *Mycobacterium bovis* BCG for coenzyme F₄₂₀ and F_O biosynthesis. *J Bacteriol* **184**: 2420–2428.
- Cook GM, Greening C, Hards K, Berney M. (2014). Energetics of pathogenic bacteria and opportunities for drug development. In: Poole RK (ed) *Advances in Bacterial Pathogen Biology*, Vol. 65. Academic Press: Cambridge, MA, USA, pp 1–62.
- Daniels L, Bakhiet N, Harmon K. (1985). Widespread distribution of a 5-deazaflavin cofactor in *Actinomyces* and related bacteria. *Syst Appl Microbiol* **6**: 12–17.
- de Poorter LMI, Geerts WJ, Keltjens JT. (2005). Hydrogen concentrations in methane-forming cells probed by the ratios of reduced and oxidized coenzyme F₄₂₀. *Microbiology* **151**: 1697–1705.
- de Vries GE, Harms N, Hoogendijk J, Stouthamer AH. (1989). Isolation and characterization of *Paracoccus denitrificans* mutants with increased conjugation frequencies and pleiotropic loss of a (nGATCn) DNA-modifying property. *Arch Microbiol* **152**: 52–57.
- Ebert S, Fischer P, Knackmuss HJ. (2001). Converging catabolism of 2,4,6-trinitrophenol (picric acid) and 2,4-dinitrophenol by *Nocardioides simplex* FJ2-1A. *Biodegradation* **12**: 367–376.
- Ebert S, Rieger P-G, Knackmuss H-J. (1999). Function of coenzyme F₄₂₀ in aerobic catabolism of 2,4,6-trinitrophenol and 2,4-dinitrophenol by *Nocardioides simplex* FJ2-1A. *J Bacteriol* **181**: 2669–2674.
- Edgar RC. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**: 1792–1797.
- Eguchi S, Nakata H, Nishio N, Nagai S. (1984). NADP⁺ reduction by a methanogen using HCOOH or H₂ as electron donor. *Appl Microbiol Biotechnol* **20**: 213–217.
- Eirich LD, Vogels GD, Wolfe RS. (1979). Distribution of coenzyme F₄₂₀ and properties of its hydrolytic fragments. *J Bacteriol* **140**: 20–27.
- Eirich LD, Vogels GD, Wolfe RS. (1978). Proposed structure for coenzyme F₄₂₀ from *Methanobacterium*. *Biochemistry* **17**: 4583–4593.
- Eker APM, Hessels JKC, van de Velde J. (1988). Photo-reactivating enzyme from the green alga *Scenedesmus acutus*. Evidence for the presence of two different flavin chromophores. *Biochemistry* **27**: 1758–1765.
- Eker APM, Pol A, van der Meyden P, Vogels GD. (1980). Purification and properties of 8-hydroxy-5-deazaflavin derivatives from *Streptomyces griseus*. *FEMS Microbiol Lett* **8**: 161–165.
- Eppl R, Carell T. (1998). Flavin- and deazaflavin-containing model compounds mimic the energy transfer step in type-II DNA-photolyases. *Angew Chemie Int Ed* **37**: 938–941.
- Felsenstein J. (2005). PHYLIP (phylogeny inference package) Distributed by the author. *Dep Genome Sci Univ Washington, Seattle*, Version 3.
- Fu L, Niu B, Zhu Z, Wu S, Li W. (2012). CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* **28**: 3150–3152.
- Gerlt JA, Bouvier JT, Davidson DB, Imker HJ, Sadkhin B, Slater DR et al. (2015). Enzyme Function Initiative-Enzyme Similarity Tool (EFI-EST): a web tool for generating protein sequence similarity networks. *Biochim Biophys Acta* **1854**: 1019–1037.
- Glas AF, Maul MJ, Cryle M, Barends TRM, Schneider S, Kaya E et al. (2009). The archaeal cofactor F_O is a light-harvesting antenna chromophore in eukaryotes. *Proc Natl Acad Sci USA* **106**: 11540–11545.
- Gorris LG, van der Drift C. (1994). Cofactor contents of methanogenic bacteria reviewed. *Biofactors* **4**: 139–145.
- Graham DE, Xu H, White RH. (2003). Identification of the 7,8-didemethyl-8-hydroxy-5-deazariboflavin synthase required for coenzyme F₄₂₀ biosynthesis. *Arch Microbiol* **180**: 455–464.
- Graupner M, White RH. (2001). Biosynthesis of the phosphodiester bond in coenzyme F₄₂₀ in the methanococci. *Biochemistry* **40**: 10859–10872.
- Graupner M, Xu H, White RH. (2002). Characterization of the 2-phospho-L-lactate transferase enzyme involved in coenzyme F₄₂₀ biosynthesis in *Methanococcus jannaschii*. *Biochemistry* **41**: 3033–3037.
- Greening C, Ahmed FH, Mohamed AE, Lee BM, Pandey G, Warden AC et al. (2016a). Physiology, biochemistry, and applications of F₄₂₀- and F_O-dependent redox reactions. *Microbiol Mol Biol Rev* **80**: 451–493.
- Greening C, Biswas A, Carere CR, Jackson CJ, Taylor MC, Stott MB et al. (2016b). Genome and metagenome surveys of hydrogenase diversity indicate H₂ is a widely-utilised energy source for microbial growth and survival. *ISME J* **10**: 761–777.
- Grochowski LL, Xu H, White RH. (2008). Identification and characterization of the 2-phospho-L-lactate guanylyl-transferase involved in coenzyme F₄₂₀ biosynthesis. *Biochemistry* **47**: 3033–3037.
- Guerra-Lopez D, Daniels L, Rawat M. (2007). *Mycobacterium smegmatis* mc² 155 *fbtC* and MSMEG_2392 are involved in triphenylmethane dye decolorization and coenzyme F₄₂₀ biosynthesis. *Microbiology* **153**: 2724–2732.
- Gurumurthy M, Rao M, Mukherjee T, Rao SPS, Boshoff HI, Dick T et al. (2013). A novel F₄₂₀-dependent anti-oxidant mechanism protects *Mycobacterium tuberculosis* against oxidative stress and bactericidal agents. *Mol Microbiol* **87**: 744–755.

- Hagemeyer CH, Shima S, Thauer RK, Bourenkov G, Bartunik HD, Ermler U. (2003). Coenzyme F₄₂₀-dependent methylenetetrahydromethanopterin dehydrogenase (Mtd) from *Methanopyrus kandleri*: a methanogenic enzyme with an unusual quaternary structure. *J Mol Biol* **332**: 1047–1057.
- Hedderich R, Whitman W. (2013). Physiology and biochemistry of the methane-producing Archaea. In: Rosenberg E, DeLong E, Lory S, Stackebrandt E, Thompson F (eds). *The Prokaryotes SE—81*. Springer: Berlin, Heidelberg, pp 635–662.
- Heiss G, Hofmann KW, Trachtmann N, Walters DM, Rouvière P, Knackmuss HJ. (2002). npd gene functions of *Rhodococcus (opacus) erythropolis* HL PM-1 in the initial steps of 2,4,6-trinitrophenol degradation. *Microbiology* **148**: 799–806.
- Houghton KM, Morgan XC, Lagutin K, MacKenzie AD, Vyssotski M, Mitchell KA et al. (2015). Thermorudis pharmacophila WKT50.2T sp. nov., a novel isolate of class Thermomicrobia isolated from geothermal soil, and emended descriptions of Thermomicrobium roseum, Thermomicrobium carboxidum, Thermorudis peleeae and Sphaerobacter thermophilus. *Int J Syst Evol Microbiol* **61**: 2482–2490.
- Isabelle D, Simpson DR, Daniels L. (2002). Large-scale production of coenzyme F₄₂₀-5,6 by using *Mycobacterium smegmatis*. *Appl Environ Microbiol* **68**: 5750–5755.
- Jackson TJ, Ramaley RF, Meinschein WG. (1973). *Thermomicrobium*, a new genus of extremely thermophilic bacteria. *Int J Syst Evol Microbiol* **23**: 28–36.
- Jacobson F, Walsh C. (1984). Properties of 7,8-didemethyl-8-hydroxy-5-deazaflavins relevant to redox coenzyme function in methanogen metabolism. *Biochemistry* **23**: 979–988.
- Janssen PH. (2006). Identifying the dominant soil bacterial taxa in libraries of 16S rRNA and 16S rRNA genes. *Appl Environ Microbiol* **72**: 1719–1728.
- Kann MG, Shoemaker BA, Panchenko AR, Przytycka TM. (2009). Correlated evolution of interacting proteins: looking behind the Mirrortree. *J Mol Biol* **385**: 91–98.
- Katoh K, Standley DM. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* **30**: 772–780.
- Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. (2015). The Phyre2 web portal for protein modeling, prediction and analysis. *Nat Protoc* **10**: 845–858.
- Kiener A, Husain I, Sancar A, Walsh C. (1989). Purification and properties of *Methanobacterium thermoautotrophicum* DNA photolyase. *J Biol Chem* **264**: 13880–13887.
- Knittel K, Lösekann T, Boetius A, Kort R, Amann R. (2005). Diversity and distribution of methanotrophic archaea at cold seeps. *Appl Environ Microbiol* **71**: 467–479.
- Lapalnik GV, Taylor MC, Warden AC, Scott C, Russell RJ, Oakeshott JG. (2012). F₄₂₀H₂-dependent degradation of aflatoxin and other furanocoumarins is widespread throughout the Actinomycetales. *PLoS One* **7**: e30114.
- Li H, Graupner M, Xu H, White RH. (2003). CofE catalyzes the addition of two glutamates to F₄₂₀-O in F₄₂₀ coenzyme biosynthesis in *Methanococcus jannaschii*. *Biochemistry* **42**: 9771–9778.
- Li W, Chou S, Khullar A, Gerratana B. (2009). Cloning and characterization of the biosynthetic gene cluster for Tomaymycin, an SJG-136 monomeric analog. *Appl Environ Microbiol* **75**: 2958–2963.
- Lin XL, White RH. (1986). Occurrence of coenzyme F₄₂₀ and its γ -monoglutamyl derivative in nonmethanogenic archaeobacteria. *J Bacteriol* **168**: 444–448.
- Markowitz VM, Chen I-MA, Palaniappan K, Chu K, Szeto E, Grechkin Y et al. (2012). IMG: the integrated microbial genomes database and comparative analysis system. *Nucleic Acids Res* **40**: D115–D122.
- Mees A, Klar T, Gnau P, Hennecke U, Eker APM, Carell T et al. (2004). Crystal structure of a photolyase bound to a CPD-like DNA lesion after in situ repair. *Science* **306**: 1789–1793.
- Meyer O, Schlegel HG. (1978). Reisolation of the carbon monoxide utilizing hydrogen bacterium *Pseudomonas carboxydovorans* (Kistner) comb. nov. *Arch Microbiol* **118**: 35–43.
- Michaelis W, Seifert R, Nauhaus K, Treude T, Thiel V, Blumenberg M et al. (2002). Microbial reefs in the black sea fueled by anaerobic oxidation of methane. *Science* **297**: 1013–1015.
- Muth E, Mörschel E, Klein A. (1987). Purification and characterization of an 8-hydroxy-5-deazaflavin-reducing hydrogenase from the archaeobacterium *Methanococcus voltae*. *Eur J Biochem* **169**: 571–577.
- Nelson-Sathi S, Sousa FL, Roettger M, Lozada-Chavez N, Thiergart T, Janssen A et al. (2015). Origins of major archaeal clades correspond to gene acquisitions from bacteria. *Nature* **517**: 77–80.
- Nocek B, Evdokimova E, Proudfoot M, Kudritska M, Grochowski LL, White RH et al. (2007). Structure of an amide bond forming F₄₂₀: γ -glutamyl ligase from *Archaeoglobus fulgidus*—a member of a new family of non-ribosomal peptide synthases. *J Mol Biol* **372**: 456–469.
- Overmars L, Kerkhoven R, Siezen RJ, Francke C. (2013). MGCV: the microbial genomic context viewer for comparative genome analysis. *BMC Genomics* **14**: 209.
- Petersen JL, Ronan PJ. (2010). Critical role of 7,8-didemethyl-8-hydroxy-5-deazariboflavin for photoreactivation in *Chlamydomonas reinhardtii*. *J Biol Chem* **285**: 32467–32475.
- Philms D, Decamps L, Berteau O, Begley TP. (2015). Biosynthetic versatility and coordinated action of 5'-deoxyadenosyl radicals in deazaflavin biosynthesis. *J Am Chem Soc* **137**: 5406–5413.
- Price MN, Dehal PS, Arkin AP. (2010). FastTree 2 – approximately maximum-likelihood trees for large alignments. *PLoS One* **5**: e9490.
- Pruitt KD, Tatusova T, Maglott DR. (2007). NCBI reference sequences (RefSeq): a curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids Res* **35**: D61–D65.
- Purwantini E, Daniels L. (1998). Molecular analysis of the gene encoding F₄₂₀-dependent glucose-6-phosphate dehydrogenase from *Mycobacterium smegmatis*. *J Bacteriol* **180**: 2212–2219.
- Purwantini E, Mukhopadhyay B. (2013). Rv0132c of *Mycobacterium tuberculosis* encodes a coenzyme F₄₂₀-dependent hydroxymycolic acid dehydrogenase. *PLoS One* **8**: e81985.
- Ray A, Lindahl E, Wallner B. (2012). Improved model quality assessment using ProQ2. *BMC Bioinformatics* **13**: 1–12.
- Shen J, Zhang J, Zuo Y, Wang L, Sun X, Li J et al. (2009). Biodegradation of 2,4,6-trinitrophenol by *Rhodococcus* sp. isolated from a picric acid-contaminated soil. *J Hazard Mater* **163**: 1199–1206.

- Shima S, Warkentin E, Grabarse W, Sordel M, Wicke M, Thauer RK *et al.* (2000). Structure of coenzyme F₄₂₀ dependent methylenetetrahydromethanopterin reductase from two methanogenic archaea. *J Mol Biol* **300**: 935–950.
- Snapper SB, Melton RE, Mustafa S, Kieser T, Jacobs WRJ. (1990). Isolation and characterization of efficient plasmid transformation mutants of *Mycobacterium smegmatis*. *Mol Microbiol* **4**: 1911–1919.
- Spang A, Poehlein A, Offre P, Zumbragel S, Haider S, Rychlik N *et al.* (2012). The genome of the ammonia-oxidizing Candidatus *Nitrososphaera gargensis*: insights into metabolic versatility and environmental adaptations. *Environ Microbiol* **14**: 3122–3145.
- Taylor MC, Jackson CJ, Tattersall DB, French N, Peat TS, Newman J *et al.* (2010). Identification and characterization of two families of F₄₂₀H₂-dependent reductases from *Mycobacteria* that catalyse aflatoxin degradation. *Mol Microbiol* **78**: 561–575.
- Thauer RK. (1998). Biochemistry of methanogenesis: a tribute to Marjory Stephenson. *Microbiology* **144**: 2377–2406.
- Walsh C. (1986). Naturally occurring 5-deazaflavin coenzymes: biological redox roles. *Acc Chem Res* **19**: 216–221.
- Wang P, Bashiri G, Gao X, Sawaya MR, Tang Y. (2013). Uncovering the enzymes that catalyze the final steps in oxytetracycline biosynthesis. *J Am Chem Soc* **135**: 7138–7141.
- Wilson MC, Mori T, Ruckert C, Uria AR, Helf MJ, Takada K *et al.* (2014). An environmental bacterial taxon with a large and distinct metabolic repertoire. *Nature* **506**: 58–62.
- Yamazaki S, Tsai L. (1980). Purification and properties of 8-hydroxy-5-deazaflavin-dependent NADP⁺ reductase from *Methanococcus vanniellii*. *J Biol Chem* **255**: 6462–6465.
- Yasui A, Takao M, Oikawa A, Kiener A, Walsh CT, Eker AP. (1988). Cloning and characterization of a photolyase gene from the cyanobacterium *Anacystis nidulans*. *Nucleic Acids Res* **16**: 4447–4463.

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)