SHORT COMMUNICATION

Evidence for a role of viruses in the thermal sensitivity of coral photosymbionts

Rachel Ashley Levin^{1,2,3}, Christian Robert Voolstra⁴, Karen Dawn Weynberg⁵ and Madeleine Josephine Henriette van Oppen^{5,6}

¹Centre for Marine Bio-Innovation, The University of New South Wales (UNSW), Sydney, NSW, Australia; ²School of Biological, Earth and Environmental Sciences, The University of New South Wales, Sydney, NSW, Australia; ³Climate Change Cluster, University of Technology Sydney, Sydney, NSW, Australia; ⁴Red Sea Research Center, Division of Biological and Environmental Science and Engineering, 4700 King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia; ⁵Australian Institute of Marine Science, Townsville, QLD, Australia and ⁶School of BioSciences, The University of Melbourne, Parkville, VIC, Australia

***Symbiodinium*, the dinoflagellate photosymbiont of corals, is posited to become more susceptible to viral infections when heat-stressed. To investigate this hypothesis, we mined transcriptome data of a thermosensitive and a thermotolerant type C1 *Symbiodinium* population at ambient (27 °C) and elevated (32 °C) temperatures. We uncovered hundreds of transcripts from nucleocytoplasmic large double-stranded DNA viruses (NCLDV) and the genome of a novel positive-sense single-stranded RNA virus (+ssRNAV). In the transcriptome of the thermosensitive population only, +ssRNAV transcripts had remarkable expression levels in the top 0.03% of all transcripts at 27 °C, but at 32 °C, expression levels of +ssRNAV transcripts decreased, while expression levels of anti-viral transcripts increased. In both transcriptomes, expression of NCLDV transcripts increased at 32 °C, but thermal induction of NCLDV transcripts involved in DNA manipulation was restricted to the thermosensitive population. Our findings reveal that viruses infecting *Symbiodinium* are affected by heat stress and may contribute to *Symbiodinium* thermal sensitivity.**

The ISME Journal (2017) 11, 808–812; doi:10.1038/ismej.2016.154; published online 2 December 2016

Tropical reef-building corals form a multipartite symbiosis with the photosynthetic dinoflagellate *Symbiodinium* and a diverse microbial community that includes bacteria, archaea, fungi, protists and viruses; collectively termed the coral holobiont (Rohwer *et al.*, 2002). The *Symbiodinium*–coral symbiosis can be disrupted by heat stress, which results in the loss of *Symbiodinium* cells from coral tissues, i.e., coral bleaching (Hoegh-Guldberg, 1999). Heat stress has been alleged to promote lytic viral infections of *Symbiodinium*, as virus-like particles have been found in heat-stressed *Symbiodinium* from the temperate sea anemone *Anemonia viridis* (Wilson *et al.*, 2001) and from the corals *Pavona danai*, *Acropora formosa* and *Stylophora pistillata* (Wilson *et al.*, 2005; Davy *et al.*, 2006). Additionally, following heat shock at 31 °C, the expression of a protein with homology to a eukaryotic viral protein

increased >100-fold in a *Symbiodinium*-enriched fraction of *Stylophora pistillata* tissue (Weston *et al.*, 2012).

Using transcriptome data generated by Levin *et al.* (2016) for two *Symbiodinium* type C1 populations cultured at 27 °C and 32 °C ($n=4$), we explore the effect of heat stress on viruses associated with *Symbiodinium* and *Symbiodinium* anti-viral responses. The *Symbiodinium* populations were originally isolated from the coral *Acropora tenuis* at South Molle Island (SM) and Magnetic Island (MI) (Great Barrier Reef, Australia). The thermosensitive SM population was found to suffer physiological damage in culture and to bleach *in hospite* at 32 °C, whereas the thermotolerant MI population was unaffected (Howells *et al.*, 2012; Levin *et al.*, 2016).

As many viral RNAs are polyadenylated (Wilson *et al.*, 2000; Priet *et al.*, 2015), they were retained in the poly(A)+ purified *Symbiodinium* RNA samples used for RNA-Seq (Levin *et al.*, 2016). Viral transcripts in each *de novo* transcriptome were identified through a robust BLASTx bit score approach adapted from Boschetti *et al.* (2012), which calculates the difference between the highest viral and the highest non-viral bit score to determine if a transcript is from a virus

Correspondence: RA Levin, Centre for Marine Bio-Innovation, The University of New South Wales (UNSW), Sydney, NSW 2052, Australia.

E-mail: rachylevin@gmail.com

Received 9 March 2016; revised 4 September 2016; accepted 18 September 2016; published online 2 December 2016

or the host, followed by GC content, trinucleotide frequency and codon usage analyses (Supplementary Materials and Methods, Supplementary Figure 1 and Supplementary Datasets 1 and 2). *Symbiodinium* anti-viral transcripts were identified by searching the transcriptomes for transcripts encoding anti-viral gene types found in corals and for transcripts with anti-viral Gene Ontology (Supplementary Materials and Methods).

The thermosensitive SM and thermotolerant MI transcriptomes contain 306 and 238 viral transcripts (Supplementary Discussion), as well as 62 and 65 anti-viral transcripts, respectively (NCBI GEO accession: GSE77911). In both transcriptomes, viral transcripts show homology to genes from NCLDVs *Mimiviridae* and *Phycodnaviridae* and the major capsid protein (MCP) gene from the dinoflagellate-specific +ssRNAV *Alvernaviridae* (*Dinornavirus*), which is in agreement with previous sequencing and transmission electron microscopy findings (Wilson *et al.*, 2005; Correa *et al.*, 2013, 2016). The +ssRNAV transcripts in the thermosensitive SM transcriptome share 94% nucleotide (nt) identity (TR74740|c13_g1_i1, 5202 nt; TR74740|c13_g1_i2, 2154 nt). The +ssRNAV transcript in the thermotolerant MI transcriptome is much shorter (TR97578|c0_g1_i1, 475 nt) but shares 100% nt identity with TR74740|c13_g1_i1. Successful PCR amplification of the MCP genes from complementary DNA reverse-transcribed from RNA—but not from genomic DNA—of both *Symbiodinium* populations supports that they are from +ssRNAVs (Supplementary

Materials and Methods and Supplementary Figure 2). However, phylogenetic analysis of the translated MCP gene sequences revealed that they are highly divergent from the *Dinornavirus* MCP gene and previously identified partial-length *Dinornavirus*-like MCP genes (Supplementary Discussion, Supplementary Table 1 and Supplementary Figure 3).

Both +ssRNAV transcripts in the thermosensitive SM transcriptome contain a putative viral internal ribosomal entry site (Supplementary Figure 4), which is related to the internal ribosomal entry site of the +ssRNA cricket paralysis virus, directly upstream from the MCP gene. The longer +ssRNAV transcript, TR74740|c13_g1_i1, also encodes an unannotated open reading frame (ORF) determined to be a +ssRNAV RNA replicase (RNA-dependent RNA-polymerase) polyprotein based on protein structure modeling (Supplementary Figure 5). The RNA replicase gene precedes the internal ribosomal entry site and MCP gene, giving the full transcript a markedly similar arrangement to the *Dinornavirus* and cricket paralysis virus complete genomes (Wilson *et al.*, 2000; Nagasaki *et al.*, 2005). Thus, we conclude TR74740|c13_g1_i1 to be the RNA genome of a novel +ssRNAV, making this the first discovered genome of any virus infecting *Symbiodinium* (Figure 1a). Surprisingly, the conserved dinoflagellate spliced leader, which is present on > 95% of *Symbiodinium* mRNAs (Zhang *et al.*, 2013), is at the 5' end of TR74740|c13_g1_i1. Although, if the +ssRNAV is dinoflagellate-specific like *Dinornavirus*, incorporation of the dinoflagellate spliced

Figure 1 Genome of the novel +ssRNAV and its expression in *Symbiodinium* transcriptomes. **(a)** Genome model of the +ssRNAV infecting *Symbiodinium* (TR74740|c13_g1_i1, GenBank accession: KX538960) and its similarities to the RNA genomes of the dinoflagellate-specific *Dinornavirus* (*Heterocapsa circularisquama* RNA virus strain 34, NCBI accession: AB218608) and cricket paralysis virus (NCBI accession: AF218039). **(b)** TMM-normalized FPKM at 27 °C (averaged across replicates for each transcript on day -1, $n=8$) for the +ssRNAV transcripts (TR74740|c13_g1_i1, 5757 FPKM; TR74740|c13_g1_i2, 1996 FPKM; TR97578|c0_g1_i1, 1 FPKM), NCLDV transcripts and non-viral transcripts in the thermosensitive SM and thermotolerant MI transcriptomes. Black lines mark the mean FPKM +s.d. for each subset of transcripts.

leader in the viral RNA genome is likely a case of molecular mimicry, a well-documented viral strategy to evade host immune responses that detect foreign nucleic acids (Elde and Malik, 2009), or for efficient cap-dependent translation by host polysomes (Zeiner *et al.*, 2003).

Viral transcripts had similar expression levels to many non-viral transcripts at 27 °C, with the exception of the two +ssRNAV transcripts in the thermosensitive SM transcriptome (Figure 1b). Both +ssRNAV transcripts in the thermosensitive SM transcriptome maintained average expression levels >1300 fragments per kb of transcript per million

mapped reads (FPKM) on all sampling time points at 27 °C, whereas the +ssRNAV transcript in the thermotolerant MI transcriptome had an average expression level <2 FPKM on all sampling time points. The vastly dissimilar expression levels of +ssRNAV transcripts between the transcriptomes suggest that the thermosensitive SM *Symbiodinium* population was experiencing a severe viral infection.

Differential expression analysis (Supplementary Materials and Methods) confirmed that no viral or anti-viral transcripts were differentially expressed between experimental groups of either population on

Figure 2 Viral infections and anti-viral responses of *Symbiodinium* under heat stress. Differential expression of (a) thermosensitive SM transcriptome +ssRNAV transcripts, (b) thermosensitive SM transcriptome NCLDV transcripts, (c) thermosensitive SM transcriptome anti-viral transcripts, (d) thermotolerant MI transcriptome NCLDV transcripts and (e) thermotolerant MI transcriptome anti-viral transcripts at 32 °C. Only transcripts with ≥ 2 -fold expression between 27 °C and 32 °C treatments on at least one sampling time point were analyzed. Transcripts were considered to have no differential expression on a sampling time point where the false discovery rate (FDR) was > 0.001 . Upregulated transcripts at 32 °C are shown in yellow. Downregulated transcripts at 32 °C are shown in purple. The gray regions represent the preheating sampling time point on day -1 when all replicates ($n=8$) were still at 27 °C. On day 0, replicates were ramped to 32 °C ($n=4$) or maintained at 27 °C ($n=4$) for the duration of the study (Supplementary Materials and Methods). Differential expression and transcript annotation results are detailed in Supplementary Tables 2–9.

day - 1 (preheating; all samples acclimated at 27 °C) (Figures 2a–e). In both transcriptomes, upregulation of NCLDV transcripts was induced at 32 °C and increased from day 9 to day 13 (Figures 2b and d, and Supplementary Tables 2–9). However, the upregulated NCLDV transcripts in the thermosensitive SM transcriptome encoded for a greater diversity of genes (F-box and FNIP repeat-containing proteins, resolvase, transposase, ankyrin repeat protein) compared to those in the thermotolerant MI transcriptome (only F-box and FNIP repeat-containing proteins). Viral F-box and FNIP repeat-containing proteins and ankyrin repeat proteins have possible roles in degrading proteins through protein–protein interactions, countering host defences and exploiting the host's ubiquitin-proteasome system to create an appropriate cellular environment for viral replication (Suhre, 2005; Sonnberg *et al.*, 2008; Fischer *et al.*, 2010), whereas resolvases and transposases directly participate in DNA manipulation (Iyer *et al.*, 2001; Schroeder *et al.*, 2009).

In the thermosensitive SM transcriptome, heat stress resulted in downregulation of the highly expressed +ssRNAV transcripts and upregulation of anti-viral transcripts (Figures 2a and c and Supplementary Tables 2–3 and 6–7). Conversely, the thermotolerant MI transcriptome showed no differential expression of its lowly expressed +ssRNAV transcript and downregulation of anti-viral transcripts at 32 °C (Figure 2e, Supplementary Tables 8–9). *Symbiodinium* anti-viral responses may therefore become activated from increased DNA manipulation by NCLDVs or initial upregulation of highly expressed +ssRNAV transcripts at 32 °C that was mitigated by the sampling time points on days 9 and 13.

Our study exemplifies how RNA-Seq data can be used to gain valuable insight into resident viruses. Our results indicate that only the thermosensitive SM *Symbiodinium* population experienced an extreme +ssRNAV infection and thermally induced NCLDV DNA manipulation. Thus, viral infections may factor into *Symbiodinium* thermal sensitivity, and consequently, coral bleaching.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

The Australian Institute of Marine Science supplied the *Symbiodinium* strains (aims-aten-C1-WSY, aims-aten-C1-MI) used in this study. Yi Jin Liew provided assistance with the bit score analysis. Rhys T Graham provided technical support for MATLAB. The Centre for Marine Bio-Innovation at the University of New South Wales, King Abdullah University of Science and Technology (KAUST), The Joyce W Vickery Scientific Research Fund Grant awarded to Rachel A Levin from the Linnean Society of New South Wales, and Future Fellowship No

FT100100088 awarded to Madeleine JH van Oppen from the Australian Research Council contributed financial support. Raw, processed and annotated sequencing data are available through NCBI GEO (accession: GSE77911). The novel +ssRNAV genome (TR74740|c13_g1_i1) and highly related, partial +ssRNAV genome (TR74740|c13_g1_i2) discovered in this study have been deposited in GenBank (accession nos: KX538960 and KX787934, poly(A) tails were trimmed from RNA-Seq reads prior to *de novo* assembly of transcripts).

References

- Boschetti C, Carr A, Crisp A, Eyres I, Wang-Koh Y, Lubzens E *et al.* (2012). Biochemical diversification through foreign gene expression in Bdelloid Rotifers. *PLoS Genet* **8**: e1003035.
- Correa AM, Ainsworth TD, Rosales SM, Thurber AR, Butler CR, Thurber RLV. (2016). Viral outbreak in corals associated with an *in situ* bleaching event: atypical herpes-like viruses and a new megavirus infecting *Symbiodinium*. *Front Microbiol* **7**: 127.
- Correa AM, Welsh RM, Thurber RLV. (2013). Unique nucleocytoplasmic dsDNA and +ssRNA viruses are associated with the dinoflagellate endosymbionts of corals. *ISME J* **7**: 13–27.
- Davy S, Burchett S, Dale A, Davies P, Davy J, Muncke C *et al.* (2006). Viruses: agents of coral disease? *Dis Aquat Organ* **69**: 101–110.
- Elde NC, Malik HS. (2009). The evolutionary conundrum of pathogen mimicry. *Nat Rev Microbiol* **7**: 787–797.
- Fischer MG, Allen MJ, Wilson WH, Suttle CA. (2010). Giant virus with a remarkable complement of genes infects marine zooplankton. *Proc Natl Acad Sci USA* **107**: 19508–19513.
- Hoegh-Guldberg O. (1999). Climate change, coral bleaching and the future of the world's coral reefs. *Mar Freshwater Res* **50**: 839–866.
- Howells EJ, Beltran VH, Larsen NW, Bay LK, Willis BL, van Oppen MJH. (2012). Coral thermal tolerance shaped by local adaptation of photosymbionts. *Nat Clim Change* **2**: 116–120.
- Iyer LM, Aravind L, Koonin EV. (2001). Common origin of four diverse families of large eukaryotic DNA viruses. *J Virol* **75**: 11720–11734.
- Levin RA, Beltran VH, Hill R, Kjelleberg S, McDougald D, Steinberg PD *et al.* (2016). Sex, scavengers, and chaperones: transcriptome secrets of divergent *Symbiodinium* thermal tolerances. *Mol Biol Evol* **33**: 2201–2215.
- Nagasaki K, Shirai Y, Takao Y, Mizumoto H, Nishida K, Tomaru Y. (2005). Comparison of genome sequences of single-stranded RNA viruses infecting the bivalve-killing dinoflagellate *Heterocapsa circularisquama*. *Appl Environ Microbiol* **71**: 8888–8894.
- Priet S, Lartigue A, Debart F, Claverie J-M, Abergel C. (2015). mRNA maturation in giant viruses: variation on a theme. *Nucleic Acids Res* **7**: 3776–3788.
- Rohwer F, Seguritan V, Azam F, Knowlton N. (2002). Diversity and distribution of coral-associated bacteria. *Mar Ecol Prog Ser* **243**: 1–10.
- Schroeder DC, Park Y, Yoon H-M, Lee YS, Kang SW, Meints RH *et al.* (2009). Genomic analysis of the

- smallest giant virus—*Feldmannia* sp. virus 158. *Virology* **384**: 223–232.
- Sonnberg S, Seet BT, Pawson T, Fleming SB, Mercer AA. (2008). Poxvirus ankyrin repeat proteins are a unique class of F-box proteins that associate with cellular SCF1 ubiquitin ligase complexes. *Proc Natl Acad Sci USA* **105**: 10955–10960.
- Suhre K. (2005). Gene and genome duplication in *Acanthamoeba polyphaga* Mimivirus. *J Virol* **79**: 14095–14101.
- Weston AJ, Dunlap WC, Shick JM, Klueter A, Iglic K, Vukelic A et al. (2012). A profile of an endosymbiont-enriched fraction of the coral *Stylophora pistillata* reveals proteins relevant to microbial–host interactions. *Mol Cell Proteomics* **11**: M111.015487.
- Wilson JE, Powell MJ, Hoover SE, Sarnow P. (2000). Naturally occurring dicistronic cricket paralysis virus RNA is regulated by two internal ribosome entry sites. *Mol Cell Biol* **20**: 4990–4999.
- Wilson W, Dale A, Davy J, Davy S. (2005). An enemy within? Observations of virus-like particles in reef corals. *Coral Reefs* **24**: 145–148.
- Wilson WH, Francis I, Ryan K, Davy SK. (2001). Temperature induction of viruses in symbiotic dinoflagellates. *Aquat Microb Ecol* **25**: 99–102.
- Zeiner GM, Sturm NR, Campbell DA. (2003). The *Leishmania tarentolae* spliced leader contains determinants for association with polysomes. *J Biol Chem* **278**: 38269–38275.
- Zhang H, Zhuang Y, Gill J, Lin S. (2013). Proof that dinoflagellate spliced leader (DinoSL) is a useful hook for fishing dinoflagellate transcripts from mixed microbial samples: *Symbiodinium kawagutii* as a case study. *Protist* **164**: 510–527.

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)