ORIGINAL ARTICLE

Metagenomic analysis of rapid gravity sand filter microbial communities suggests novel physiology of *Nitrospira* spp.

Alejandro Palomo^{1,2}, S Jane Fowler¹, Arda Gülay¹, Simon Rasmussen²,
Thomas Sicheritz-Ponten² and Barth F Smets¹

¹Department of Environmental Engineering, Technical University of Denmark, Kongens Lyngby, Denmark and

²Department of Systems Biology, Center for Biological Sequence Analysis, Technical University of Denmark, Kongens Lyngby, Denmark

Rapid gravity sand filtration is a drinking water production technology widely used around the world. Microbially catalyzed processes dominate the oxidative transformation of ammonia, reduced manganese and iron, methane and hydrogen sulfide, which may all be present at millimolar concentrations when groundwater is the source water. In this study, six metagenomes from various locations within a groundwater-fed rapid sand filter (RSF) were analyzed. The community gene catalog contained most genes of the nitrogen cycle, with particular abundance in genes of the nitrification pathway. Genes involved in different carbon fixation pathways were also abundant, with the reverse tricarboxylic acid cycle pathway most abundant, consistent with an observed *Nitrospira* dominance. From the metagenomic data set, 14 near-complete genomes were reconstructed and functionally characterized. On the basis of their genetic content, a metabolic and geochemical model was proposed. The organisms represented by draft genomes had the capability to oxidize ammonium, nitrite, hydrogen sulfide, methane, potentially iron and manganese as well as to assimilate organic compounds. A composite *Nitrospira* genome was recovered, and *amo*-containing *Nitrospira* genome contigs were identified. This finding, together with the high *Nitrospira* abundance, and the abundance of atypical *amo* and *hao* genes, suggests the potential for complete ammonium oxidation by *Nitrospira*, and a major role of *Nitrospira* in the investigated RSFs and potentially other nitrifying environments.

The ISME Journal (2016) 10, 2569–2581; doi:10.1038/ismej.2016.63; published online 29 April 2016

Introduction

Microorganisms have an essential role in Earth's global biogeochemical cycles, and they provide a multitude of nature's ecosystem services. Microbes also dominate the technical systems used in wastewater treatment, drinking-water production, waste management and bioenergy generation (Rittmann, 2006). Although these technical systems have been in use for many years, the study of their microbial composition has only recently been initiated. Unraveling the microbial composition and processes occurring in these systems will not only help optimize the technical processes but can also lead to the discovery of novel pathways and organisms. In the era of next-generation sequencing techniques, metagenomics-based approaches have enabled the

discovery of a suite of novel microbial pathways (Schleper *et al.*, 2005; Bryant *et al.*, 2007; Ettwig *et al.*, 2010; Carrión *et al.*, 2015) and the expansion of the tree of life (Brown *et al.*, 2015). Some of these discoveries have been made in technical systems; for example, novel candidate phyla and expanded metabolic versatility have been identified in aerobic and anaerobic bioreactors (Wexler *et al.*, 2005; Rinke *et al.*, 2013; Nobu *et al.*, 2015).

Rapid sand filters (RSFs), are engineered microbial systems used throughout the world for the production of drinking water from both surface and groundwater. With sufficiently high groundwater quality, rapid sand filtration—often in two serial filters—can be the sole treatment of groundwater before its distribution. Inlet water is initially aerated, which aims to physically remove compounds such as sulfide and methane and oxygenate the water. The main electron donors entering the serial filters are ammonia, reduced manganese and iron, small amounts of assimilable organic carbon, and residual methane and hydrogen sulfide (Tatari *et al.*, 2013; Albers *et al.*, 2015). Most of these compounds are oxidatively

Correspondence: BF Smets, Department of Environmental Engineering, Technical University of Denmark, Miljøvej 113, Kongens Lyngby 2800, Denmark.

E-mail: bfsm@env.dtu.dk

Received 23 November 2015; revised 29 February 2016; accepted 19 March 2016; published online 29 April 2016

transformed by microbes in the RSFs (Lee *et al.*, 2014). Hence, microorganisms colonize the filter material and its mineral coatings (Gülay *et al.*, 2014). In contrast to other engineered microbial systems, such as wastewater treatment plants and anaerobic digesters, RSFs treating groundwater can be considered oligotrophic (with influent concentrations of individual constituents below mM) and receive predominantly inorganic electron donors, supporting chemolithotrophic metabolisms (Gülay *et al.*, 2016).

Central metabolic functions in RSFs have previously been assigned to taxa based on their activity in other environments—*Nitrosomonas*, *Nitrospira*, *Gallionella* and *Bacillus* clades have been proposed as important ammonium (van der Wielen *et al.*, 2009; Lautenschlager *et al.*, 2014), nitrite (Gülay *et al.*, 2014; Albers *et al.*, 2015; LaPara *et al.*, 2015), iron (Søgaard, 2001; de Vet *et al.*, 2012; Li *et al.*, 2013) and manganese oxidizers (Mouchet, 1992; Cerrato *et al.*, 2010), respectively. Ammonia-oxidizing archaea (AOA) have been detected in RSFs at low abundance (van der Wielen *et al.*, 2009; Albers *et al.*, 2015), although in a few examples they outnumbered ammonia-oxidizing bacteria (AOB; Wang *et al.*, 2014; Nitzsche *et al.*, 2015). However, the total AOB and AOA densities are often insufficient to explain the high abundances of *Nitrospira* relative to the ammonium oxidizers (reviewed in Gülay *et al.*, 2016). The role of other dominant clades within RSF microbial communities (for example, *Acidobacteria*, *Actinobacteria*, *OD1* and *Chloroflexi*; Pinto *et al.*, 2012) are, similarly, undescribed.

Our recent investigation into the microbial communities of several groundwater-fed RSFs (Gülay *et al.*, 2016) revealed an astonishing and consistent abundance and microdiversity of *Nitrospira* spp. over all other taxa: up to 27% of all amplicons in pre-filters and up to 45% in after-filters were unambiguously identified as *Nitrospira* of lineages I, II and IV, although most phylotypes belonged to previously undescribed lineages. In addition, the *Rhizobiales*, *Chloracidobacterium*, *Da023*, *Acidimicrobiales*, *Gemmatimonadales* and *Burkholderiales* taxa were abundant, suggesting a central, yet to be identified, metabolic role for those taxa. Here, we apply shotgun metagenomic sequencing on replicate samples from an after-filter of one of the earlier described waterworks, where *Nitrospira* accounted for 65% of the total amplicon abundance (Gülay *et al.*, 2016), to determine the putative metabolic role of the *Nitrospira* clade and to elucidate the potential functions of the dominant taxa in investigated RSFs. We focused on two primary questions: can *Nitrospira* use any other energy source beyond nitrite in the RSFs? And which taxa contribute the central functions in the RSF? The metagenomic data set was assembled and a microbial gene catalog was constructed describing the community metabolic potential. By reassembly of taxonomically clustered contigs, we were able to identify and obtain 14 near-complete draft genomes for members of several abundant taxa

described in Gülay *et al.* (2016). A highly abundant composite *Nitrospira* genome was identified, which harbored genes for complete ammonium oxidation.

Materials and methods

Sample collection and extraction of DNA

The studied waterworks follows a treatment chain consisting of aeration and two sequential filtration steps. The aerated water flows downward through a first RSF, designed to remove ferrous iron, and then through a second RSF. DNA samples obtained from the study of Gülay *et al.* (2016) were subjected to shotgun metagenomic sequencing. Briefly, filter material samples were taken from Islevbro waterworks in Denmark and collected using a 60-cm-long core sampler 6 days after filter backwash. Core samples were obtained at three random positions in the filter to obtain biological replicates. Samples were extruded and sliced into five sections aseptically (0–5, 5–15, 15–25, 25–35 and 35–45 cm). DNA extracts from the top section (0–5 cm, ISLTop) were used to represent the uppermost microbial community. For each replicate, a composite DNA sample was generated by mixing equal volumes of DNA extracts of the other core sections (5–45 cm, ISLBulk) to represent the microbial community deeper in the filter. Thus, three samples from the top and three from the bulk of the filter were obtained.

Library preparation, sequencing and de novo assembly

DNA-shearing and library preparation were performed according to the NEXTflex Rapid DNA-Seq Kit, V13.08 (Bioo Scientific, Austin, TX, USA). Briefly, 250 ng genomic DNA was sheared with the Covaris E210 System using 10% duty cycle, intensity of 5, cycles per burst of 200 for 300 s to create 200-bp fragments. The samples were end-repaired and adenylated to produce an A-overhang. Adapters containing unique barcodes were ligated to the DNA. The samples were purified using bead-size selection for range ~300–400 bp with the Agencourt AMPure XP beads (Beckman Coulter, Beverly, MA, USA). The purified DNA libraries were amplified according to the manufacturer's protocol: initial denaturation (2 min, 98 °C), followed by 12 cycles of denaturation (98 °C, 30 s), annealing (65 °C, 30 s), extension (72 °C, 1 min) and final extension (72 °C, 4 min). DNA was quantified using NanoDrop ND-1000 UV-VIS Spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA), and quality was checked on an Agilent 2100 Bioanalyzer using the High Sensitivity DNA kit (Agilent Technologies, Santa Clara, CA, USA). The DNA libraries were mixed in equimolar ratios. Sequencing was performed as a 100-bp pair-end run on HiSeq 2000 (Illumina Int., San Diego, CA, USA) at BGI (Copenhagen, Denmark). *Trimmomatic* v0.22 (Bolger *et al.*, 2014) was used to remove adapters and trim the reads (threshold quality = 15; minimum

length=45). Quality control was carried out using *FastQC* (Babraham Bioinformatics (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>)). High-quality reads from each sample were assembled into scaffolds using *IDBA-UD* (Peng *et al.*, 2012) with default parameters. High-quality reads of each sample have been submitted to MG-RAST using the 'join fastq-formatted paired-reads' option retaining the nonoverlapping reads. They can be found under accession numbers 4629971.3, 4622590.3, 4629689.3, 4631157.3, 4630162.3 and 4631739.3.

Construction of a non-redundant gene catalog and quantification of reference gene abundance

Gene calling on the assembled scaffolds was performed using the metagenome implementation ('-p meta') of *Prodigal* 2.50 (Hyatt *et al.*, 2010). Predicted genes from all samples (2.85 M in total) were clustered using *UCLUST* (Edgar, 2010). Any two genes with greater than 90% nucleotide identity were clustered together, resulting in a set of 1 242 515 non-redundant genes. High-quality reads were mapped to the reference gene catalog using Burrows–Wheeler Aligner (Li and Durbin, 2010), *BWA-MEM*. Mapped reads were subsequently filtered by removing those with a poor alignment ($-q30$). The remaining mapped reads were used to form an abundance matrix of the number of reads mapped to each gene in every sample. When both read pairs mapped to the same gene, it was counted as a single hit. If each pair mapped to different genes then this is counted as a hit to each gene. The abundance matrix was normalized based on data set size and gene length.

Taxonomic and functional annotation of the gene catalog

Predicted coding sequences were annotated using *USEARCH* (Edgar, 2010) *-ublast* against the UniProt database (best hit with $E < 1e^{-5}$, Bitscore > 60 and sequence similarity $> 30\%$). An amino-acid similarity of 65% or higher was used as a threshold for phylum-level assignment (Li *et al.*, 2014). Clusters of orthologous groups (COG) and non-supervised orthologous groups (NOG) annotations of unassembled sequences were conducted in MG-RAST ($E < 1e^{-5}$, sequence similarity $> 30\%$; Meyer *et al.*, 2008) and compared with 25 metagenomes randomly selected from MG-RAST from five pre-selected biomes (Supplementary Table S1).

Binning, composite genome assembly and annotation

For each sample, contigs larger than 1000 nucleotides were clustered into putative taxonomic groups based on pentanucleotide signatures using VizBin (Laczny *et al.*, 2015). Contigs of the resultant clusters were used to recruit sequence reads from the original quality-filtered data set. Recruited reads in each cluster were *de novo* assembled as described above to construct composite genomes (CGs). Resultant

reconstructed CGs were manually evaluated through contig depth and GC content to validate accurate bin segregation. Completeness and potential contamination of each CG was evaluated using 107 essential single-copy genes (Albertsen *et al.*, 2013). When the same draft CG could be reconstructed from several samples the one best assembled, most complete, and with lowest contamination was retained. To improve assembly of higher-abundance organisms, taxonomic clusters were subsampled by segregation based on read depth and were subjected to *de novo* assembly. The best assemblies were selected based on N50, completeness and taxonomic affiliation of single-copy genes (for further details see Supplementary Information). CGs that contained at least 75% of the single-copy genes were further analyzed. Genomes were identified to appropriate taxonomic levels based on two approaches: (i) A set of 107 essential single-copy genes were blasted (BLASTP, $E < 1e^{-5}$) against the NCBI-nr database with follow-up analysis in MEGAN (Huson and Weber, 2013) to identify the lowest common ancestor of the blast output, (ii) using the standard settings of PhyloSift (Darling *et al.*, 2014). A taxon was assigned when at least 75% of the identified essential single-copy genes resulted in a concordant taxonomy. For genomic comparison between CGs and reference genomes, the average amino-acid identity (AAI; Konstantinidis and Tiedje, 2005) was calculated based on reciprocal best hits (two-way AAI) between two genomic data sets of proteins using the AAI calculator (<http://enve-omics.ce.gatech.edu/aa/>). The relative abundance of the organisms represented by the CGs was calculated by mapping the original, quality-filtered reads from each sample against the CG sequence. Gene calling and annotation of CGs was conducted as described above. Furthermore, to confirm protein functional assignment, Kyoto encyclopedia of genes and genomes (KEGG) annotations of the predicted proteins in each CG were obtained using the WebMGA server (Wu *et al.*, 2011). Presence of glycoside hydrolases and extracellular peptidases was evaluated by dbCAN ($E < 1e^{-5}$ and cover fraction > 0.4 ; Yin *et al.*, 2012) and MEROPS ($E < 1e^{-10}$; Rawlings *et al.*, 2014), respectively.

Phylogenetic analysis of ammonium oxidation genes

Predicted amino-acid sequences from the metagenomic samples and CGs were aligned with reference sequences using MUSCLE (Edgar, 2004) or T-Coffee (Notredame *et al.*, 2000). Multiple alignments were manually revised before phylogenetic analysis. Bootstrapped maximum likelihood trees were constructed in MEGA6 using the Jones Taylor Thornton model with 100 replicates (Tamura *et al.*, 2013).

Results and discussion

Metagenomic assembly and taxonomy

We sequenced whole-community DNA for six samples from a RSF (three samples from the top 5 cm of the filter, ISLTop and three composite

samples representing filter depths from 5 to 45 cm, ISLBulk), generating an average of 2.6 ± 0.1 Gbp of high-quality, paired-end sequence data per sample. The estimated abundance-weighted average coverage (Rodriguez-R and Konstantinidis, 2014) of the top and bulk metagenomes were 62% and 82%, respectively (Supplementary Figure S1), indicating that the majority of the microbial community was captured in the study. Reads from each data set were *de novo* assembled, producing an average of 293 391 contigs (Table 1). Prediction of coding sequences resulted in an average of 475 155 putative genes per data set, which were combined across the six samples to a non-redundant gene catalog of 1 242 515 genes. The annotation of the non-redundant gene catalog yielded a total of 768 197 genes with function assigned (Table 1).

Within all of the communities, *Bacteria* were by far the most abundant domain (ISLBulk, 97.85 ± 0.23 ; ISLTop, 98.63 ± 0.02). *Archaea* represented less than 1% of the filter community in both top and bulk samples (ISLBulk, 0.74 ± 0.10 ; ISLTop, 0.43 ± 0.02), consistent with previous studies examining the microbial communities in rapid gravity filtrations filters (Bai *et al.*, 2013; Wang *et al.*, 2014) and with a previous pyrosequencing analysis of the same community (Gülay *et al.*, 2016). At the phylum level, *Proteobacteria* and *Nitrospirae* dominate bulk sample communities with a relative abundance of $18.93\% \pm 1.31\%$ and $9.22\% \pm 4.72\%$, respectively; in the top of the filter these phyla were also dominant, with *Nitrospirae* most abundant ($26.08\% \pm 0.94\%$), followed by *Proteobacteria* ($14.47\% \pm 1.07\%$; Figure 1). These results are in-line with other observations of very high *Nitrospira* abundances (13% to 50% of all community 16S rRNA amplicons or clones) in similar oligotrophic water treatment and distribution systems (Martiny *et al.*, 2005; White *et al.*, 2012; LaPara *et al.*, 2015; Gülay *et al.*, 2016).

ISLBulk samples had higher alpha diversity and evenness than ISLTop samples. The average Shannon index for ISLBulk was 6.02 ± 0.40 , whereas for ISLTop samples it was 4.37 ± 0.10 . On the other hand, the Pielou evenness was 0.71 ± 0.05 in ISLBulk and 0.52 ± 0.02 in ISLTop. Together, these results

indicate a stratified distribution in the filter environment: the top-few centimeters of the filter are dominated by a few phylotypes—potentially involved in the removal of most of the groundwater contaminants (Gülay *et al.*, 2016)—whereas the remainder of the filter contains a higher diversity of less dominant microbes.

The functional potential of the RSF communities

The functional potential of the RSF microbial communities was contrasted with 25 metagenomes from different natural and engineered biomes (Supplementary Table S1) by comparison of COG and NOG categories (Supplementary Figure S2). This analysis showed that the RSF microbial community has relatively high proportions of chaperones and genes involved in post-translational modification and protein turnover (category O) as well as coenzyme transport and metabolism (H). These two categories have been detected to be abundant in oligotrophic ecosystems in comparison with higher

Figure 1 Average relative occurrence (%) of the most abundant phyla in top 5 cm (ISLTop, outer circle) and lower regions of filter (ISLBulk, inner circle) based on best blast hits of predicted protein sequences from the gene catalog to the Uniprot database (65% amino-acid identity for phylum-level assignment).

Table 1 Metagenome characteristics and general features of the gene catalog from the RSF samples

Metagenome characteristics	Top1	Top2	Top3	Bulk1	Bulk2	Bulk3
Sequenced reads (100 bp, millions)	29.69	30.62	33.35	30.89	30.26	33.54
Reads after quality filtering (millions)	24.72	25.80	27.95	25.01	24.33	26.93
Contigs	267 037	267 411	295 824	285 059	319 719	325 297
Mean size	822	814	833	875	854	906
N50	1104	1102	1184	1157	1028	1198
Predicted CDS	459 432	419 534	470 333	459 469	508 071	534 092
Non-redundant CDS			1 242 515			
Assigned putative functions			768 197			
Average assigned putative function length			163 ± 120			
% Reads mapped to gene catalog	76.31	79.29	75.50	68.80	60.83	62.61

Abbreviations: CDS, coding sequences; RFS, rapid sand filter.

Non-redundant predicted CDS from different samples were merged into a single gene catalog.

nutrient availability ecosystems (Ortiz *et al.*, 2013). The abundance of chaperones may help bacteria to deal with low nutrient availability and have been related to environmental stress tolerance (Storz and Hengge, 2011). The high abundance of secondary metabolism genes (Q) together with the low number of genes associated with transcription (K) in the filter samples is consistent with another study in which these two categories differentiated oligotrophic from copiotrophic microorganisms (Lauro *et al.*, 2009).

The abundance of transposases in the RSF metagenomes was striking: they accounted for $1.18\% \pm 0.09\%$ and $1.49\% \pm 0.04\%$ of the total annotated genes in the ISLBulk and ISLTop, respectively, substantially higher than in a study of 2137 complete genomes where the average transposase abundance was 0.83% (Aziz *et al.*, 2010). The high number of transposases may be associated with the filter environment, as mobile genetic elements have been found with greater prevalence in surface-attached communities (Stewart, 2013; Madsen *et al.*, 2012). Thus, transposases could provide the means by which phylogenetically distinct bacteria share genetic material, facilitating functional similarities as has been suggested in other studies (Hooper *et al.*, 2009).

Potential for inorganic electron-donor usage by the microbial community was inferred from the relative abundance of specific marker genes (Figure 2). Ammonium and nitrite oxidation were the dominant potential electron-donor processes in ISLTop. In ISLBulk potentials for the oxidation of ammonium, nitrite, reduced sulfur species, manganese and, to a lesser extent, methane, iron and hydrogen oxidation were more equal in abundance. This is consistent with experimental observations that nitrification is a dominant biogeochemical process in the first few centimeters of these filters (Tatari *et al.*, 2013; Lee *et al.*, 2014). Given the abundance of the ammonium oxidation potential, we investigated the phylogenetic diversity of *amoA* (as marker gene of ammonium oxidation) including reference *amoA* sequences and *amoA* genes recovered from the gene catalog (Figure 3). The RSF metagenome *amoA* sequences separated into four different clusters. In all, 1% and 7% (ISLBulk and ISLTop, respectively) of *amoA* genes were related to a cluster of oligotrophic AOB *Nitrosomonas* spp. including strain Is79A3 and *N. oligotropha*. Ammonia-oxidizing archaea-associated *amoA* made up 1% of the *amoA* in the ISLBulk samples; however, this cluster was not detected in the top of the filter. *amoA* associated with heterotrophic ammonia oxidizers made up 46% and 12% of all *amoA* (ISLBulk and ISLTop, respectively). Finally, the majority of the *amoA* genes (52% of *amoA* genes of the ISLBulk and 81% in ISLTop) could not be assigned to any known ammonia-oxidizing prokaryote (AOP) clades (40% dissimilar to known AOB at the protein level). Representative *pmoA* reference sequences were included to reject the possible misclassification of *amoA* sequences with the marker gene of methane oxidation, as they

Figure 2 Log-transformed relative abundances of marker genes for carbon fixation pathways, electron donors and electron acceptors in top 5 cm (ISLTop) and lower regions of filter (ISLBulk). Marker genes used for carbon fixation pathways were RuBisCo (CBB cycle), pyruvate: ferredoxin oxidoreductase (reverse TCA cycle) and 4-hydroxybutyryl-CoA dehydratase (4-hydroxybutyrate cycle).

share substantial sequence similarity (Holmes *et al.*, 1995). Consistent clustering patterns were observed for other ammonium oxidation-related genes: *amoB*, *amoC* and *hao* genes clustered with the characterized AOP clades, as well as an additional novel clade (Supplementary Figure S3).

In addition to potential electron-donor usage, we also investigated potential electron-acceptor usage (Figure 2). As expected from the predominantly aerobic conditions within the filters, cytochrome *c* oxidases involved in oxygen reduction accounted for around 80% of genes involved in electron-accepting processes. Genes involved in the reduction of nitrogen oxides (NO₃⁻ and NO₂⁻) were also relatively abundant in the gene catalog, showing the capacity of certain members of the community to use alternate electron acceptors under anoxic or micro-oxic conditions.

As the majority of characterized nitrifiers are autotrophic, we examined the diversity and relative abundance of genes for carbon fixation pathways in the gene catalog. The reverse tricarboxylic acid cycle (rTCA) pathway was dominant (Figure 2). The rTCA pathway is not found in typical AOB and *Alphaproteobacterial* nitrite-oxidizing bacteria (both

Figure 3 Phylogenetic reconstruction of *amoA* and *pmoA* reference protein sequences with putative *amoA* sequences recovered from the metagenomes (bold). Relative abundance of putative *amoA* clusters in the metagenomes is shown with colors corresponding to phylogenetic groups. Bootstrap support greater than 60 is indicated (based on 100 replicates).

have the Calvin–Benson–Bassham (CBB) cycle (Badger and Bek, 2007)), but is characteristic of the *Nitrospira* genus. In fact, all marker genes of the rTCA cycle found in the catalog were taxonomically assigned to *Nitrospira* spp. Carbon fixation via the CBB cycle was the second most abundant. The CBB genes displayed a wider phylogenetic diversity. The 4-hydroxybutyrate cycle pathway was also detected but was rare (Figure 2). Genes of the acetyl-CoA or 3-hydroxypropionate carbon fixation pathway were not

detected in the gene catalog. These results further underline the dominance of *Nitrospira* spp. among the autotrophic organisms in the examined filters.

Genome reconstruction and identification

Assembled contigs (length > 1000 bp) from each sample were used to generate taxonomically restricted bins using pentanucleotide frequency (Laczny *et al.*, 2015; Figure 4 and Supplementary Figure S4).

Genomic coverage and GC content were analyzed to validate bin segregation. Fourteen near-complete (>75% of essential genes) draft genomes were reconstructed from the metagenomic data sets (Table 2). Reconstructed genomes were classified as *Betaproteobacteria* (CG5, CG13 and CG26), *Alphaproteobacteria* (CG3, CG18 and CG6), *Acidobacteria* (CG10 and CG15), *Nitrospirae* (CG24), *Gammaproteobacteria* (CG7), *Gemmatimonadetes* (CG33) and *Planctomycetes* (CG4). Two of the recovered genomes could not be classified to currently defined phyla (CG1 and CG2). *Nitrospira* CG24 and *Acidobacteria* CG15 represent bins containing more than one genome based on the presence of multiple single-copy genes (Supplementary Table S2). The difficulty to separate these genomes probably resides in the existence of microdiversity, which is a hindrance to genome reconstruction and segregation from metagenomic data (Wilmes et al., 2009).

Composite genomes were genetically compared with their most closely related reference genomes (Supplementary Table S3). On the basis of average AAI, just two of the genomes had values approaching the species level of 85% (Luo et al., 2014). CG7 had an AAI of $78\% \pm 17\%$ with *Methyloglobulus morosus*, and CG26 had an AAI of $87\% \pm 15\%$ with *Nitrosomonas* sp. IS79. This analysis suggests that most of the draft genomes described herein are not closely related to previously sequenced organisms.

The most abundant CG both in ISLTop and ISLBulk samples was *Nitrospira* CG24, with an average abundance of 29.59% and 9.57%, respectively. CG10, affiliated to *Acidobacteria* subdivision 4, had a relative read proportion of 2.45% in the top of the filter and

2.43% in bulk samples. *Burkholderiales* CG13 and *Rhizobiales* CG3 appeared with greater than 1% abundance in both ISLTop and ISLBulk. *Acidobacteria* CG15 bin accounted for 5.37% of the reads in ISLBulk and the 0.70% in ISLTop. In addition, *Betaproteobacteria* CG5 and a poorly classified CG (CG1) were present with greater than 2% abundance in the bulk samples (Table 2).

Composite-genome metabolic potential

The metabolic potential of CGs was examined, focusing on energy and carbon metabolism, including electron donor, electron acceptor and carbon fixation pathways (Supplementary Figure S5).

Ammonium and nitrite oxidation

The near-complete CG26 contained a complete ammonia monooxygenase (*amoCAB*) operon. However, the gene set lacked *hao* (hydroxylamine oxidase), possibly because the genome was not complete. The *Nitrosomonas* bin encoded genes for the CBB cycle. Furthermore, it harbored an aa3-type cytochrome oxidase and nitrite reductase (*nirK*). The closest relative to CG26 was *Nitrosomonas* sp. IS79 (AAI of $87\% \pm 15\%$), a strain adapted to low ammonium concentrations (Bollmann et al., 2013).

CG24, a *Nitrospira* cluster, composed of five genomes (Supplementary Table S2), contained nitrite oxidoreductase (*nxr*) genes as well as cytochrome bd-type oxidase involved in oxygen reduction. This bin also encoded for nitrite reductase (NO-forming). CG24 harbored genes for the reductive citric acid cycle

Figure 4 Scatter plot of contigs (≥ 1 kb) assembled from metagenomic reads from sample Top2. Individual points represent contigs, with point size proportional to the natural log of contig length, and opacity proportional to the natural log of contig coverage. The manually placed red polygons highlight selected clusters used for the assembly of CGs as marked. A star shape highlights contigs annotated to contain the *amoA* (red) and *nxrB* (yellow) genes.

Table 2 Characteristics of composite genomes reconstructed from the RSF metagenomes

ID	Phylogenetic affiliation ^a	No. of contigs	Genetic size (Mb)	N50	GC (%)	Average sequencing depth	No. of genes	No. of single-copy genes ^b	No. of multiple single-copy genes	Relative abundance (%) TOP/BULK
CG7	<i>Methylococcaceae</i>	790	3.6	9864	48	14	3883	104/105	5	0.94/0.40
CG10	<i>Acidobacteria</i> subdivision 4	268	4.2	66 255	52	41	4015	104/106	4	2.45/2.43
CG5	<i>Betaproteobacteria</i>	1228	4.4	17 803	65	22	4938	104/106	13	0.45/2.13
CG1	<i>Bacteria</i>	897	5.6	25 218	69	18	5009	101/105	2	0.55/2.30
CG3	<i>Rhizobiales</i>	528	3.7	18 144	64	16	3754	99/105	3	1.14/1.44
CG33	<i>Gemmatimonadetes</i>	1567	4.5	8048	67	21	5141	96/104	8	0.30/1.91
CG2	<i>Bacteria</i>	868	2.8	4962	63	10	3062	91/105	3	0.14/0.57
CG13	<i>Burkholderiales</i>	540	5.1	19 632	68	15	5132	90/106	1	1.68/1.64
CG6	<i>Alphaproteobacteria</i>	871	2.0	3722	39	8	2502	88/105	4	0.21/0.06
CG4	<i>Planctomycetaceae</i>	1632	6.3	7253	62	10	6136	87/102	7	0.43/1.36
CG26	<i>Nitrosomonas</i>	1728	3.3	4142	43	10	4378	81/106	14	0.33/0.04
CG18	<i>Sphingomonas</i>	873	3.1	11 151	61	17	3666	80/105	6	0.91/0.53
CG24 ^c	<i>Nitrospira</i>	17 640	22.4	2136	56	70	33 880	105/105	102	29.59/9.57
CG15 ^c	<i>Acidobacteria</i>	2078	10.0	55 629	65	27	10 572	103/105	98	0.70/5.37

Abbreviation: RSF, rapid sand filter.

^aA taxon was assigned when at least 75% of the identified essential single-copy genes resulted in a concordant taxonomy.

^bThe number is given in relation to the average number of essential genes in all sequenced finished genomes in the given phylum (Albertsen *et al.*, 2013).

^cRepresent multiple genomes from the same clade.

(rTCA) involved in carbon fixation. In addition, it contained NAD⁺-dependent formate dehydrogenase and formate transporters, suggesting that it could oxidize formate coupled with the reduction of oxygen, nitrate or nitrite under anoxic conditions, as has been observed in other *Nitrospira* sp. (Koch *et al.*, 2015). Another feature recently found in *Nitrospira*, which was also encoded in this bin, is urease which is involved in the hydrolysis of urea into ammonia and carbamate (Koch *et al.*, 2015). CG24 contained urease subunits alpha, beta and gamma. Furthermore, it encoded chlorite dismutase, an enzyme involved in the conversion of chlorite to chloride and molecular oxygen. Surprisingly, *Nitrospira* CG24 also harbored three complete ammonia monooxygenase (*amoCAB*) operons. Considering the novelty of this finding, scaffolds were visualized for evenness of mapped read depth to identify potential chimeric regions using Integrative Genome Viewer (Thorvaldsdottir *et al.*, 2013). The visualization revealed homogeneity of read depth with other genes in the same scaffold and with other scaffolds in the same bin (Supplementary Figure S6). These were the most abundant *amo* genes in the gene catalog; however, they shared low identity with those previously described in classical AOB (and are therefore referred to as atypical-AMOX genes (Figures 3 and 5 and Supplementary Figure S3)). In comparison, typical AOB *amo* genes (here referred to as typical-AMOX genes) accounted for a low percentage of total *amo* genes. The same pattern was observed for other genes involved in ammonia oxidation (Supplementary Table S4). To rule out potential contamination during genome binning, depth and GC content of atypical-AMOX genes and typical-AMOX genes were compared with average depth and GC content of *Nitrospira* CG24 and *Nitrosomonas* CG26 (Supplementary Figure S7). In both cases, the distribution of atypical-AMOX genes

was identical to *Nitrospira* CG24 (Wilcoxon test, P -value > 0.05) but different from *Nitrosomonas* CG26. The comparison was also performed with other draft genomes, but none of them presented similar distributions (Supplementary Figure S7). To further support these findings, the covariance between the abundance of genes of interest (Supplementary Table S4) in the gene catalog and investigated bins (CG24 and CG26) along the six samples was analyzed. All atypical-AMOX genes statistically correlated with *Nitrospira* CG24 (R^2 > 0.97–0.99, P -value < 0.001) but not with *Nitrosomonas* CG26 (R^2 > 0.24–0.51, P -value > 0.1; Supplementary Figure S7). Together, these results strongly support the finding that *Nitrospira* CG24 contains genes encoding for ammonia oxidation.

Nitrospira CG24 was compared with two other *Nitrospira* genomes based on average AAI. CG24 has 70% AAI with *N. moscoviensis* and 65% with '*Candidatus Nitrospira defluvii*' (Supplementary Figure S8). These AAIs are far from the 85% species-level cutoff value but would comply with genus-level similarity (60–80%; Luo *et al.*, 2014). Thus, CG24 would represent a different species of the *Nitrospira* genus. An *amo* operon-containing contig from CG24 was compared with *amo* operons and flanking regions from *Nitrosomonas* sp. IS79 (the reference genome most closely related to the AOB found in this RSF) and an AOB reference genome from another genus (*Nitrosospira multiformis*). Predicted amino-acid sequence identities were highest for *amoC*, which shared 70% and 71% identity with *Nitrosomonas* and *Nitrosospira*, respectively. Identity was lower for *amoA* and *amoB*, ranging from 56% to 59%, respectively (Figure 5). This is in contrast to the *amo* AAI between *Nitrosomonas* and *Nitrosospira* spp., which are upwards of 80%. These results suggest that, whereas these proteins still share a common function, they

Figure 5 Comparison of *amo* operons and surrounding regions of *Nitrosomonas* sp. IS79A3 and *Nitrospira multiformis* with putative *amo* operon-containing contig belonging to CG24/*Nitrospira* sp. draft genome.

are derived from a more distantly related clade. Examination of the neighboring regions of the *amo* operon in CG24 revealed no similarity with neighboring regions in the *Nitrosomonas* and *Nitrospira* genomes (Figure 5).

Methane oxidation

A reconstructed genome CG7 contained particulate methane monooxygenase (*pmo*) and methanol dehydrogenase (*mdh*). This draft genome possessed the electron transport chain as well as aa3-type and bd-type cytochrome oxidases. In relation to carbon metabolism, it encoded for the TCA cycle, pentose phosphate pathway and Entner–Doudoroff pathway. On the basis of this genetic content, CG7 likely thrives on residual methane present in the water.

Oxidation of reduced sulfur, manganese and iron

A genomic bin (CG13) associated with *Burkholderiales* encoded dissimilatory sulfite reductases (*dsrA* and *dsrB*) for sulfide oxidation as well as adenylylsulfate reductases (*aprA* and *aprB*) and sulfate adenylyltransferase (*sat*) for complete oxidation of sulfite to sulfate.

CG13 carried genes for near-complete CBB cycle, including ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO) gene. CG13 contained cytochrome *cbb3*-type oxidase, periplasmic nitrate reductase (*napA*) and nitrite reductase (NO-forming). Furthermore, this genome contained *mtrAB* genes with homology (amino-acid sequence identity from 29% to 55%) to those found in lithotrophic iron-oxidizing bacteria (Jiao and Newman, 2007; Beckwith *et al.*, 2015). Thus, CG13 could be involved in the oxidation of ferrous iron, although the presence of *mtrABC* (periplasmic and outer membrane cytochromes) could also indicate electron transfer to minerals (Weber *et al.*, 2006; Gülay *et al.*, 2014). The CG13 bin also contained multicopper oxidases homologous (60–75% at the protein level) to those found in the known manganese oxidizers *Leptothrix cholodnii* str. SP-6 and *Pedomicrobium* sp. ACM 3067, suggesting a potential capacity for manganese oxidation.

Rhizobiales CG3 encoded for two multicopper oxidases with sequence identity (59% to 75% at protein level) related to those found in the manganese oxidizer *Aurantimonas manganoxydans* SI85-9A1 (Dick *et al.*, 2008). CG3 encoded for cytochrome

cbb3-type oxidase and nitrate reductase (*narG*). *Rhizobiales* CG3 also harbored genes of the TCA cycle, pentose phosphate pathway (non-oxidative phase) and beta oxidation pathway. Moreover, this genetic bin possessed extracellular peptidase and glycoside hydrolases involved in protein and carbohydrate metabolism, respectively (Supplementary Figure S9). These genetic features together with the absence of any carbon fixation pathway suggest the potential use of Mn^{2+} as an energy source, coupled to organic compound degradation as a carbon source. This chemolithoheterotrophic behavior has previously been observed in other manganese oxidizers (Francis *et al.*, 2001; Templeton *et al.*, 2005).

Heterotrophy/organic carbon degradation

To explore the capability of the microbial community to degrade organic carbon present in the influent water as well as products derived from microbial growth and decay, we screened reconstructed genomes for carbon metabolism pathways, glycoside hydrolases, extracellular peptidases and sugar and amino-acid transporters. Most of CGs encoded for complete or near-complete TCA cycle, beta oxidation and glycolysis pathways. Partial or complete pentose phosphate pathways were also present in most of the CGs. On the other hand, the Entner–Doudoroff pathway was less abundant, being complete, or mostly complete in only four of the CGs. In relation to glycoside hydrolases, most of the CGs contained genes for the degradation of cellulose, peptidoglycan, N-acetylglucosamine and starch (Supplementary Figure S9). Glycoside hydrolases were particularly abundant and diverse in both *Acidobacteria* CG15 and CG10 and *Planctomycetaceae* CG4. Reconstructed genomes CG1 and *Sphingomonas* CG18 had glycoside hydrolases for the degradation of hemicellulose. Furthermore, extracellular peptidases were present in most of CGs. CG1, *Gemmatimonadetes* CG33 and *Acidobacteria* CG10 possessed several genes of different families of extracellular peptidases potentially involved in the decomposition of bacterial cell walls and proteins (Supplementary Figures S8 and S9). These observations, together with the presence of sugar (CG10, CG2, CG1 and CG4) and amino-acid (CG15 and CG3) transporters in several of the reconstructed genomes, indicate the potential of several of the CGs to degrade and uptake organic carbon in this system. In particular, the high diversity of genes involved in carbohydrate degradation and uptake in CG1 and both *Acidobacteria* CG15 and CG10 combined with their high abundance suggests that these organisms may have a key role in organic carbon flow in the filter.

Predicted metabolic and geochemical model

On the basis of metabolic reconstruction of CGs and the chemical characteristics of the RSF, a metabolic and geochemical model of the RSF community is presented

Figure 6 Model of predicted metabolic and geochemical processes facilitating the degradation of groundwater contaminants in rapid gravity sand filters based on metagenomic analysis. Gray arrows denote metabolic capability, whereas blue arrows denote putative metabolic capability.

(Figure 6). Reconstructed genome *Methylococcaceae* CG7 could be involved in methane oxidation, as methane may not be completely stripped during aeration. *Burkholderiales* CG13 harbors the metabolic capabilities to realize complete oxidation of sulfide to sulfate and may be also associated with autotrophic manganese and iron oxidation. The ability to oxidize Fe^{2+} and Mn^{2+} indiscriminately has also been observed in other bacteria (Gouzinis *et al.*, 1998). Manganese could potentially also be oxidized by the heterotrophic *Rhizobiales* CG3. A large number of CGs encoded heterotrophic metabolisms, which are likely involved in the degradation of organic molecules present in the groundwater or the decay and metabolic by-products of autotrophic biomass. Ammonium in the filter could be oxidized by the typical autotrophic AOB *Nitrosomonas* CG26. The nitrite produced by CG26 could then be utilized by *Nitrospira* CG24. However, given its metabolic potential, the dominance of *Nitrospira* CG24 and the high abundance of *Nitrospira*-AMOX genes compared with other AMO genes, *Nitrospira* CG24 might mediate complete ammonium oxidation in this system. This would be in agreement with the suggestion that a microorganism that completely oxidizes ammonia to nitrate exists (Costa *et al.*, 2006). The *Nitrospira* abundance observed in this study is consistent with earlier pyrosequencing and quantitative PCR-based analysis of the same system, where abundances of up to 65% and 18%, respectively, were measured, and *Nitrospira* 16S rRNA gene sequences were often up to two orders of magnitude more abundant than *Nitrosospira* and *Nitrosomonas* 16S rRNA sequences combined (Gülay *et al.*, 2016). Further, ammonium oxidation *Nitrospira* activity may help to explain the high abundance of *Nitrospira* and the unusual *Nitrospira*/AOB ratios observed in other rapid gravity filters (Feng *et al.*, 2012; Albers *et al.*,

2015; Cai *et al.*, 2015; LaPara *et al.*, 2015). Whereas *Nitrospira* may be the primary ammonium oxidizer, the high abundance of heterotrophic *amoA* sequences, especially in the bulk samples, suggests that heterotrophic ammonium oxidation may also occur in the RSFs.

During the revision of this manuscript, evidence of complete ammonia oxidation (comammox) by *Nitrospira* organisms was described by others (Daims *et al.*, 2015; van Kessel *et al.*, 2015). Metagenomic evidence of comammox *Nitrospira* in a drinking water distribution system was reported by Pinto *et al.* (2016). Average AAI of CG24 was compared with the recently published comammox genomes: *Nitrospira* CG24 would represent a different species within the *Nitrospira* genus (AAI of $68\% \pm 16\%$ with '*Candidatus Nitrospira inopinata*', $71 \pm 16\%$ '*Candidatus Nitrospira nitrosa*', $73 \pm 17\%$ '*Candidatus Nitrospira nitrificans*' and $75 \pm 18\%$ with the *Nitrospira* sp. genome from a drinking-water system) sharing the metabolic potential for complete ammonia oxidation. As CG24 consisted of more than a single genome, we investigated the heterogeneity within CG24. We examined the average number of essential single-copy genes and *amo* operons (one in recently published comammox genomes; Daims *et al.*, 2015; Pinto *et al.*, 2016; van Kessel *et al.*, 2015) present in CG24. The *Nitrospira* bin CG24 has an average of 4.7 essential single-copy genes and three *amo* operons. On the basis of this, we predict that approximately three out of the five potential genomes represented by CG24 are complete ammonia oxidizers. Although we were unable to assign the comammox *Nitrospira* in this system to a specific lineage, previous analysis of this system revealed that the majority of *Nitrospira* sequences in this filter belong to novel lineages (Gülay *et al.*, 2016). Thus, we predict that the comammox *Nitrospira* present in this system may belong to a novel lineage, and not to lineage II as for other described comammox *Nitrospira*. Further sequencing of the comammox *Nitrospira* in this system is required to elucidate the lineage of the comammox *Nitrospira* described herein. In conclusion, our metagenomic survey provides insights into the metabolic capabilities of the microbial communities in an oligotrophic engineered system. Genome reconstructions allowed us to predict the roles of dominant community members in nitrogen, carbon, manganese, iron, methane and sulfur cycling—the main biological processes occurring in the sand filters. The results of our analysis point toward the novel metabolic capability of complete ammonia oxidation in the genus *Nitrospira*, which was highly abundant and ubiquitous in the investigated RSFs.

The metabolic and geochemical model of the RSF microbial communities can be used to inform further modeling efforts with the ultimate goal of enhancing contaminant removal and improving the design or function of RSFs. Improved understanding of the function of novel taxa in RSFs may be of considerable utility in restoring malfunctioning filters where common microbiological data have, thus far, been

insufficient to explain the observed patterns of contaminant removal. Transcriptomic analysis of the RSFs will help to validate the predictions made in this study and to elucidate the ecological importance of the various taxa and functions in RSFs.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

We thank K Amoako, MD Dalgaard, and BJR Cockx for supporting sample collection, library preparation and NG sequencing, and graphics. This research was financially supported by MERMAID (An initial training network funded by the People Programme - Marie Skłodowska-Curie Actions- of the European Union's Seventh Framework Programme FP7/2007-2013/ under REA grant agreement n°607492).

Author contributions

The original study design was conceived by AG, AP, TSP and BFS. Sampling and DNA extraction were performed by AG. AP performed bioinformatic analyses, supported by SJF, TSP and SR. AP and SJF lead interpretation of the results. AP drafted the manuscript; all authors contributed to its revision and completion, and approved the final submission.

References

- Albers CN, Ellegaard-Jensen L, Harder CB, Rosendahl S, Knudsen BE, Ekelund F *et al.* (2015). Groundwater chemistry determines the prokaryotic community structure of waterworks sand filters. *Environ Sci Technol* **49**: 839–846.
- Albertsen M, Hugenholtz P, Skarshewski A, Nielsen KL, Tyson GW, Nielsen PH. (2013). Genome sequences of rare, uncultured bacteria obtained by differential coverage binning of multiple metagenomes. *Nat Biotechnol* **31**: 533–538.
- Aziz RK, Breitbart M, Edwards RA. (2010). Transposases are the most abundant, most ubiquitous genes in nature. *Nucleic Acids Res* **38**: 4207–4217.
- Badger MR, Bek EJ. (2007). Multiple Rubisco forms in proteobacteria: their functional significance in relation to CO₂ acquisition by the CBB cycle. *J Exp Bot* **59**: 1525–1541.
- Bai Y, Liu R, Liang J, Qu J. (2013). Integrated metagenomic and physiochemical analyses to evaluate the potential role of microbes in the sand filter of a drinking water treatment system. *PLoS One* **8**: e61011.
- Beckwith CR, Edwards MJ, Lawes M, Shi L, Butt JN, Richardson DJ *et al.* (2015). Characterization of MtoD from *Sideroxydans lithotrophicus*: a cytochrome c electron shuttle used in lithoautotrophic growth. *Front Microbiol* **6**: 332.

- Bolger AM, Lohse M, Usadel B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.
- Bollmann A, Sedlacek CJ, Norton J, Laanbroek HJ, Suwa Y, Stein LY et al. (2013). Complete genome sequence of *Nitrosomonas* sp. Is79, an ammonia oxidizing bacterium adapted to low ammonium concentrations. *Stand Genomic Sci* **7**: 469–482.
- Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A et al. (2015). Unusual biology across a group comprising more than 15% of domain bacteria. *Nature* **523**: 208–211.
- Bryant DA, Costas AMG, Maresca JA, Chew AGM, Klatt CG, Bateson MM et al. (2007). Candidatus Chloracidobacterium thermophilum: an aerobic phototrophic Acidobacterium. *Science* **317**: 523–526.
- Cai Y, Li D, Liang Y, Luo Y, Zeng H, Zhang J. (2015). Effective start-up biofiltration method for Fe, Mn, and ammonia removal and bacterial community analysis. *Bioresour Technol* **176**: 149–155.
- Carrión O, Curson ARJ, Kumaresan D, Fu Y, Lang AS, Mercadé E et al. (2015). A novel pathway producing dimethylsulphide in bacteria is widespread in soil environments. *Nat Commun* **6**: 6579.
- Cerrato JM, Falkinham JO, Dietrich AM, Knocke WR, McKinney CW, Pruden A. (2010). Manganese-oxidizing and -reducing microorganisms isolated from biofilms in chlorinated drinking water systems. *Water Res* **44**: 3935–3945.
- Costa E, Pérez J, Kreft J-U. (2006). Why is metabolic labour divided in nitrification? *Trends Microbiol* **14**: 213–219.
- Daims H, Lebedeva EV, Pjevac P, Han P, Herbold C, Albertsen M et al. (2015). Complete nitrification by *Nitrospira* bacteria. *Nature* **528**: 504–509.
- Darling AE, Jospin G, Lowe E, Matsen FA, Bik HM, Eisen JA. (2014). PhyloSift: phylogenetic analysis of genomes and metagenomes. *PeerJ* **2**: e243.
- de Vet WWJM, Dinkla IJT, Abbas BA, Rietveld LC, van Loosdrecht MCM. (2012). *Gallionella* spp. in trickling filtration of subsurface aerated and natural groundwater. *Biotechnol Bioeng* **109**: 904–912.
- Dick GJ, Podell S, Johnson HA, Rivera-Espinoza Y, Bernier-Latmani R, McCarthy JK et al. (2008). Genomic insights into Mn(II) oxidation by the marine alphaproteobacterium *Aurantimonas* sp. strain SI85-9A1. *Appl Environ Microbiol* **74**: 2646–2658.
- Edgar RC. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**: 1792–1797.
- Edgar RC. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**: 2460–2461.
- Ettwig KF, Butler MK, Le Paslier D, Pelletier E, Mangenot S, Kuypers MMM et al. (2010). Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. *Nature* **464**: 543–548.
- Feng S, Xie S, Zhang X, Yang Z, Ding W, Liao X et al. (2012). Ammonium removal pathways and microbial community in GAC-sand dual media filter in drinking water treatment. *J Environ Sci (China)* **24**: 1587–1593.
- Francis CA, Co EM, Tebo BM. (2001). Enzymatic manganese(II) oxidation by a marine alpha-proteobacterium. *Society* **67**: 4024–4029.
- Gouzinis A, Kosmidis N, Vayenas DV, Lyberatos G. (1998). Removal of Mn and simultaneous removal of NH₃, Fe and Mn from potable water using a trickling filter. *Water Res* **32**: 2442–2450.
- Gülay A, Musovic S, Alberchtsen H-J, Al-Soud WA, Sørensen S, Smets BF. (2016). Ecological patterns, diversity and core taxa of microbial communities in groundwater-fed rapid gravity filters. *ISME J*; e-pub ahead of print 8 March 2016; doi:10.1038/ismej.2016.16.
- Gülay A, Tatari K, Musovic S, Mateiu RV, Albrechtsen H-J, Smets BF. (2014). Internal porosity of mineral coating supports microbial activity in rapid sand filters for groundwater treatment. *Appl Environ Microbiol* **80**: 7010–7020.
- Holmes AJ, Costello A, Lidstrom ME, Murrell JC. (1995). Evidence that particulate methane monooxygenase and ammonia monooxygenase may be evolutionarily related. *FEMS Microbiol Lett* **132**: 203–208.
- Hooper SD, Mavromatis K, Kyrpides NC. (2009). Microbial co-habitation and lateral gene transfer: what transposases can tell us. *Genome Biol* **10**: R45.
- Huson DH, Weber N. (2013). Microbial community analysis using MEGAN. *Methods Enzymol* **531**: 465–485.
- Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. (2010). Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**: 119.
- Jiao Y, Newman DK. (2007). The pio operon is essential for phototrophic Fe(II) oxidation in *Rhodospseudomonas palustris* TIE-1. *J Bacteriol* **189**: 1765–1773.
- Koch H, Lückner S, Albertsen M, Kitzinger K, Herbold C, Spieck E et al. (2015). Expanded metabolic versatility of ubiquitous nitrite-oxidizing bacteria from the genus *Nitrospira*. *Proc Natl Acad Sci USA* **112**: 11371–11376.
- Konstantinidis KT, Tiedje JM. (2005). Towards a genome-based taxonomy for prokaryotes. *J Bacteriol* **187**: 6258–6264.
- Laczny CC, Sternal T, Plugaru V, Gawron P, Atashpendar A, Margossian H et al. (2015). VizBin - an application for reference-independent visualization and human-augmented binning of metagenomic data. *Microbiome* **3**: 1.
- LaPara TM, Hope Wilkinson K, Strait JM, Hozalski RM, Sadowksy MJ, Hamilton MJ. (2015). The bacterial communities of full-scale biologically active, granular activated carbon filters are stable and diverse and potentially contain novel ammonia-oxidizing microorganisms. *Appl Environ Microbiol* **81**: 6864–6872.
- Lauro FM, McDougald D, Thomas T, Williams TJ, Egan S, Rice S et al. (2009). The genomic basis of trophic strategy in marine bacteria. *Proc Natl Acad Sci USA* **106**: 15527–15533.
- Lautenschlager K, Hwang C, Ling F, Liu W-T, Boon N, Köster O et al. (2014). Abundance and composition of indigenous bacterial communities in a multi-step biofiltration-based drinking water treatment plant. *Water Res* **62**: 40–52.
- Lee CO, Boe-Hansen R, Musovic S, Smets BF, Albrechtsen H-J, Binning P. (2014). Effects of dynamic operating conditions on nitrification in biological rapid sand filters for drinking water treatment. *Water Res* **64C**: 226–236.
- Li H, Durbin R. (2010). Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* **26**: 589–595.
- Li J, Jia H, Cai X, Zhong H, Feng Q, Sunagawa S et al. (2014). An integrated catalog of reference genes in the human gut microbiome. *Nat Biotechnol* **32**: 834–841.

- Li X, Chu Z, Liu Y, Zhu M, Yang L, Zhang J. (2013). Molecular characterization of microbial populations in full-scale biofilters treating iron, manganese and ammonia containing groundwater in Harbin, China. *Bioresour Technol* **147**: 234–239.
- Luo C, Rodriguez-R LM, Konstantinidis KT. (2014). MyTaxa: An advanced taxonomic classifier for genomic and metagenomic sequences. *Nucleic Acids Res* **42**: 1–12.
- Madsen JS, Burmølle M, Hansen LH, Sørensen SJ. (2012). The interconnection between biofilm formation and horizontal gene transfer. *FEMS Immunol Med Microbiol* **65**: 183–195.
- Martiny AC, Albrechtsen H, Arvin E, Molin S. (2005). Identification of bacteria in biofilm and bulk water samples from a nonchlorinated model drinking water distribution system: detection of a large nitrite-oxidizing population associated with *Nitrospira* spp. *Appl Environ Microbiol* **71**: 8611–8617.
- Meyer F, Paarmann D, D'Souza M, Olson R, Glass E, Kubal M et al. (2008). The metagenomics RAST server – a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics* **9**: 386.
- Mouchet P. (1992). From conventional to biological removal of iron and manganese in France. *J Am Water Work Assoc* **84**: 158–167.
- Nitzsche KS, Weigold P, Lösekann-Behrens T, Kappler A, Behrens S. (2015). Microbial community composition of a household sand filter used for arsenic, iron, and manganese removal from groundwater in Vietnam. *Chemosphere* **138**: 47–59.
- Nobu MK, Narihiro T, Rinke C, Kamagata Y, Tringe SG, Woyke T et al. (2015). Microbial dark matter ecogenomics reveals complex synergistic networks in a methanogenic bioreactor. *ISME J* **9**: 1710–1722.
- Notredame C, Higgins DG, Heringa J. (2000). T-Coffee: a novel method for fast and accurate multiple sequence alignment. *J Mol Biol* **302**: 205–217.
- Ortiz M, Legatzki A, Neilson JW, Fryslie B, Nelson WM, Wing RA et al. (2013). Making a living while starving in the dark: metagenomic insights into the energy dynamics of a carbonate cave. *ISME J* **8**: 478–491.
- Peng Y, Leung HCM, Yiu SM, Chin FYL. (2012). IDBA-UD: a de novo assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics* **28**: 1420–1428.
- Pinto AJ, Marcus DN, Ijaz Z, Bautista-de los Santos QM, Dick GJ, Raskin L. (2016). Metagenomic evidence for the presence of comammox *Nitrospira*-like bacteria in a drinking water system. *mSphere* **1**: e00054–15.
- Pinto AJ, Xi C, Raskin L. (2012). Bacterial community structure in the drinking water microbiome is governed by filtration processes. *Environ Sci Technol* **46**: 8851–8859.
- Rawlings ND, Waller M, Barrett AJ, Bateman A. (2014). MEROPS: the database of proteolytic enzymes, their substrates and inhibitors. *Nucleic Acids Res* **42**: D503–D509.
- Rinke C, Schwientek P, Sczyrba A, Ivanova NN, Anderson IJ, Cheng J-F et al. (2013). Insights into the phylogeny and coding potential of microbial dark matter. *Nature* **499**: 431–437.
- Rittmann BE. (2006). Microbial ecology to manage processes in environmental biotechnology. *Trends Biotechnol* **24**: 261–266.
- Rodriguez-R LM, Konstantinidis KT. (2014). Estimating coverage in metagenomic data sets and why it matters. *ISME J* **8**: 2349–2351.
- Schleper C, Jurgens G, Jonuscheit M. (2005). Genomic studies of uncultivated archaea. *Nat Rev Microbiol* **3**: 479–488.
- Søgaard E. (2001). Conditions for biological precipitation of iron by *Gallionella ferruginea* in a slightly polluted ground water. *Appl Geochemistry* **16**: 1129–1137.
- Stewart FJ. (2013). Where the genes flow. *Nat Geosci* **6**: 688–690.
- Storz G, Hengge R. (2011). *Bacterial Stress Responses*. ASM Press: Washington, DC, USA.
- Tamura K, Stecher G, Peterson D, Filipinski A, Kumar S. (2013). MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* **30**: 2725–2729.
- Tatari K, Smets BF, Albrechtsen H-J. (2013). A novel bench-scale column assay to investigate site-specific nitrification biokinetics in biological rapid sand filters. *Water Res* **47**: 6380–6387.
- Templeton AS, Staudigel H, Tebo BM. (2005). Diverse Mn (II)-oxidizing bacteria isolated from submarine basalts at Loihi Seamount. *Geomicrobiol J* **22**: 127–139.
- Thorvaldsdottir H, Robinson JT, Mesirov JP. (2013). Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform* **14**: 178–192.
- van der Wielen PWJJ, Voost S, van der Kooij D. (2009). Ammonia-oxidizing bacteria and archaea in groundwater treatment and drinking water distribution systems. *Appl Environ Microbiol* **75**: 4687–4695.
- van Kessel MAHJ, Speth DR, Albertsen M, Nielsen PH, Op den Camp HJM, Kartal B et al. (2015). Complete nitrification by a single microorganism. *Nature* **528**: 555–559.
- Wang H, Narihiro T, Straub AP, Pugh CR, Tamaki H, Moor JF et al. (2014). MS2 bacteriophage reduction and microbial communities in biosand filters. *Environ Sci Technol* **48**: 6702–6709.
- Weber KA, Achenbach LA, Coates JD. (2006). Microorganisms pumping iron: anaerobic microbial iron oxidation and reduction. *Nat Rev Microbiol* **4**: 752–764.
- Wexler M, Bond PL, Richardson DJ, Johnston AWB. (2005). A wide host-range metagenomic library from a waste water treatment plant yields a novel alcohol/aldehyde dehydrogenase. *Environ Microbiol* **7**: 1917–1926.
- White CP, Debry RW, Lytle DA. (2012). Microbial survey of a full-scale, biologically active filter for treatment of drinking water. *Appl Environ Microbiol* **78**: 6390–6394.
- Wilmes P, Simmons SL, Denev VJ, Banfield JF. (2009). The dynamic genetic repertoire of microbial communities. *FEMS Microbiol Rev* **33**: 109–132.
- Wu S, Zhu Z, Fu L, Niu B, Li W. (2011). WebMGA: a customizable web server for fast metagenomic sequence analysis. *BMC Genomics* **12**: 444.
- Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y. (2012). dbCAN: a web resource for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res* **40**: W445–W451.

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)