SHORT COMMUNICATION

Recent expansion of heat-activated retrotransposons in the coral symbiont *Symbiodinium microadriaticum*

Jit Ern Chen¹, Guoxin Cui¹, Xin Wang, Yi Jin Liew and Manuel Aranda

Red Sea Research Center, Division of Biological and Environmental Science and Engineering, King Abdullah University of Science and Technology, Thuwal, Saudi Arabia

Rising sea surface temperature is the main cause of global coral reef decline. Abnormally high temperatures trigger the breakdown of the symbiotic association between corals and their photosynthetic symbionts in the genus *Symbiodinium*. Higher genetic variation resulting from shorter generation times has previously been proposed to provide increased adaptability to *Symbiodinium* compared to the host. Retrotransposition is a significant source of genetic variation in eukaryotes and some transposable elements are specifically expressed under adverse environmental conditions. We present transcriptomic and phylogenetic evidence for the existence of heat stress-activated Ty1-*copia*-type LTR retrotransposons in the coral symbiont *Symbiodinium microadriaticum*. Genome-wide analyses of emergence patterns of these elements further indicate recent expansion events in the genome of *S. microadriaticum*. Our findings suggest that acute temperature increases can activate specific retrotransposons in the *Symbiodinium* genome with potential impacts on the rate of retrotransposition and the generation of genetic variation under heat stress.

The ISME Journal (2018) 12, 639–643; doi:10.1038/ismej.2017.179; published online 20 October 2017

Coral reefs are notable for being one of the world's most productive and diverse ecosystems despite flourishing in highly oligotrophic environments (de Goeij *et al.*, 2013). This paradox exists because of the endosymbiotic relationship between corals and unicellular algae of the genus *Symbiodinium*, whereby corals provide nutrients and shelter to the algae in exchange for energy in the form of photosynthetically fixed carbon (Muscattine and Porter, 1977). In the past few decades, most coral ecosystems around the world have been gradually declining, a process that has been accelerated by the warming of the oceans (Ainsworth *et al.*, 2016). The coral *Symbiodinium* symbiosis is notably sensitive to increases in seawater temperature, which can lead to widespread expulsion of the symbiont and eventual starvation of the coral hosts if the symbiotic relationship is not restored (Hoegh-Guldberg and Smith, 1989). However, the presence of thermotolerant corals, such as those in the Red Sea (Fine *et al.*, 2013), strongly suggests that there is room for adaptation to higher ocean temperatures. One component that determines coral thermotolerance is the presence of

Symbiodinium strains that are more resistant to heat stress (Howells *et al.*, 2012). Given the shorter generation times of *Symbiodinium* relative to their coral hosts, it has been speculated that the higher rate at which genetic variation increases in *Symbiodinium* could play a major role in the evolution of coral–algal symbiosis with greater thermotolerance (Csaszar *et al.*, 2010; van Oppen *et al.*, 2017).

One way in which rapid and environmentally triggered genomic variation could be generated is via retrotransposons, whose capability to replicate independently of the host genome makes them particularly interesting as one of the causal agents of natural insertional mutagenesis (Kazazian, 2004). While most retrotransposons are transcriptionally silenced by the host genome, several lines of evidence suggest that these genetic elements can be activated under certain stress conditions such as wounding (Takeda *et al.*, 1998) or UV irradiation (Ramallo *et al.*, 2008).

A specific class of retrotransposons, defined by the presence of flanking long terminal repeats (LTR), is notable for being transcriptionally activated by a range of stress conditions in fungi (Anaya and Roncero, 1996), metazoans (Faure *et al.*, 1996) and higher plants (Ito *et al.*, 2011). An example of this is ONSEN, a Ty1-*copia*-type LTR retrotransposon in *Arabidopsis thaliana* that was shown to be activated by elevated temperatures and capable of generating *A. thaliana* lines resistant to abscisic acid, thereby linking heat stress with the generation of novel

Correspondence: M Aranda, Red Sea Research Center, Division of Biological and Environmental Science and Engineering, King Abdullah University of Science and Technology, Ibn Al Haytham, 4700 KAUST, Thuwal, Makkah 23955-6900, Saudi Arabia.
E-mail: manuel.aranda@kaust.edu.sa

¹These authors contributed equally to this work.

Received 12 February 2017; revised 28 August 2017; accepted 15 September 2017; published online 20 October 2017

Figure 2 Genome-wide analyses of *copia*-like elements. **(a)** Maximum likelihood phylogeny of 365 reverse transcriptase amino-acid sequences curated *in silico* from the *S. microadriaticum* genome. Green lines indicate known Ty1-*copia*-type sequences, orange lines indicate SmTCL1-4, pink lines indicate retrovirus, caulimovirus and Ty3-*gypsy*-like outgroup sequences as seen in Figure 1c and Supplementary Figure S1, black lines indicate Ty1-*copia*-like sequences. Sequences considered to form the SmTCL subclades are highlighted in yellow. Bootstrap values below 70 were removed from the phylogeny. Scale bar indicates number of substitutions per site. **(b)** Jukes-Cantor distribution of selected retrotransposon families in the *S. microadriaticum* genome. Insert plot shows the distribution of retrotransposons within the 0–0.01 Jukes-Cantor distance fraction, at bin sizes of 0.001. SmTCL (purple line) shows the genome coverage of sequences associated with the heat responsive elements SmTCL1-4. A sequence with 50% of its nucleotide positions different from its consensus sequence would have a Jukes-Cantor distance of 0.824 according to the equation, $d = -(\frac{3}{4}) \times \ln(1 - (\frac{4}{3})p)$, where p is the proportion of sites with different nucleotides and d is the Jukes-Cantor distance.

Figure 1 Evidence for putative heat stress-activated Ty1-*copia*-like retrotransposons. **(a)** qPCR expression analysis of selected genes under heat stress. SmTCL, *S. microadriaticum* Ty-*copia*-like; Fox-1, RNA-binding protein fox-1 homolog; CYPLIA8, Cytochrome P450. Error bars show the standard error of the means. **one-tailed *t*-test $P < 0.01$; *** $P < 0.001$. **(b)** Domain arrangements of various LTR retroelements. LTR, long terminal repeats; PBS, primer binding site; gag, group-specific antigen; pol, pol polyprotein; PR, protease; INT, integrase; RT, reverse transcriptase; RH, ribonuclease H; PPT, polypurine tract; env, envelope protein. **(c)** Maximum likelihood phylogeny of 32 reverse transcriptase or RNA-dependent RNA polymerase amino-acid domains of various retrotransposons and viruses. Dotted lines indicate polyphyletic groupings and the scale bar indicates number of substitutions per site. Node support values indicate bootstrap support.

phenotypes via retrotransposon-mediated mutagenesis (Ito *et al.*, 2011, 2016).

We have identified four sequences in the *Symbiodinium microadriaticum* genome (Aranda *et al.*, 2016) as putative Ty1-*copia* class LTR retrotransposons using previously published transcriptomic data (Liew *et al.*, 2017), henceforth referred to as *S. microadriaticum* Ty1-*Copia-Like* (*SmTCL*) 1, 2, 3 and 4. The first two were identified as being the two most upregulated genes in response to 36 °C temperature stress (Supplementary Table S1) in the assembled transcriptome. In addition, we identified two further heat stress-activated *SmTCLs* by conducting an independent differential expression analysis on a set of genome-curated sequences that were pre-identified as Ty1-*copia* sequences (Supplementary Table S2).

To confirm that the overexpression of these sequences was indeed a direct consequence of the heat stress treatment, we repeated the RNA extraction experiment twice and validated the differential expression of *SmTCL*1, 2 and 3 using quantitative reverse transcription polymerase chain reaction (RT-qPCR) (Figure 1a). All three *SmTCLs* were significantly upregulated in heat-stressed samples. In addition, we also tested the expression of *Fox-1* and *CYPLIIA8*, which were the two most down-regulated transcripts in our RNA-Seq data set (Supplementary Table S1).

Analysis of the genetic structure showed that all four *SmTCLs* possess a reverse transcriptase (RT) domain, an integrase domain upstream of said RT domain, a RNase H domain and are flanked by LTRs (Figure 1b). The position of the integrase domain plays a key role in the classification of the two major classes of LTR retrotransposons: Ty1-*copia* and Ty3-*gypsy*. The domain arrangement of Ty3-*gypsy* is very similar to that of retroviruses (Figure 1b), with the integrase domain being downstream of the reverse transcriptase domain. We also built a phylogenetic tree (Figure 1c) using the aligned amino-acid sequences of 28 model reverse transcriptases (Supplementary Figure S1) from a diverse variety of retroelements (Xiong and Eickbush, 1990) in order to classify the *SmTCLs*. We found that all *SmTCLs* clustered together with the Ty1-*copia* family of retrotransposons. To our knowledge, this is the first reported example of heat-activated retrotransposons in *S. microadriaticum* in particular, and dinoflagellates in general.

We determined the prevalence of *SmTCL* RT isoforms in the genome using genome-wide BLAST analysis and identified 343 RT sequences, which were classified as Ty1-*copia*-like sequences based on phylogenetic analysis relative to a retrovirus and Ty3-*gypsy* outgroup (Figure 2a). The *SmTCLs* grouped separately, indicating that these sequences diverged some time ago.

A genome-wide analysis of the Jukes–Cantor distribution of LTR retrotransposons (Chapman *et al.*, 2010) in the *Symbiodinium* genome indicates

that *copia*-like sequences have had a recent increase in copy numbers, as evidenced by a large number of elements showing almost no sequence divergence (0–0.01 Jukes–Cantor distance) (Figure 2b). In terms of genome coverage, the plurality of *copia* sequences are considered near-identical to their consensus sequences indicating that they have not had the time to acquire nucleotide changes. Interestingly, we found that the heat responsive elements *SmTCLs*1–4 constituted ~26% of the genome coverage of the lowest divergence fraction (0–0.001 Jukes–Cantor distance), indicating that these elements expanded recently (Supplementary Table S3). For *SmTCL*3, for instance, we identified five near-identical full-length copies (>99.9% similarity over ~10 kb) as well as a 500 bp copy with the same LTRs (Supplementary Data S1).

In the diatom *Phaeodactylum tricornutum*, expression of the Ty1-*copia*-like retrotransposons *Black-beard* was found to be induced by nitrate starvation (Maumus *et al.*, 2009). Different accessions of *P. tricornutum* were shown to possess different retrotransposition events that were distinguishable at the population level. Like *Symbiodinium*, *P. tricornutum* has not been directly observed to undergo sexual reproduction (Levin *et al.*, 2016). Unlike *Phaeodactylum*, *Symbiodinium* has a haploid genome (Santos and Coffroth, 2003) and thus retrotransposition into a functional genomic region would likely have a stronger phenotypic effect. Retrotransposon-mediated stress-activated mutagenesis could thus be a mechanism to generate additional genetic variation under adverse conditions, where the cost of deleterious mutations is outweighed by the possibility of acquiring novel adaptations (Rey *et al.*, 2016).

Our results do not provide evidence for new, heat stress-induced retrotransposition events, but they suggest that mutation rates for *S. microadriaticum* could increase substantially due to stress-mediated transcriptional activation of transposons, leading to a greater-than-expected increase in genetic variation under acute heat stress (Chakravarti *et al.*, 2017). Further work investigating whether isolated, post-heat-stressed *S. microadriaticum* possess more LTR retrotransposon copy numbers compared to non-stressed control cultures would be the first step to establishing whether these new retrotransposition events are capable of generating new phenotypes from a clonal parental population. This would provide some indication whether it would be feasible to generate phenotypes with increased coral symbiosis resilience to heat stress via heat stress-induced ‘experimental evolution’ of *S. microadriaticum* strains (van Oppen *et al.*, 2017). Additionally, *in natura* analyses comparing LTR retrotransposon copy numbers in *Symbiodinium* populations from sites differing in temperature fluctuations could provide further evidence for the ecological relevance of such a mechanism under natural conditions.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

This publication is based on the work supported by the King Abdullah University of Science and Technology (KAUST) Office of Sponsored Research (OSR) under Award No. URF/1/1705-01.

References

- Ainsworth TD, Heron SF, Ortiz JC, Mumby PJ, Grech A, Ogawa D *et al.* (2016). Climate change disables coral bleaching protection on the Great Barrier Reef. *Science* **352**: 338–342.
- Anaya N, Roncero MIG. (1996). Stress-induced rearrangement of *Fusarium* retrotransposon sequences. *Mol Gen Genet* **253**: 89–94.
- Aranda M, Li Y, Liew YJ, Baumgarten S, Simakov O, Wilson MC *et al.* (2016). Genomes of coral dinoflagellate symbionts highlight evolutionary adaptations conducive to a symbiotic lifestyle. *Sci Rep* **6**: 39734.
- Chakravarti LJ, Beltran VH, van Oppen MJH. (2017). Rapid thermal adaptation in photosymbionts of reef-building corals. *Glob Chang Biol* 2017; **23**: 4675–4688.
- Chapman JA, Kirkness EF, Simakov O, Hampson SE, Mitros T, Weinmaier T *et al.* (2010). The dynamic genome of *Hydra*. *Nature* **464**: 592–596.
- Csaszar NBM, Ralph PJ, Frankham R, Berkelmans R, van Oppen MJH. (2010). Estimating the potential for adaptation of corals to climate warming. *PLoS One* **5**: e9751.
- de Goeij JM, van Oevelen D, Vermeij MJ, Osinga R, Middelburg JJ, de Goeij AF *et al.* (2013). Surviving in a marine desert: the sponge loop retains resources within coral reefs. *Science* **342**: 108–110.
- Faure E, BestBelpomme M, Champion S. (1996). X-irradiation activates the *Drosophila* 1731 retrotransposon LTR and stimulates secretion of an extracellular factor that induces the 1731-LTR transcription in nonirradiated cells. *J Biochem* **120**: 313–319.
- Fine M, Gildor H, Genin A. (2013). A coral reef refuge in the Red Sea. *Glob Change Biol* **19**: 3640–3647.
- Hoegh-Guldberg O, Smith GJ. (1989). The effect of sudden changes in temperature, light and salinity on the population-density and export of zooxanthellae from the reef corals *Stylophora pistillata* Esper and *Seriatopora hystris* Dana. *J Exp Mar Biol Ecol* **129**: 279–303.
- Howells EJ, Beltran VH, Larsen NW, Bay LK, Willis BL, van Oppen MJH. (2012). Coral thermal tolerance shaped by local adaptation of photosymbionts. *Nat Clim Change* **2**: 116–120.
- Ito H, Gaubert H, Bucher E, Mirouze M, Vaillant I, Paszkowski J. (2011). An siRNA pathway prevents transgenerational retrotransposition in plants subjected to stress. *Nature* **472**: 115–119.
- Ito H, Kim JM, Matsunaga W, Saze H, Matsui A, Endo TA *et al.* (2016). A stress-activated transposon in *Arabidopsis* induces transgenerational abscisic acid insensitivity. *Sci Rep* **6**: 23181.
- Kazazian HH Jr. (2004). Mobile elements: drivers of genome evolution. *Science* **303**: 1626–1632.
- Levin RA, Beltran VH, Hill R, Kjelleberg S, McDougald D, Steinberg PD *et al.* (2016). Sex, scavengers, and chaperones: transcriptome secrets of divergent *Symbiodinium* thermal tolerances. *Mol Biol Evol* **33**: 2201–2215.
- Liew YJ, Li Y, Baumgarten S, Voolstra CR, Aranda M. (2017). Condition-specific RNA editing in the coral symbiont *Symbiodinium microadriaticum*. *PLoS Genet* **13**: e1006619.
- Maumus F, Allen AE, Mhiri C, Hu HH, Jabbari K, Vardi A *et al.* (2009). Potential impact of stress activated retrotransposons on genome evolution in a marine diatom. *BMC Genomics* **10**: 624.
- Muscatine L, Porter JW. (1977). Reef corals - mutualistic symbioses adapted to nutrient-poor environments. *BioScience* **27**: 454–460.
- Ramallo E, Kalendar R, Schulman AH, Martinez-Izquierdo JA. (2008). Reme1, a Copia retrotransposon in melon, is transcriptionally induced by UV light. *Plant Mol Biol* **66**: 137–150.
- Rey O, Danchin E, Mirouze M, Loot C, Blanchet S. (2016). Adaptation to global change: a transposable element-epigenetics perspective. *Trends Ecol Evol* **31**: 514–526.
- Santos SR, Coffroth MA. (2003). Molecular genetic evidence that dinoflagellates belonging to the genus *Symbiodinium freudenthal* are haploid. *Biol Bull* **204**: 10–20.
- Takeda S, Sugimoto K, Otsuki H, Hirochika H. (1998). Transcriptional activation of the tobacco retrotransposon Tto1 by wounding and methyl jasmonate. *Plant Mol Biol* **36**: 365–376.
- van Oppen MJH, Gates RD, Blackall LL, Cantin N, Chakravarti LJ, Chan WY *et al.* (2017). Shifting paradigms in restoration of the world's coral reefs. *Glob Chang Biol* **23**: 3437–3448.
- Xiong Y, Eickbush TH. (1990). Origin and evolution of retroelements based upon their reverse transcriptase sequences. *EMBO J* **9**: 3353–3362.

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>

© The Author(s) 2018

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)