MINI REVIEW

Capturing the genetic makeup of the active microbiome *in situ*

Esther Singer¹, Michael Wagner² and Tanja Woyke¹¹US Department of Energy Joint Genome Institute, Walnut Creek, CA, USA and ²University of Vienna, Department of Microbial Ecology and Ecosystem Science, Division of Microbial Ecology, University of Vienna, Vienna, Austria

More than any other technology, nucleic acid sequencing has enabled microbial ecology studies to be complemented with the data volumes necessary to capture the extent of microbial diversity and dynamics in a wide range of environments. In order to truly understand and predict environmental processes, however, the distinction between active, inactive and dead microbial cells is critical. Also, experimental designs need to be sensitive toward varying population complexity and activity, and temporal as well as spatial scales of process rates. There are a number of approaches, including single-cell techniques, which were designed to study *in situ* microbial activity and that have been successively coupled to nucleic acid sequencing. The exciting new discoveries regarding *in situ* microbial activity provide evidence that future microbial ecology studies will indispensably rely on techniques that specifically capture members of the microbiome active in the environment. Herein, we review those currently used activity-based approaches that can be directly linked to shotgun nucleic acid sequencing, evaluate their relevance to ecology studies, and discuss future directions.

The ISME Journal (2017) 11, 1949–1963; doi:10.1038/ismej.2017.59; published online 2 June 2017

Introduction

Microbial communities perform essential roles in natural and engineered ecosystems. They contribute to the dynamics and complexity of global biogeochemical cycles and are the main drivers for important biotechnological processes. One of the main challenges in evaluating the microbial impact in a given system is that organism abundance, diversity and (re-)action potential can vary across systems and across different temporal and spatial scales and may be correlated with a combination of biological, chemical and physical stimuli. The identification of important microbial factors in a given habitat using 16S ribosomal RNA (rRNA) gene surveys (Tringe and Hugenholtz, 2008), shotgun metagenomics (Handelsman, 2004) and single-cell genomics (Lasken, 2007) has provided important insights into the potential functions of particular microbes. However, using these techniques in their traditional sense does not offer the possibility to fully capture the plasticity of microbial communities and its effect on local and global processes. Instead, it is becoming essential to complement sequencing with methods that distinguish cells that are growing and metabolically active from those that are dormant or

dead (or even extracellular DNA), and to understand under what circumstances microbial dormancy could be followed by stochastic awakening (Buerger *et al.*, 2012). Meaningful ecological units, such as microbial guilds (Simberloff and Dayan, 1991), could be identified by monitoring the activity of individual microbes in relevant time series experiments coupled with spatially comprehensive study designs. Several of the discussed tools offer single-cell resolution and thus provide spatial resolution in addition to information on phylogeny and metabolism (Table 1, Figure 1). This review is aimed at highlighting the different ecological and physiological insights into active community members of a given microbiome that can be gained with currently used approaches (in chronological order of published studies) in conjunction with shotgun nucleic acid sequencing. We note that methods that yield activity-labeled samples *in situ* cannot provide answers to all ecological challenges, however, when applicable, these tools present valuable assets in microbial ecology research. We further discuss prospects of exciting emerging technologies in this field.

An array of methods available for studying microbial *in situ* activity linked to shotgun sequencing

Methods detecting microbial *in situ* activity target a variety of physiological processes to determine the

Correspondence: T Woyke, US Department of Energy Joint Genome Institute, Walnut Creek, CA 94598, USA.
E-mail: twoyke@lbl.gov

Received 1 November 2016; revised 2 March 2017; accepted 10 March 2017; published online 2 June 2017

Table 1 Comparison of basic characteristics associated with selected techniques for studying active microbial populations *in situ* that can be combined with shotgun sequencing

<i>Cell process targeted</i>	<i>Technique</i>	<i>Advantages</i>	<i>Disadvantages</i>	<i>Biases/assumptions</i>	<i>Reference</i>
DNA synthesis	BrdU labeling (B, S)	Can be used as single-cell tool if combined with fluorescent antibody staining	Low labeling efficiency	Rate of uptake varies by cell; may be toxic to some cells	Borneman, 1999; Yin <i>et al.</i> , 2000*
	DNA-SIP (B)	Link of metabolic function to identity	Long incubation time; cross-feeding problem; contamination prone; high GC DNA problematic; dependent on commercially available labeled compounds ^a	Cell division may be stimulated by substrate addition and not reflect <i>in situ</i> growth rates	Radajewski <i>et al.</i> , 2000*
	Peak-to-trough ratios in metagenomes (S+B), iRep	No sample incubation necessary; samples can be frozen immediately	Reliant on database; affected by community complexity.		Korem <i>et al.</i> , 2015;* Brown <i>et al.</i> , 2016*
Transcription	RNA-SIP (B)	High phylogenetic resolution; link of metabolic function to taxonomic identity	Dependent on commercially available labeled compounds; cross-feeding potential ^b	Cell division may be stimulated by substrate addition and not reflect <i>in situ</i> growth rates	Radajewski <i>et al.</i> , 2003*
	Metatranscriptomics (S+B)	Samples can be frozen immediately; inexpensive; fast	mRNA abundance not necessarily indicative of protein levels and enzymatic activities; high transcriptome coverage necessary; modulation may occur during sampling	RNA abundance does not directly correlate with cell activity	Leininger <i>et al.</i> , 2006
	BONCAT (+FISH/FACS) (S+B)	No known effect on growth and translation activity	High amino acid conc. in sample diminishes/suppresses signal	Cell growth may be stimulated by added amino acids; dependence on uptake mechanism	Hatzenpichler <i>et al.</i> , 2014; Hatzenpichler <i>et al.</i> , 2016*
Lipid, carbohydrate, amino-acid biosynthesis	D ₂ O+Raman microscopy (S)	D ₂ O concentration required not known to be toxic to any cells; no stimulation effect expected in hydrated samples; can be combined with FISH; stimulation experiments with unlabeled substrates possible	Low-throughput so far	Different physiologies lead to different incorporation rates of deuterium (for example, autotrophs vs heterotrophs)	Berry <i>et al.</i> , 2015*
Bacterial reduction potential	Redox sensor green+FACS (S+B)	Rapid response to redox activity within cell; stable fluorescence signal; quick cell penetration; no detrimental effect on core cell functions known; stimulation experiments with unlabeled substrates possible	Incubated cells cannot be frozen; not all microbial species may transport redox sensor green equally well into their cells		Kalyuzhnaya <i>et al.</i> , 2008*

Techniques are further characterized as allowing for single-cell approaches (S) and/or bulk community analysis (B). References given denote those of either first use of the respective method or the first combined use with (any) sequence data (marked with an asterisk).
Abbreviations: BONCAT, biorthogonal non-canonical amino-acid tagging; BrdU, bromodeoxyuridine; FACS, fluorescence-activated cell sorting; FISH, fluorescence *in situ* hybridization; iRep, index of replication; SIP, stable isotope probing.
^aDNA-SIP may be used with ¹⁸O-H₂O, which—to our knowledge—has not been used with metagenomics or single-cell sequencing.
^bRNA-SIP may be used with ¹⁸O-H₂O, which—to our knowledge—has not been used with metagenomics or single-cell sequencing.

Figure 1 Methods that yield activity-labeled samples and are targeting cell processes in an 'active' microbial cell that can be coupled with shotgun sequencing. Colors denote resources (green), cell components (blue) and cell processes (orange). Raman: Raman microspectroscopy. For DNA- and RNA-SIP, total nucleic acids are extracted from the samples, and labeled and unlabeled DNA/RNA is separated by density gradient centrifugation. The 'heavier' labeled nucleic acid fractions can be used for construction of metagenomic libraries (Neufeld *et al.*, 2007; Whiteley *et al.*, 2007), whereas PLFAs are analyzed on a mass spectrometer and cannot be combined with nucleic acid sequencing (Jehmlich *et al.*, 2010). Many ^{13}C -, ^{18}O -, ^{15}N -labeled fine chemicals are available (for example, phenol, methanol, ammonia, methane, carbonate, etc.), but the wide-ranging application of SIP is limited by the commercial availability of complex labeled compounds that require expensive custom synthesis. Furthermore, sensitivity of the SIP technique is a function of substrate concentration and the duration of substrate incorporation. Successful SIP is dependent on optimization of substrate concentration to guarantee a significant signal-to-noise ratio and incubation length and avoid enrichment bias (Neufeld *et al.*, 2006) (Table 1). 'Cross-feeding', that is, the flow of the isotope label from primary metabolizers to secondary consumers has also been documented (Hutchens *et al.*, 2004; Dumont *et al.*, 2006). RSG is a fluorogenic redox indicator dye available from Molecular Probes, Invitrogen (Carlsbad, CA, USA). RSG yields green fluorescence (488 nm excitation) when modified by bacterial reductases, many of which are parts of electron transport systems. SIP-Raman microspectroscopy has been performed using ^{13}C -, ^{15}N -labeled compounds, as well as with D_2O . The addition of D_2O (up to a certain concentration and for limited time) is expected to have negligible effects on the microbial community composition and activity patterns, for example, compared with nutrient substrates (Lester *et al.*, 1960; Berry *et al.*, 2015; Kopf *et al.*, 2015) that are traditionally used for SIP experiments. Incorporation of D_2O -derived deuterium into the biomass of autotrophic and heterotrophic bacteria and archaea can be unambiguously detected via C-D signature peaks in single-cell Raman spectra (Ashkin, 1970; Berry *et al.*, 2015). However, for comparative studies between active taxa it should be kept in mind that microbes with different physiologies will incorporate different amounts of deuterium at similar growth rates.

activity state of a cell. The earliest developed microbial method that yields activity-labeled samples *in situ* and can be combined with shotgun sequencing takes advantage of the incorporation of bromodeoxyuridine (BrdU) into newly synthesized DNA (and RNA) of a viable cell and subsequent density gradient centrifugation or immunocapture of BrdU-labeled DNA (Urbach *et al.*, 1999). BrdU labeling has been successfully used in a variety of environments to identify DNA-synthesizing populations. Studies have investigated phosphate stimulation in soil (Borneman, 1999), the active population of lake water (Urbach *et al.*, 1999), bacterial functional redundancy with respect to various

carbon substrates in soil (Yin *et al.*, 2000), microbial association with arbuscular mycorrhizal hyphae (Artursson and Jansson, 2003), microbial response to inoculation of various plants with arbuscular mycorrhizal (Artursson *et al.*, 2005), nitrogen addition in Alaskan boreal forest (Handelsman, 2004; Allison *et al.*, 2008), vertical redox gradients in Baltic Sea sediment (Lasken, 2007; Edlund *et al.*, 2008), fungal response to different carbon sources in forest soils (Hanson *et al.*, 2008; Buerger *et al.*, 2012; McGuire *et al.*, 2012) and microbial response to carbon sources of varying recalcitrance in soil (Urbach *et al.*, 1999; Goldfarb *et al.*, 2011). In addition, studies have targeted bacterial and archaeal

(Borneman, 1999; Yin *et al.*, 2000; Artursson and Jansson, 2003; Artursson *et al.*, 2005; Goldfarb *et al.*, 2011) as well as fungal (Urbach *et al.*, 1999; Allison and Treseder, 2008; Allison *et al.*, 2008; Hanson *et al.*, 2008) DNA synthesis activity in response to various environmental stimuli in soils and sediments (Yin *et al.*, 2000; Edlund *et al.*, 2008). Although BrdU-labeling experiments can help identify replicating microorganisms and determine average activity in response to various changing environmental conditions, organisms vary in their ability to incorporate the relatively large BrdU molecule into their DNA, and the rate at which incorporation occurs can deviate up to 10-fold between cells (Urbach *et al.*, 1999; Artursson and Jansson, 2003; Hellman *et al.*, 2011). Such variability may lead to incomplete and skewed representation of the active microbial population in a given environment. In addition, owing to low labeling efficiency, BrdU labeling linked to sequencing requires a relatively large number of sample replicates to obtain the amount of DNA needed for 16S rRNA gene and metagenome sequencing, which renders this technique relatively costly and labor intensive (Tables 1 and 2). The aforementioned studies have yielded important insights into the functioning of microbial communities, yet combining BrdU labeling with either fluorescence-activated cell sorting (FACS) and subsequent high-throughput sequencing or solely with high-throughput sequencing has found limited application to date (Mou *et al.*, 2008; David *et al.*, 2014; Hamasaki *et al.*, 2016). Published studies have predominantly included PCR-amplified rRNA gene surveys and only a few metagenome studies are available (Figure 2, Table 3). However, as DNA requirements for shotgun sequencing decrease, sequencing of BrdU-labeled environmental DNA may become more prevalent.

Stable isotope probing (SIP) labeling technology can be used to selectively enrich the DNA or RNA of active microbial populations (Radajewski *et al.*, 2003). For SIP analysis, stable isotope (for example, ^2H , ^{13}C , ^{15}N or ^{18}O) labeled substrates are added to

the environment and as these substrates are consumed by active organisms, labeled atoms are incorporated in their DNA or RNA. Radajewski *et al.* (2000) first demonstrated this approach by adding various ^{13}C -enriched carbon sources ($^{13}\text{CH}_3\text{OH}$, $^{13}\text{CH}_4$) to soil and subsequently identifying active methylotrophs. The authors found that methylotrophy was confined to *Alphaproteobacteria* lineages and identified various new methanol dehydrogenase alpha subunit (*mxoF*) sequences. DNA-SIP is currently the most applied method, having been used by many research groups to study *in situ* microbial activity (Figure 2). RNA-SIP provided insights that led to the isolation and subsequent genome sequencing of a *Thauera* sp., which was identified as key phenol degrading organisms in an industrial wastewater treatment plant (Manefield *et al.*, 2002). Recently, Dumont *et al.* (2013) obtained a $^{13}\text{CH}_4$ labeled-RNA metatranscriptome from methanotrophs in lake sediment. The labeled metatranscriptome was predominantly enriched in sequences from *Methylococcaceae* and transcripts of methane monooxygenase (*pmoCAB*) genes. The authors suggested that SIP and metatranscriptomics can be broadly applicable, but this approach has only been used one other time to our knowledge (Fortunato and Huber, 2015). Although limitations of DNA- and RNA-SIP include availability of labeled compounds, cross-feeding, sensitivity and potential for enrichment bias (Figure 1, Table 1), continuous efforts to optimize SIP, applying it to wider ranges of environments, and combining it with other techniques, such as Raman microspectroscopy and fluorescence *in situ* hybridization (FISH; Huang *et al.*, 2007) make DNA-SIP the best current method for sequence-based characterization of key populations synthesizing DNA in the presence of isotope-labeled compounds.

Metatranscriptome studies enable the analysis of expressed genes in environmental communities, portraying the near real-time condition of microorganisms because of the relatively short half-life of mRNA (Moran *et al.*, 2012). Studies of microbial communities in soil and marine environments were

Table 2 Comparison of practical considerations of activity-based techniques in combination with sequencing

Technique	Incubation time	Labor intensity	Cost of disposables
BrdU labeling	Hours to days	H	\$\$-\$\$\$
DNA-SIP	Hours to days	M	\$\$
Peak-to-trough ratios in metagenomes, iRep	NA	L-M	\$
RNA-SIP	Minutes	M	\$\$
Metatranscriptomics	NA	L	\$
BONCAT (+FACS)	Hours to days	H	\$\$\$*
D ₂ O+Raman spectrometry	Hours to days	H (currently because of lack of automation)	\$\$\$*
Redox sensor green+FACS	10 min	H	\$\$\$*

Labor intensity is a rough estimate of time required to handle 10 retrieved samples/data sets, where 1 sample may either be 1 cell or a population: L ~ <1 week; M ~ 1–2 weeks; H ~ 2–4 weeks. Cost of disposables is divided into three brackets for processing and analysis of 10 individual samples (excluding replicates): \$ ~ <\$100; \$\$ ~ \$100–500; \$\$\$ ~ >\$500. These costs do not include labor or equipment time. Cost differences that may occur depending on whether samples are defined as single cells, which require MDA, or as sub-populations were determined to be negligible for 10 samples. Costs affected by this sample differentiation are denoted with an asterisk.

Abbreviations: BONCAT, bioorthogonal non-canonical amino-acid tagging; BrdU, bromodeoxyuridine; FACS, fluorescence-activated cell sorting; iRep, index of replication; MDA, multiple displacement amplification; NA, not applicable; SIP, stable isotope probing.

Figure 2 Project statistics by method over the last decade. Counts displayed exclusively include projects using high-throughput sequencing. Metagenome (MetaG) and metatranscriptome (MetaT) projects are depicted by lines (primary y axis), whereas BrdU, DNA-SIP, RNA-SIP and PTR projects are displayed as bars (secondary y axis). Project abundances are cumulative. Number of MetaG and MetaT projects include public sequencing projects as recorded in the Genomes OnLine Database (GOLD) (Pagani *et al.*, 2011) retrieved 15 January 2016. Number of all other activity-based projects include published records that feature high-throughput (next-generation) shotgun sequence data.

the first to use high-throughput sequencing technology for mRNA analysis (Leininger *et al.*, 2006; Frias-Lopez *et al.*, 2008; Urich *et al.*, 2008). Since then, several studies in various environments have shown that only a subset of the total microbial community contributes to the sequenced RNA pool, and can therefore be considered transcriptionally active. For example, Lesniewski *et al.* (2012) were able to show that transcriptional microbial activity of hydrothermal plume communities was largely attributed to water column microorganisms rather than members from the seafloor. mRNA is regularly used as an indicator of microbial activity and is logistically advantageous because samples can be frozen instantaneously, thereby enabling delayed processing after freezing, which is not an option for other methods that yield activity-labeled samples *in situ* (Table 1). This is relevant when *in situ* conditions cannot easily be maintained during sampling or downstream sample processing. A critical issue is how deeply a community transcriptome is 'covered' by the sequence data. If too shallow, libraries will be dominated by transcripts from metabolic pathways shared by most cells and lacking in those representing specialized biogeochemical pathways (Hewson *et al.*, 2009; Gifford *et al.*, 2010; Poretsky *et al.*, 2010). As a consequence, unique expression patterns within a community may be missed, and comparative analyses between communities can become insensitive. Also, depletion of rRNA before metatranscriptome sequencing is not always trivial. Nevertheless, the near-real-time response of mRNA levels to environmental changes experienced by a cell renders assessing inventories of mRNA pools highly informative for identifying ecologically

relevant processes while also determining the identities of their microbial drivers. Declining costs, greater sequencing depth and more comprehensive genomic databases render metatranscriptomics an increasingly powerful and popular tool.

With the advent of fluorescence microscopy activity stains have found an application in microbial ecology. These include stains that rely on cell membrane integrity, enzyme activity or the presence of biomolecules (lipids/nucleic acid). For instance, redox sensor green (RSG) and denaturing gradient gel electrophoresis (DGGE) profiling were used to detect and enumerate microbes actively metabolizing the C_1 substrates methane, methanol, methylamine and formaldehyde in Lake Washington sediment (Kalyuzhnaya *et al.*, 2008). According to Kalyuzhnaya *et al.*, RSG does not suppress cellular metabolism as was previously found for tetrazolium salts, such as CTC (5-cyano-2,3-ditolyl tetrazolium chloride) (Ullrich *et al.*, 1996). Using FACS, RSG could likely be scaled to separate bacterial and archaeal cells with an electron membrane potential in the presence of environmental stimuli, therefore studying population activity levels.

One of the most recently developed methods that yield activity-labeled samples *in situ* is bioorthogonal non-canonical amino-acid tagging (BONCAT). This technique is based on the *in vivo* incorporation of artificial amino acids that carry modifiable chemical tags into newly synthesized proteins (Dieterich *et al.*, 2006). These chemical tags can then be fluorescently labeled via click chemistry (Hatzenpichler *et al.*, 2014). BONCAT has been demonstrated to be effective in labeling the proteomes of a wide range of taxonomically and

Table 3 Availability of nucleic acid sequence data retrieved using *in situ* microbial activity approaches

Technique	Environment	Sequence data	References for shotgun sequencing approaches
BrdU labeling		16S rRNA, MetaG	Mou <i>et al.</i> , 2008; David <i>et al.</i> , 2014
DNA-SIP		16S rRNA, MetaG	Kalyuzhnaya <i>et al.</i> , 2008c; Dumont <i>et al.</i> , 2006; Chen <i>et al.</i> , 2008; Kalyuzhnaya <i>et al.</i> , 2008b; Neufeld <i>et al.</i> , 2008; Sul <i>et al.</i> , 2009; Gutierrez <i>et al.</i> , 2013; Zhang <i>et al.</i> , 2013; Chemerys <i>et al.</i> , 2014; Kim <i>et al.</i> , 2014; Pinnell <i>et al.</i> , 2014; Verastegui <i>et al.</i> , 2014; Werner <i>et al.</i> , 2014; Eyice <i>et al.</i> , 2015; Abu Laban <i>et al.</i> , 2015; Schwartz <i>et al.</i> , 2016
RNA-SIP		16S rRNA, MetaT	Borneman, 1999; Mou <i>et al.</i> , 2008; Haroon <i>et al.</i> , 2013; Fortunato and Huber, 2015
Metatranscriptomics		16S rRNA, MetaT	Leininger <i>et al.</i> , 2006; Dumont <i>et al.</i> , 2006; Frias-Lopez <i>et al.</i> , 2008; Chen <i>et al.</i> , 2008; Urich <i>et al.</i> , 2008; Kalyuzhnaya <i>et al.</i> , 2008b; Neufeld <i>et al.</i> , 2008; Shi <i>et al.</i> , 2009; Sul <i>et al.</i> , 2009; He <i>et al.</i> , 2010; Lin <i>et al.</i> , 2010; Helbling <i>et al.</i> , 2011; Feike <i>et al.</i> , 2011; Damon <i>et al.</i> , 2011; Bomar <i>et al.</i> , 2011; Burow <i>et al.</i> , 2012; Ottman <i>et al.</i> , 2012; Vila-Costa <i>et al.</i> , 2012; Zhang <i>et al.</i> , 2013; Chaparro <i>et al.</i> , 2013; Gutierrez <i>et al.</i> , 2013; Embree <i>et al.</i> , 2013; Carvalhais <i>et al.</i> , 2013; Sanders <i>et al.</i> , 2013; Baker <i>et al.</i> , 2013; Ursell and Knight, 2013; Lim <i>et al.</i> , 2013; Lehenbre <i>et al.</i> , 2013; Orsi <i>et al.</i> , 2013; Chemerys <i>et al.</i> , 2014; Kim <i>et al.</i> , 2014; Pinnell <i>et al.</i> , 2014; Verastegui <i>et al.</i> , 2014; Werner <i>et al.</i> , 2014; Franzosa <i>et al.</i> , 2014; Jorth <i>et al.</i> , 2014; Schwab <i>et al.</i> , 2014; Lamendella <i>et al.</i> , 2014; Satinsky <i>et al.</i> , 2014; Klindworth <i>et al.</i> , 2014; Hua <i>et al.</i> , 2014; Simón-Soro <i>et al.</i> , 2014; Hilton <i>et al.</i> , 2014; Eyice <i>et al.</i> , 2015; Abu Laban <i>et al.</i> , 2015; Alexander <i>et al.</i> , 2015a; Chapelle <i>et al.</i> , 2015; Bikel <i>et al.</i> , 2015; Bello-Ortí <i>et al.</i> , 2015; Ishii <i>et al.</i> , 2015; Alexander <i>et al.</i> , 2015b; Eren <i>et al.</i> , 2015; Jones <i>et al.</i> , 2015; Steen <i>et al.</i> , 2016; Jiang <i>et al.</i> , 2016; Marzano and Domier, 2016; Kodzius and Gojobori, 2016; Mondot and Lepage, 2016; De Filippis <i>et al.</i> , 2016; Radajewski <i>et al.</i> , 2000; Kalyuzhnaya <i>et al.</i> , 2008a, c; Konopka <i>et al.</i> , 2010
Redox sensor green		16S rRNA	
BONCAT		16S rRNA	Hatzenpichler <i>et al.</i> , 2014; Korem <i>et al.</i> , 2015
Raman+D ₂ O		16S rRNA	Berry <i>et al.</i> , 2015; Brown <i>et al.</i> , 2016
Peak-to-trough ratios in metagenomes, iRep		MetaG	Radajewski <i>et al.</i> , 2003; Leininger <i>et al.</i> , 2006; Hatzenpichler <i>et al.</i> , 2014; Korem <i>et al.</i> , 2015; Brown <i>et al.</i> , 2016; Hatzenpichler <i>et al.</i> , 2016

Sequence data denote those from published studies in listed references. : sediment or soil; : marine or fresh water; : (enrichment) culture or bioreactor; : association studies between bacteria and/or archaea and eukaryotes. References do not include molecular marker studies. Abbreviations: BONCAT, bioorthogonal non-canonical amino-acid tagging; BrdU, bromodeoxyuridine; iRep, index of replication; rRNA, ribosomal RNA; SIP, stable isotope probing. References for metatranscriptomics studies are too comprehensive to list, and thus cover a broad selection of studies.

physiologically distinct archaea and bacteria (Hatzenpichler *et al.*, 2014). For example, dominant cells in an aerobic methanotrophic enrichment culture from deep-sea sediments were identified as *Methylococcaceae*-related gammaproteobacterium based on FISH analyses. The translational activity of these *Methylococcaceae* cells was found to be dependent on the addition of methane to the enrichment culture. Genome information from this *Methylococcaceae* strain confirmed the methanotrophic nature of this bacterium. Rather than

studying the bulk proteome, BONCAT is able to specifically target proteins that have been expressed in response to an experimental condition. The method can also be used in conjunction with rRNA-targeted FISH and thereby allow the simultaneous identification of a microbial cell and its translational activity. In a very recent effort to describe anaerobic methane oxidizers from methane seep sediments, BONCAT was combined with FACS to sort translationally active methane-oxidizing consortia. Whole-genome amplification and 16S rRNA

Figure 3 High-throughput workflows of current and emerging *in situ* microbial activity approaches linked to sequencing. Metatranscriptomics and stable isotope labeling are the most commonly used techniques coupled with next-generation shotgun sequencing technology. Emerging methods that are currently still subject to development and/or optimization involve the incubation of cells and cell clusters with, for example, fluorescent compounds or D_2O before selective sorting of active cells using FACS or Raman OT.

gene sequencing of the aggregates revealed a novel archaeal–bacterial association (Hatzepichler *et al.*, 2016). This technique presents great prospects for future high-throughput sequencing coupled to BONCAT and is expected to help answer questions regarding cellular translational activities in response to environmental cues at micrometer resolution, as

well as in a high-throughput manner across larger scales (Figure 3). Limitations of BONCAT include the diversity of amino-acid uptake mechanisms in the environment, the potential for artificial growth stimulation following addition of amino acids and suppression of signal when environmental concentrations of amino acids are high. Aside from these potential

limitations, we expect BONCAT to be combined with shotgun metagenomics in the near future.

Raman microspectroscopy is an established vibrational spectroscopic technique with emerging prospects in microbial ecology. This non-destructive technique enables chemical fingerprinting of individual eukaryotic cells, organelles, or bacterial and archaeal cells. Chemical information derived from a Raman spectrum provides comprehensive and intrinsic information on the chemical composition of the analyzed cell without the need of external labeling, which, for example, enables microbiologists to identify storage compounds, cytochromes and pigments on a single-cell level within a few seconds (Gruber-Vodicka *et al.*, 2011; Majed *et al.*, 2012; Milucka *et al.*, 2012). Interestingly, the incorporation of the stable isotopes ^{13}C , deuterium and ^{15}N into microbial biomass can also be detected via characteristic band shifts in the Raman spectra of the labeled microbes (Wang *et al.*, 2016), but it should be noted that the ^{15}N -induced shifts are difficult to interpret in complex samples. As Raman microspectroscopy can be combined with FISH (Huang *et al.*, 2007) and optical tweezers (OTs), it offers a culture-independent approach to study the physiology of uncultivated microorganisms in their natural ecosystem. For example, Raman-FISH was applied to identify and measure *in situ* ^{13}C -naphthalene degraders within a microbial community from a complex groundwater system and proved that an uncultured species (*Acidovorax* sp.) had the key role in naphthalene biodegradation, rather than the cultivated naphthalene biodegrading *Pseudomonas* sp. obtained from the same groundwater (Huang *et al.*, 2008). OT-Raman (Raman microspectroscopy coupled with OTs) has been applied to trap and measure single cells of blood (Xie *et al.*, 2002) or *Escherichia coli* (Xie and Li, 2003) and can also be used to sort isotope-labeled microbial cells for subsequent whole genome amplification and sequence-based 16S rRNA gene identification (Huang *et al.*, 2009; Berry *et al.*, 2015). Alternatively, Raman-activated cell ejection using laser-induced forward transfer can be exploited as a low-throughput method for physically removing individual cells from a sample based on their Raman spectrum (Wang *et al.*, 2013; Song *et al.*, 2016).

In a recent study, active microbial cells were identified under near-natural conditions in complex systems via combined use of heavy water (D_2O) and Raman microspectroscopy (Berry *et al.*, 2015). D_2O -Raman spectroscopy can be combined with FISH for parallel identification of the metabolically active microbes. Furthermore, unlabeled substrates can be added and changes in activities of microbial groups of interest can be quantified (Berry *et al.*, 2015; Eichorst *et al.*, 2015). For example, it was revealed that in the cecal microbiota of mice *Akkermansia muciniphila* and *Bacteroides acidifaciens*, two host-compound foragers, exhibited distinctive response patterns to amendments of mucin and sugars (Berry

et al., 2015). By applying D_2O -labeling and OT-Raman, cecal microbial cells stimulated by mucin or glucosamine were obtained for multiple displacement amplification and subsequent identification through 16S rRNA gene sequencing (Berry *et al.*, 2015). Although offering direct access to genomic information from microbes with a defined activity, sorting via OTs is slow. In this context, it is interesting to note that recently a fast Raman-activated cell sorting system (RACS), was designed and used to collect carotenoid-producing yeast cells (Zhang *et al.*, 2015a). Other recent advances include microfluidics-based cell manipulation techniques, such as flow-based OT-RACS and Dielectrophoresis (DEP-) RACS, and advanced Raman techniques, for example, Surface-enhanced Raman spectroscopy (SERS) flow cytometry, Resonance Raman spectroscopy (RRS) cell counting and Coherent anti-Stokes Raman spectroscopy (CARS) flow cytometry, which are discussed elsewhere (Zhang *et al.*, 2015b). Microfluidic device-based Raman sorting can be readily integrated with downstream processes, such as cell incubation, chemical analysis and PCR (Wang *et al.*, 2005). Unlike FACS, Raman sorting can use more criteria, including the incorporation of stable isotopes, to identify and sort individual cells (Table 1).

Finally, two very recently developed non-invasive bioinformatic approaches for linking metagenomics and bacterial activity use sequence read depth across microbial genomes. DNA replication typically begins at a single origin of replication in bacterial and archaeal chromosomes (Mott and Berger, 2007). Thus, during DNA replication, regions that have already been passed by the replication fork will have two copies while the yet unreplicated regions will have a single copy. Korem *et al.* (2015) show that metagenome read coverage patterns for different microbial genomes contain a single trough and a single peak, the latter coinciding with the bacterial origin of replication. The peak-to-trough ratio (PTR) was also demonstrated to provide a quantitative measure of a species growth rate *in vitro* and *in vivo*, under different growth conditions and in complex bacterial communities. For instance, it was shown that the proliferative behavior of virulent and non-virulent strains of *C. rodentium* could be accurately predicted by their PTRs. The application of PTRs is promising, not least, because it considers genomic variance between strains, copy number variation of different genomic regions and variable coverage resulting from sequencing depth. However, it requires information on origin of replication sites of the investigated bacteria and archaea and, on a case-by-case basis, manual correction for relative abundance levels of taxa (Table 1). In an effort to expand this method to metagenome data with draft genomes, Brown *et al.* (2016) developed the index of replication (iRep), which is based on mapping metagenome sequencing reads to the collection of assembled sequences that represent a draft genome. Excluding extremely high and low coverage regions, the overall

slope of coverage across a genome is used to calculate iRep (Brown *et al.*, 2016). Applied to metagenomic data, iRep showed, for instance, that Candidate Phyla Radiation bacteria are generally characterized by slower growth rates compared with various human microbiome organisms. iRep requires a $\geq 75\%$ complete genome in order to produce accurate measurements and was shown to be then correlated to PTRs and experimentally determined growth rates (Brown *et al.*, 2016).

Choosing an appropriate method for assessing microbial activity *in situ*

Our view of microbial activity is strongly shaped by the method and parameters chosen. Microbial contribution to processes can neither always be definitively assessed in every environment nor at a scale that allows interpretation at the systems level. But when experiments are designed carefully to address specific ecological questions at relevant and yet tangible temporal or spatial scales and under consideration of the limitations of the chosen tool (as discussed in the following), methods that yield activity-labeled samples *in situ*, are a great resource. The choice of an appropriate tool for targeting the *in situ* active microbial population will depend on various factors. The physicochemical characteristics of the environment of interest, for example, water saturation, pH, amino acid and sulfide concentration, determine whether a water- or bioorthogonal amino-acid-based labeling method is practical. Although microbial activity is generally correlated with the availability of water in a system, the addition of water may lead to artificially increased microbial response, for example, in systems that are heterogeneously wetted, such as soil aggregates. On the other hand, adding isotope-labeled substrates other than water as typically done in conventional SIP studies changes the nutrient composition of the system and often leads to biases in microbial activity patterns (Table 1). The presence of a reductase activity, DNA synthesis, transcription and translation may occur in cell maintenance mode, as well as during growth and division, which is problematic for the correlation of these processes to quantitative rates of specific functions (Blazewicz *et al.*, 2013). Estimates for maintenance requirements between cultures and microbes living, for example, in low activity subsurface environments vary widely (Hoehler and Jørgensen, 2013). Similarly, ranges of metabolic rates for the same reaction should be modeled for their specific environmental parameters to optimize incubation durations (Hoehler and Jørgensen, 2013). As defining meaningful scales for microbiology research is a recurring challenge (Paerl and Steppe, 2003), the scale of the microbial impact radius determines the choice of method. For example, in a soil environment techniques based on single-cell resolution rarely provide adequate

insights for systems-level mass balance and flux calculations. Each method should be tested for the particular type of microbial community studied before results are interpreted, because discrimination against various cell types or taxa may occur (Berney *et al.*, 2007; Teske and Sørensen, 2007; Narasingarao *et al.*, 2011; Franklin *et al.*, 2013; Lloyd *et al.*, 2013; Eloe-Fadrosh *et al.*, 2016).

As the numbers of bulk metagenome and meta-transcriptome studies have greatly increased over the last few years (Figure 2), we expect other *in situ* activity targeted approaches coupled to shotgun sequencing to follow this trend, as in part already observed by the recent increase in DNA-SIP projects. Although DNA- and RNA-SIP methods are still hampered by throughput, high-throughput pipelines will likely become available for broader community use. FACS-based enrichment of active microbial cells may contribute sequence data in the near future because of its accessibility to users and compatibility with a variety of staining or tagging assays (Figure 3). Similarly, SIP in combination with Raman microspectroscopy and sorting is a promising approach, which may soon become available as high-throughput RACS (Zhang *et al.*, 2015b) (Figure 3). A future venture will be determined by the amenability of methods that yield activity-labeled samples *in situ* to unicellular eukaryotic microbes, including protists and fungi. Although many nucleic acid sequence-based studies encompass well-characterized microbial communities providing a level of understanding of their role in a particular habitat, allowing community modeling (Ruff *et al.*, 2015), we often lack an understanding of eukaryotic microbes. This group of organisms is extremely heterogeneous and there is evidence for its significance in various systems, for example, in airways of cystic fibrosis (Eickmeier *et al.*, 2014), soil (Taylor *et al.*, 2014) and contaminated aquifers (Holmes *et al.*, 2013). On the other hand, inactive eukaryotic microbes may significantly add to the pool of relic DNA. Hence, methods that yield activity-labeled samples *in situ* should include these organisms. Availability and cost-effectiveness of the necessary equipment, as well as the scalability of data throughput will likely guide the future of methods that yield activity-labeled samples *in situ* linked to genomics.

Integrating genomic and functional information from the active microbiome

Capturing the genetic makeup of the active microbiome *in situ* holds great promise to substantially advance our understanding of microbial community functioning. It enables a shift from cataloging the mere presence of taxa and encoded (potential) functions to understanding their ecological roles, including contributions to ecosystem processes. Some of the intriguing and timely research foci in

microbial ecology revolve around expanding our knowledge of microbial guilds, groups of species that exploit the same resource, and of reactions that are thermodynamically feasible enough to sustain microbial growth that are lacking associated microorganisms to date. In these cases, nucleic acid sequencing combined with specific labeled substrate assays and coupled to continuous geochemical profiling would, for instance, allow targeting novel organisms that perform redox reactions of interest and enable comprehensive flux modeling of systems (LaRowe and Amend, 2016). Another timely ecological topic is the significance of so-called ‘strongly interacting species’ (SISs) and temporarily dormant cells. SISs are species that have strong positive or negative impacts on the species they directly interact with (Gibson *et al.*, 2016). SISs are not necessarily keystone species as the removal of SISs from a community does not coercively result in mass extinction (Paine, 1995), but they may include organisms of relatively low abundance (Gibson *et al.*, 2016). SISs can have an important role in shaping the steady states of microbial ecosystems regardless of their relative abundance (Gibson *et al.*, 2016). Hence, identifying SISs and studying the processes in their ‘interaction hub’ in a comparative analysis can help reveal metabolic strategies of ecologically relevant microorganisms and interaction strength heterogeneity, for example, across geochemical gradients. SISs may be periodically dormant and hence require time series sampling and/or stimulation experiments (Epstein, 2013). Dormancy represents a bet-hedging strategy, in which an organism can enter a state of low metabolic activity when environmental conditions are disadvantageous and resuscitate when conditions become favorable (Lennon and Jones, 2011). Dormant cells thus generate seed banks, which are likely to impact diversity dynamics and maintenance in a given ecosystem (Jorgensen, 2011). To better understand the contribution of key population size vs organism state to variable environmental conditions, data on total community (for example, targeted using total DNA), population growth (for example, targeted using BrdU, metagenomics) and specific (for example, targeted using SIP, metatranscriptomics, BONCAT and D₂O-Raman microspectroscopy with substrate amendments), as well as general activity (for example, targeted using BONCAT, metatranscriptomics, RSG, D₂O-Raman microspectroscopy, iRep, PTR) should each be collected in time series experiments. Similarly important is the characterization of the ecological niche of SISs, which may be difficult because of micro-environmental heterogeneity, for instance in soil (Pester *et al.*, 2010) and biofilms (Augspurger *et al.*, 2010). Owing to intricate structures, including channels and voids, a biofilm can host multiple functional groups of cells in various spatial relationships of each other (Gieseke *et al.*, 2003). These spatial arrangements affect nutrient accessibility and therefore influence levels

of competition and mutualism between microbial factors (Santegoeds and Damgaard, 1999). The opportunity to resolve the spatial distribution of active microbial cells will disclose the relevance of proximity for microbial interactions with each other, higher organisms or substrate surfaces. Relevant scaling of experimental designs can be achieved using geochemical profiling and flux analysis before and paralleling the use of community analysis. Retrieval of shotgun sequence information remains essential, for instance, when studying genomic variation and gene regulation within species, resulting in increased ecosystem functioning complexity (Darch *et al.*, 2015). Culture-based approaches, metaproteomics and metabolomics studies will effectively complement these studies. Finally, novel single-molecule sequencing methods, such as nanopore technologies (Oxford Nanopore Technologies (ONT; Schneider, 2012), may soon enable the direct measurement and sequencing of chemically modified bases incorporated into the nucleic acids of transcriptionally active or replicating cells. ONT has recently demonstrated the detection of 5-methylcytosine in bacterial and human DNA (Simpson *et al.*), highlighting its potential to detect other non-canonical bases in the near-term future (Woyke and Rubin, 2014).

Exciting technological advances have shaped the trajectory of microbial community studies, resulting in fascinating insights into our microbial world. We expect this trend to continue as sequence quality improves at reduced costs with a shift from bulk microbial community analyses to more refined studies that target active populations of interest. These improved and novel technologies will assist in exploring the surging scientific questions and applications that have been engaging the scientific community in the field of microbial ecology.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

This work was conducted by the US Department of Energy Joint Genome Institute, a DOE Office of Science User Facility, and was supported under contract no. DE-AC02-05CH11231. Research in the Wagner lab on single-cell activity assays was supported by an ERC Advanced Grant (NITRICARE, 294343), as well as by the Emerging Technologies Opportunity Program, which is supported by the Office of Science of the US Department of Energy under Contract DE-AC02-05CH11231.

References

- Abu Laban N, Tan B, Dao A, Foght J. (2015). Draft genome sequence of uncultivated *Desulfosporosinus* sp. Strain

- Tol-M, obtained by stable isotope probing using [^{13}C]6 toluene. *Gen Announc* 3: e01422–14.
- Alexander H, Jenkins BD, Rynearson TA, Dyhrman ST. (2015a). Metatranscriptome analyses indicate resource partitioning between diatoms in the field. *PNAS* 112: E2182–E2190.
- Alexander H, Rouco M, Haley ST, Wilson ST, Karl DM, Dyhrman ST. (2015b). Functional group-specific traits drive phytoplankton dynamics in the oligotrophic ocean. *PNAS* 112: E5972–E5979.
- Allison SD, Czimczik CI, Treseder KK. (2008). Microbial activity and soil respiration under nitrogen addition in Alaskan boreal forest. *Global Change Biol* 14: 1156–1168.
- Allison SD, Treseder KK. (2008). Warming and drying suppress microbial activity and carbon cycling in boreal forest soils. *Global Change Biol* 14: 2898–2909.
- Artursson V, Finlay RD, Jansson JK. (2005). Combined bromodeoxyuridine immunocapture and terminal-restriction fragment length polymorphism analysis highlights differences in the active soil bacterial metagenome due to *Glomus mosseae* inoculation or plant species. *Environmental Microbiology* 7: 1952–1966.
- Artursson V, Jansson JK. (2003). Use of Bromodeoxyuridine Immunocapture To Identify Active Bacteria Associated with Arbuscular Mycorrhizal Hyphae. *Appl Environ Microbiol* 69: 6208–6215.
- Ashkin A. (1970). Acceleration and trapping of particles by radiation pressure. *Physical review letters* 24: 156–159.
- Augsburger C, Karwautz C, Mußmann M, Daims H, Battin TJ. (2010). Drivers of bacterial colonization patterns in stream biofilms. *FEMS Microbiology Ecology* 72: 47–57.
- Baker BJ, Sheik CS, Taylor CA, Jain S, Bhasi A, Cavalcoli JD et al. (2013). Community transcriptomic assembly reveals microbes that contribute to deep-sea carbon and nitrogen cycling. *ISME J* 7: 1962–1973.
- Bello-Ortí B, Howell KJ, Tucker AW, Maskell DJ, Aragon V. (2015). Metatranscriptomics reveals metabolic adaptation and induction of virulence factors by *Haemophilus parasuis* during lung infection. *Vet Res*, 1–13.
- Berney M, Hammes F, Bosshard F, Weilenmann HU, Egli T. (2007). Assessment and interpretation of bacterial viability by using the live/dead bacLight kit in combination with flow cytometry. *Appl Environ Microbiol* 73: 3283–3290.
- Berry D, Mader E, Lee TK, Woebken D, Wang Y, Zhu D et al. (2015). Tracking heavy water (D $_2$ O) incorporation for identifying and sorting active microbial cells. *PNAS* 112: E194–E203.
- Bikel S, Valdez-Lara A, Cornejo-Granados F, Rico K, Canizales-Quinteros S, Soberón X et al. (2015). Combining metagenomics, metatranscriptomics and viromics to explore novel microbial interactions: towards a systems-level understanding of human microbiome. *CSBJ* 13: 390–401.
- Blazewicz SJ, Barnard RL, Daly RA, Firestone MK. (2013). Evaluating rRNA as an indicator of microbial activity in environmental communities: limitations and uses. *ISME J* 7: 2061–2068.
- Bomar L, Maltz M, Colston S, Graf J. (2011). Directed culturing of microorganisms using metatranscriptomics. *mBio* 2: e00012–11–e00012–11.
- Borneman J. (1999). Culture-independent identification of microorganisms that respond to specified stimuli. *Appl Environ Microbiol* 65: 3398–3400.
- Brown CT, Olm MR, Thomas BC, Banfield JF. (2016). In situ replication rates for uncultivated bacteria in microbial communities. *bioRxiv* e-pub ahead of print, doi:10.1101/057992.
- Buerger S, Spoering A, Gavrish E, Leslin C, Ling L, Epstein SS. (2012). Microbial scout hypothesis, stochastic exit from dormancy, and the nature of slow growers. *Appl Environ Microbiol* 78: 3221–3228.
- Burrow LC, Woebken D, Marshall IP, Lindquist EA, Bebout BM, Prufert-Bebout L et al. (2012). Anoxic carbon flux in photosynthetic microbial mats as revealed by metatranscriptomics. *ISME J* 7: 817–829.
- Carvalhais LC, Muzzi F, Tan C-H, Hsien-Choo J, Schenk PM. (2013). Plant growth in *Arabidopsis* is assisted by compost soil-derived microbial communities. *Front Plant Sci* 4: 235.
- Chaparro JM, Badri DV, Vivanco JM. (2013). Rhizosphere microbiome assemblage is affected by plant development. *ISME J* 8: 790–803.
- Chapelle E, Mendes R, Bakker PAH, Raaijmakers JM. (2015). Fungal invasion of the rhizosphere microbiome. *ISME J* 10: 265–268.
- Chemerys A, Pelletier E, Cruaud C, Martin F, Violet F, Jouanneau Y. (2014). Characterization of novel polycyclic aromatic hydrocarbon dioxygenases from the bacterial metagenomic DNA of a contaminated soil. *Appl Environ Microbiol* 80: 6591–6600.
- Chen Y, Dumont MG, Neufeld JD, Bodrossy L, Stralis-Pavese N, McNamara NP et al. (2008). Revealing the uncultivated majority: combining DNA stable-isotope probing, multiple displacement amplification and metagenomic analyses of uncultivated *Methylocystis* in acidic peatlands. *Environ Microbiol* 10: 2609–2622.
- Damon C, Vallon L, Zimmermann S, Haider MZ, Galeote V, Dequin S et al. (2011). A novel fungal family of oligopeptide transporters identified by functional metatranscriptomics of soil eukaryotes. *ISME J* 5: 1871–1880.
- Darch SE, McNally A, Harrison F, Corander J, Barr HL, Paszkiewicz K et al. (2015). Recombination is a key driver of genomic and phenotypic diversity in a *Pseudomonas aeruginosa* population during cystic fibrosis infection. *Scientific Reports* 5: 7649.
- David MM, Cecillon S, Warne BM, Prestat E, Jansson JK, Vogel TM. (2014). Microbial ecology of chlorinated solvent biodegradation. *Environ Microbiol* 17: 4835–4850.
- De Filippis F, Genovese A, Ferranti P, Gilbert JA, Ercolini D. (2016). Metatranscriptomics reveals temperature-driven functional changes in microbiome impacting cheese maturation rate. *Sci Rep*, 1–11.
- Dieterich DC, Link AJ, Graumann J, Tirrell DA, Schuman EM. (2006). Selective identification of newly synthesized proteins in mammalian cells using bioorthogonal noncanonical amino acid tagging (BONCAT). *Proceedings of the National Academy of Sciences* 103: 9482–9487.
- Dumont MG, Pommerenke B, Casper P. (2013). Using stable isotope probing to obtain a targeted metatranscriptome of aerobic methanotrophs in lake sediment. *Environmental Microbiology Reports* 5: 757–764.
- Dumont MG, Radajewski SM, Miguez CB, McDonald IR, Murrell JC. (2006). Identification of a complete methane monooxygenase operon from soil by combining stable isotope probing and metagenomic analysis. *Environ Microbiol* 8: 1240–1250.

- Edlund A, Hårdeman F, Jansson JK, Sjöling S. (2008). Active bacterial community structure along vertical redox gradients in Baltic Sea sediment. *Environmental Microbiology* **10**: 2051–2063.
- Eichorst SA, Strasser F, Woyke T, Schintlmeister A, Wagner M, Woebken D. (2015). Advancements in the application of NanoSIMS and Raman microspectroscopy to investigate the activity of microbial cells in soils Häggblom M (ed). *FEMS Microbiology Ecology* **91**: 1–14.
- Eickmeier O, Hector A, Singh A, Chotirmall SH, Hartl D. (2014). Fungi in cystic fibrosis: recent findings and unresolved questions. *Curr Fungal Infect Rep* **9**: 1–5.
- Eloe-Fadrosh EA, Ivanova NN, Woyke T, Kyrpides NC. (2016). Metagenomics uncovers gaps in amplicon-based detection of microbial diversity. *Nature Microbiology* **1**: 15032.
- Embree M, Nagarajan H, Movahedi N, Chitsaz H, Zengler K. (2013). Single-cell genome and metatranscriptome sequencing reveal metabolic interactions of an alkane-degrading methanogenic community. *ISME J* **8**: 757–767.
- Epstein SS. (2013). ScienceDirectThe phenomenon of microbial uncultivability. *Current Opinion in Microbiology* **16**: 636–642.
- Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML et al. (2015). Anvi'o: an advanced analysis and visualization platform for 'omics data. *PeerJ* **3**: e1319.
- Eyice OZ, Namura M, Chen Y, Mead A, Samavedam S. (2015). SIP metagenomics identifies uncultivated Methylophilaceae as dimethylsulphide degrading bacteria in soil and lake sediment. *ISME J* **9**: 2336–2348.
- Feike J, Jürgens K, Hollibaugh JT, Krüger S, Jost G, Labrenz M. (2011). Measuring unbiased metatranscriptomics in suboxic waters of the central Baltic Sea using a new in situ fixation system. *ISME J* **6**: 461–470.
- Fortunato CS, Huber JA. (2015). Coupled RNA-SIP and metatranscriptomics of active chemolithoautotrophic communities at a deep-sea hydrothermal vent. *ISME J* **10**: 1925–1938.
- Franklin RB, Luria C, Shozo Ozaki L, Bukaveckas PA. (2013). Community composition and activity state of estuarine bacterioplankton assessed using differential staining and metagenomic analysis of 16S rDNA and rRNA. *Aquat Microb Ecol* **69**: 247–261.
- Franzosa EA, Morgan XC, Segata N, Waldron L, Reyes J, Earl AM et al. (2014). Relating the metatranscriptome and metagenome of the human gut. *PNAS* **111**: E2329–E2338.
- Frias-Lopez J, Shi Y, Tyson GW, Coleman ML, Schuster SC, Chisholm SW et al. (2008). Microbial community gene expression in ocean surface waters. *PNAS* **105**: 3805–3810.
- Gibson TE, Bashan A, Cao H-T, Weiss ST, Liu Y-Y. (2016). On the Origins and Control of Community Types in the Human Microbiome. In: Gore J (ed). *PLoS Comput Biol* **12**: e1004688.
- Gieseke A, Bjerrum L, Wagner M, Amann R. (2003). Structure and activity of multiple nitrifying bacterial populations co-existing in a biofilm. *Environmental Microbiology* **5**: 355–369.
- Gifford SM, Sharma S, Rinta-Kanto JM, Moran MA. (2010). Quantitative analysis of a deeply sequenced marine microbial metatranscriptome. *The ISME Journal* **5**: 461–472.
- Goldfarb KC, Karaoz U, Hanson CA, Santee CA, Bradford MA, Treseder KK et al. (2011). Differential growth responses of soil bacterial taxa to carbon substrates of varying chemical recalcitrance. *Frontiers in Microbiology* **2**: 94.
- Gruber-Vodicka HR, Dirks U, Leisch N, Baranyi C, Stoecker K, Bulgheresi S et al. (2011). Paracatenula, an ancient symbiosis between thiotrophic Alphaproteobacteria and catenulid flatworms. *Proceedings of the National Academy of Sciences* **108**: 12078–12083.
- Gutierrez T, Singleton DR, Berry D, Yang T, Aitken MD, Teske A. (2013). Hydrocarbon-degrading bacteria enriched by the Deepwater Horizon oil spill identified by cultivation and DNA-SIP. *ISME J* **7**: 2091–2104.
- Hamasaki K, Taniguchi A, Tada Y, Kaneko R, Miki T. (2016). Active populations of rare microbes in oceanic environments as revealed by bromodeoxyuridine incorporation and 454 tag sequencing. *Gene* **576**: 650–656.
- Handelsman J. (2004). Metagenomics: application of genomics to uncultured microorganisms. *Microbiology and Molecular Biology Reviews* **68**: 669–685.
- Hanson CA, Allison SD, Bradford MA, Wallenstein MD, Treseder KK. (2008). Fungal Taxa Target Different Carbon Sources in Forest Soil. *Ecosystems* **11**: 1157–1167.
- Haroon MF, Hu S, Shi Y, Imelfort M, Keller J, Hugenholtz P et al. (2013). Anaerobic oxidation of methane coupled to nitrate reduction in a novel archaeal lineage. *Nature*, 1–7.
- Hatzenpichler R, Connon SA, Goudeau D, Malmstrom RR, Woyke T, Orphan VJ. (2016). Visualizing in situ translational activity for identifying and sorting slow-growing archaeal–bacterial consortia. *PNAS* **113**: E4069–E4078.
- Hatzenpichler R, Scheller S, Tavormina PL, Babin BM, Tirrell DA, Orphan VJ. (2014). In situ visualization of newly synthesized proteins in environmental microbes using amino acid tagging and click chemistry. *Environ Microbiol* **16**: 2568–2590.
- He S, Wurtzel O, Singh K, Froula JL, Yilmaz S, Tringe SG et al. (2010). Validation of two ribosomal RNA removal methods for microbial metatranscriptomics. *Nat Methods* **7**: 807–812.
- Helbling DE, Ackermann M, Fenner K, Kohler H-PE, Johnson DR. (2011). The activity level of a microbial community function can be predicted from its metatranscriptome. *ISME J* **6**: 902–904.
- Hellman M, Berg J, Brandt KK, Hallin S. (2011). Journal of Microbiological Methods. *Journal of microbiological methods* **86**: 376–378.
- Hewson I, Poretsky RS, Dyhrman ST, Zielinski B, White AE, Tripp HJ et al. (2009). Microbial community gene expression within colonies of the diazotroph, *Trichodesmium*, from the Southwest Pacific Ocean. *The ISME Journal* **3**: 1286–1300.
- Hilton JA, Satinsky BM, Doherty M, Zielinski B, Zehr JP. (2014). Metatranscriptomics of N₂-fixing cyanobacteria in the Amazon River plume. *ISME J* **9**: 1557–1569.
- Hoehler TM, Jørgensen BB. (2013). Microbial life under extreme energy limitation. *Nat Rev Micro* **11**: 83–94.
- Holmes DE, Giloteaux L, Williams KH, Wrighton KC, Wilkins MJ, Thompson CA et al. (2013). Enrichment of specific protozoan populations during in situ bioremediation of uranium-contaminated groundwater. *The ISME Journal* **7**: 1286–1298.

- Hua Z-S, Han Y-J, Chen L-X, Liu J, Hu M, Li S-J *et al.* (2014). Ecological roles of dominant and rare prokaryotes in acid mine drainage revealed by metagenomics and metatranscriptomics. *ISME J* **9**: 1280–1294.
- Huang WE, Ferguson A, Singer AC, Lawson K, Thompson IP, Kalin RM *et al.* (2008). Resolving genetic functions within microbial populations: in situ analyses using rRNA and mrna stable isotope probing coupled with single-cell Raman-fluorescence in situ hybridization. *Appl Environ Microbiol* **75**: 234–241.
- Huang WE, Stoecker K, Griffiths R, Newbold L, Daims H, Whiteley AS *et al.* (2007). Raman-FISH: combining stable-isotope Raman spectroscopy and fluorescence in situ hybridization for the single cell analysis of identity and function. *Environmental Microbiology* **9**: 1878–1889.
- Huang WE, Ward AD, Whiteley AS. (2009). Raman tweezers sorting of single microbial cells. *Environmental Microbiology Reports* **1**: 44–49.
- Hutchens E, Radajewski S, Dumont MG, McDonald IR, Murrell JC. (2004). Analysis of methanotrophic bacteria in Movile Cave by stable isotope probing. *Environmental Microbiology* **6**: 111–120.
- Ishii S, Suzuki S, Tenney A, Norden-Krichmar TM, Neelson KH, Bretschger O. (2015). Microbial metabolic networks in a complex electrogenic bio lm recovered from a stimulus-induced metatranscriptomics approach. *Sci Rep*, 1–14.
- Jehmlich N, Schmidt F, Taubert M, Seifert J, Bastida F, Bergen Von M *et al.* (2010). Protein-based stable isotope probing. *Nat Protoc* **5**: 1957–1966.
- Jiang Y, Xiong X, Danska J, Parkinson J. (2016). Metatranscriptomic analysis of diverse microbial communities reveals core metabolic pathways and microbiome-specific functionality. *Microbiome*, 1–18.
- Jones DS, Flood BE, Bailey JV. (2015). Metatranscriptomic insights into polyphosphate metabolism in marine sediments. *ISME J* **10**: 1015–1019.
- Jorgensen BB. (2011). Deep subseafloor microbial cells on physiological standby. *PNAS* **108**: 18193–18194.
- Jorth P, Turner KH, Gumus P, Nizam N, Buduneli N, Whiteley M. (2014). Metatranscriptomics of the human oral microbiome during health and disease. *mBio* **5**: e01012–14–e01012–14.
- Kalyuzhnaya MG, Hristova KR, Lidstrom ME, Chistoserdova L. (2008a). characterization of a novel methanol dehydrogenase in representatives of burkholderiales: implications for environmental detection of methylotrophy and evidence for convergent evolution. *J Bacteriol* **190**: 3817–3823.
- Kalyuzhnaya MG, Lapidus A, Ivanova N, Copeland AC, Mchardy AC, Szeto E *et al.* (2008b). High-resolution metagenomics targets specific functional types in complex microbial communities. *Nat Biotechnol* **26**: 1029–1034.
- Kalyuzhnaya MG, Lidstrom ME, Chistoserdova L. (2008). Real-time detection of actively metabolizing microbes by redox sensing as applied to methylotroph populations in Lake Washington. *The ISME Journal* **2**: 696–706.
- Kalyuzhnaya MG, Lidstrom ME, Chistoserdova L. (2008c). Real-time detection of actively metabolizing microbes by redox sensing as applied to methylotroph populations in Lake Washington. *ISME J* **2**: 696–706.
- Kim SJ, Park S-J, Jung MY, Kim JG, Madsen EL, Rhee SK. (2014). An uncultivated nitrate-reducing member of the genus herminiimonas degrades toluene. *Appl Environ Microbiol* **80**: 3233–3243.
- Klindworth A, Mann AJ, Huang S, Wichels A, Quast C, Waldmann J *et al.* (2014). Diversity and activity of marine bacterioplankton during a diatom bloom in the North Sea assessed by total RNA and pyrotag sequencing. *Mar Genom* **18**: 185–192.
- Kodzias R, Gojobori T. (2016). Single-cell technologies in environmental omics. *Gene* **576**: 701–707.
- Konopka MC, Strovas TJ, Ojala DS, Chistoserdova L, Lidstrom ME, Kalyuzhnaya MG. (2010). Respiration response imaging for real-time detection of microbial function at the single-cell level. *Appl Environ Microbiol* **77**: 67–72.
- Kopf SH, McGlynn SE, Green-Saxena A, Guan Y, Newman DK, Orphan VJ. (2015). Heavy water and 15 N labelling with NanoSIMS analysis reveals growth rate-dependent metabolic heterogeneity in chemostats. *Environmental Microbiology* **17**: 2542–2556.
- Korem T, Zeevi D, Suez J, Weinberger A, Avnit-Sagi T, Pompan-Lotan M *et al.* (2015). Growth dynamics of gut microbiota in health and disease inferred from single metagenomic samples. *Science* **349**: 1101–1106.
- Lamendella R, Strutt S, Borglin S, Chakraborty R, Tas N, Mason OU *et al.* (2014). Assessment of the Deepwater Horizon oil spill impact on Gulf coast microbial communities. *Front Microbiol* **5**. e-pub ahead of print, doi: 10.3389/fmicb.2014.00130/abstract.
- LaRowe DE, Amend JP. (2016). The energetics of ano-bolism in natural settings. *ISME J*. **10**: 1285–1295.
- Lasken RS. (2007). Single-cell genomic sequencing using Multiple Displacement Amplification. *Current Opinion in Microbiology* **10**: 510–516.
- Lehembre F, Doillon D, David E, Perrotto S, Baude J, Foulon J *et al.* (2013). Soil metatranscriptomics for mining eukaryotic heavy metal resistance genes. *Environ Microbiol* **15**: 2829–2840.
- Leininger S, Urich T, Schlöter M, Schwark L, Qi J, Nicol GW *et al.* (2006). Archaea predominate among ammonia-oxidizing prokaryotes in soils. *Nature* **442**: 806–809.
- Lennon JT, Jones SE. (2011). Microbial seed banks: the ecological and evolutionary implications of dormancy. *Nat Rev Micro* **9**: 119–130.
- Lesniewski RA, Jain S, Anantharaman K, Schloss PD, Dick GJ. (2012). The metatranscriptome of a deep-sea hydrothermal plume is dominated by water column methanotrophs and lithotrophs. *The ISME Journal* **6**: 2257–2268.
- Lester W, Sun SH, Seber A. (1960). Observations on the influence of deuterium on bacterial growth. *Annals of the New York Academy of Sciences* **84**: 667–677.
- Lim YW, Schmieder R, Haynes M, Willner D, Furlan M, Youle M *et al.* (2013). Metagenomics and metatranscriptomics: windows on CF-associated viral and microbial communities. *J Cyst Fibros* **12**: 154–164.
- Lin S, Zhang H, Zhuang Y, Tran B. (2010). Spliced leader-based metatranscriptomic analyses lead to recognition of hidden genomic features in dinoflagellates. *Proc Natl Acad Sci* **107**: 20033–20038.
- Lloyd KG, May MK, Kevorkian RT, Steen AD. (2013). Meta-analysis of quantification methods shows that archaea and bacteria have similar abundances in the subseafloor. *Appl Environ Microbiol* **79**: 7790–7799.
- Majed N, Chernenko T, Diem M, Gu AZ. (2012). Identification of functionally relevant populations in enhanced biological phosphorus removal processes based on intracellular polymers profiles and insights into the

- metabolic diversity and heterogeneity. *Environmental Science & Technology* **46**: 5010–5017.
- Manefield M, Whiteley AS, Griffiths RI, Bailey MJ. (2002). RNA Stable Isotope Probing, a Novel Means of Linking Microbial Community Function to Phylogeny. *Appl Environ Microbiol* **68**: 5367–5373.
- Marzano S-YL, Domier LL. (2016). Novel mycoviruses discovered from metatranscriptomics survey of soybean phyllosphere phytobiomes. *Virus Res* **213**: 332–342.
- McGuire KL, Bent E, Borneman J, Majumder A, Allison SD, Treseder KK. (2012). Functional diversity in resource use by fungi. *Ecology* **91**: 2324–2332.
- Milucka J, Ferdelman TG, Polerecky L, Franzke D, Wegener G, Schmid M et al. (2012). Zero-valent sulphur is a key intermediate in marine methane oxidation. *Nature* **491**: 541–546.
- Mondot S, Lepage P. (2016). The human gut microbiome and its dysfunctions through the meta-omics prism. *Ann N Y Acad Sci*, 1–11; <http://doi.org/10.1111/nyas.13033>.
- Moran MA, Satinsky B, Gifford SM, Luo H, Rivers A, Chan L-K et al. (2012). Sizing up metatranscriptomics. *ISME J* **7**: 237–243.
- Mott ML, Berger JM. (2007). DNA replication initiation: mechanisms and regulation in bacteria. *Nat Rev Micro* **5**: 343–354.
- Mou X, Sun S, Edwards RA, Hodson RE, Moran MA. (2008). Bacterial carbon processing by generalist species in the coastal ocean. *Nature* **451**: 708–711.
- Narasingarao P, Podell S, Ugalde JA, Brochier-Armanet CEL, Emerson JB, Brocks JJ et al. (2011). De novo metagenomic assembly reveals abundant novel major lineage of Archaea in hypersaline microbial communities. *The ISME Journal* **6**: 81–93.
- Neufeld JD, Chen Y, Dumont MG, Murrell JC. (2008). Marine methylotrophs revealed by stable-isotope probing, multiple displacement amplification and metagenomics. *Environ Microbiol* **10**: 1526–1535.
- Neufeld JD, Dumont MG, Vohra J, Murrell JC. (2006). Methodological Considerations for the Use of Stable Isotope Probing in Microbial Ecology. *Microb Ecol* **53**: 435–442.
- Neufeld JD, Vohra J, Dumont MG, Lueders T, Manefield M, Friedrich MW et al. (2007). DNA stable-isotope probing. *Nat Protoc* **2**: 860–866.
- Orsi WD, Edgcomb VP, Christman GD, Biddle JF. (2013). Gene expression in the deep biosphere. *Nature* **499**: 205–208.
- Ottman N, Smidt H, De Vos WM, Belzer C. (2012). The function of our microbiota: who is out there and what do they do? *Front Cell Inf Microbiol* **2**: 104.
- Paerl HW, Steppe TF. (2003). Scaling up: the next challenge in environmental microbiology. *Environmental Microbiology* **5**: 1025–1038.
- Pagani I, Liolios K, Jansson J, Chen I-MA, Smirnova T, Nosrat B et al. (2011). The Genomes OnLine Database (GOLD) v.4: status of genomic and metagenomic projects and their associated metadata. *Nucleic Acids Research* **40**: D571–D579.
- Paine RT. (1995). A conversation on refining the concept of keystone species.
- Pester M, Bittner N, Deevong P, Wagner M, Loy A. (2010). A 'rare biosphere' microorganism contributes to sulfate reduction in a peatland. *The ISME Journal*, 1–12.
- Pinnell LJ, Dunford E, Ronan P, Hausner M, Neufeld JD. (2014). Recovering glycoside hydrolase genes from active tundra cellulolytic bacteria. *Can J Microbiol* **60**: 469–476.
- Poretsky RS, Sun S, Mou X, Moran MA. (2010). Transporter genes expressed by coastal bacterioplankton in response to dissolved organic carbon. *Environmental Microbiology* **12**: 616–627.
- Radajewski S, Ineson P, Parekh NR, Murrell JC. (2000). Stable-isotope probing as a tool in microbial ecology. *Nature* **403**: 646–649.
- Radajewski S, McDonald IR, Murrell JC. (2003). Stable-isotope probing of nucleic acids: a window to the function of uncultured microorganisms. *Current Opinion in Biotechnology* **14**: 296–302.
- Ruff SE, Biddle JF, Teske AP, Knittel K, Boetius A, Ramette A. (2015). Global dispersion and local diversification of the methane seep microbiome. *PNAS* **112**: 4015–4020 201421865.
- Sanders JG, Beinart RA, Stewart FJ, DeLong EF, Girguis PR. (2013). Metatranscriptomics reveal differences in in situ energy and nitrogen metabolism among hydrothermal vent snail symbionts. *ISME J* **7**: 1556–1567.
- Santegoeds CM, Damgaard LR. (1999). Distribution of sulfate-reducing and methanogenic bacteria in anaerobic aggregates determined by microsensor and molecular analyses. *Appl Environ Microbiol* **65**: 4618–4629.
- Satinsky BM, Crump BC, Smith CB, Sharma S, Zielinski BL, Doherty M et al. (2014). Microspatial gene expression patterns in the Amazon River Plume. *PNAS* **111**: 11085–11090.
- Schneider G. F, Dekker C. (2012). *News and Views*. Nature Publishing Group **30**(4): 326–328; <http://doi.org/10.1038/nbt.2181>.
- Schwab C, Tveit AT, Schleper C, Urich T. (2014). Gene expression of lactobacilli in murine forestomach biofilms. *Microbial Biotechnol* **7**: 347–359.
- Schwartz E, Hayer M, Hungate BA, Koch BJ, McHugh TA, Mercurio W et al. (2016). Stable isotope probing with 18O-water to investigate microbial growth and death in environmental samples. *Curr Opin Biotechnol* **41**: 14–18.
- Shi Y, Tyson GW, Delong EF. (2009). Metatranscriptomics reveals unique microbial small RNAs in the ocean's water column. *Nature* **459**: 266–269.
- Simberloff D, Dayan T. (1991). The guild concept and the structure of ecological communities. *Annu Rev Ecol Syst* **22**: 115–143.
- Simpson JT, Workman R, Zuzarte PC, David M, Dursi LJ, Timp W Detecting DNA Methylation using the Oxford Nanopore Technologies MinION sequencer. e-pub ahead of print10.1101/047142.
- Simón-Soro A, Guillen-Navarro M, Mira A. (2014). Metatranscriptomics reveals overall active bacterial composition in caries lesions. *J Oral Microbiol* **6**: 1252.
- Song Y, Yin H, Huang WE. (2016). ScienceDirectRaman activated cell sorting. *Current opinion in chemical biology* **33**: 1–8.
- Steen IH, Dahle H, Stokke R, Roalkvam I, Daae F-L, Rapp HT et al. (2016). Novel Barite Chimneys at the Loki's Castle Vent Field Shed Light on Key Factors Shaping Microbial Communities and Functions in Hydrothermal Systems. *Fron Microbiol* **6**: 4248.
- Sul WJ, Park J, Quensen JF, Rodrigues JLM, Seliger L, Tsoi TV et al. (2009). DNA-stable isotope probing integrated with metagenomics for retrieval of biphenyl dioxygenase genes from polychlorinated biphenyl-contaminated river sediment. *Appl Environ Microbiol* **75**: 5501–5506.
- Taylor DL, Hollingsworth TN, McFarland JW. (2014). A first comprehensive census of fungi in soil reveals both hyperdiversity and fine-scale niche partitioning. *Ecological Monographs* **84**: 3–20.

- Teske A, Sørensen KB. (2007). Uncultured archaea in deep marine subsurface sediments: have we caught them all? *The ISME Journal* **2**: 3–18.
- Tringe SG, Hugenholtz P. (2008). A renaissance for the pioneering 16 S rRNA gene. *Current Opinion in Microbiology* **11**: 442–446.
- Ullrich S, Karrasch B, Hoppe H, Jeskulke K, Mehrens M. (1996). Toxic effects on bacterial metabolism of the redox dye 5-cyano-2,3-ditolyl tetrazolium chloride. *Appl Environ Microbiol* **62**: 4587–4593.
- Urbach E, Vergin KL, Giovannoni SJ. (1999). Immunochemical detection and isolation of DNA from metabolically active bacteria. *Appl Environ Microbiol* **65**: 1207–1213.
- Urich T, Lanzén A, Qi J, Huson DH, Schleper C, Schuster SC. (2008). Simultaneous assessment of soil microbial community structure and function through analysis of the meta-transcriptome Ward N (ed). *PLoS One* **3**: e2527.
- Ursell LK, Knight R. (2013). Xenobiotics and the human gut microbiome: metatranscriptomics reveal the active players. *Cell Metab* **17**: 317–318.
- Verastegui Y, Cheng J, Engel K, Kolczynski D, Mortimer S, Lavigne J et al. (2014). Multisubstrate isotope labeling and metagenomic analysis of active soil bacterial communities. *mBio* **5**: e01157–14–e01157–14.
- Vila-Costa M, Sharma S, Moran MA, Casamayor EO. (2012). Diel gene expression profiles of a phosphorus limited mountain lake using metatranscriptomics. *Environ Microbiol* **15**: 1190–1203.
- Wang T-H, Peng Y, Zhang C, Wong PK, Ho C-M. (2005). Single-molecule tracing on a fluidic microchip for quantitative detection of low-abundance nucleic acids. *Journal of the American Chemical Society* **127**: 5354–5359.
- Wang Y, Huang WE, Cui L, Wagner M. (2016). Single cell stable isotope probing in microbiology using Raman microspectroscopy. *Current Opinion in Biotechnology* **41**: 34–42.
- Wang Y, Ji Y, Wharfe ES, Meadows RS, March P, Goodacre R et al. (2013). Raman Activated Cell Ejection for Isolation of Single Cells. *Anal Chem* **85**: 10697–10701.
- Werner JJ, Garcia ML, Perkins SD, Yarasheski KE, Smith SR, Muegge BD et al. (2014). Microbial community dynamics and stability during an ammonia-induced shift to syntrophic acetate oxidation. *Appl Environ Microbiol* **80**: 3375–3383.
- Whiteley AS, Thomson B, Lueders T, Manefield M. (2007). RNA stable-isotope probing. *Nat Protoc* **2**: 838–844.
- Woyke T, Rubin EM. (2014). Searching for new branches on the tree of life. *Science* **346**: 698–699.
- Xie C, Dinno MA, Li Y-Q. (2002). Near-infrared Raman spectroscopy of single optically trapped biological cells. *Opt Lett* **27**: 249–251.
- Xie C, Li Y-Q. (2003). Confocal micro-Raman spectroscopy of single biological cells using optical trapping and shifted excitation difference techniques. *J Appl Phys* **93**: 2982.
- Yin B, Crowley D, Sparovek G, De Melo WJ, Borneman J. (2000). Bacterial functional redundancy along a soil reclamation gradient. *Appl Environ Microbiol* **66**: 4361–4365.
- Zhang P, Ren L, Zhang X, Shan Y, Wang Y. (2015a). Raman-activated cell sorting based on dielectrophoretic single-cell trap and release. *Anal Chem* **87**: 2282–2289.
- Zhang Q, Zhang P, Gou H, Mou C, Huang WE, Yang M et al. (2015b). Towards high-throughput micro-fluidic Raman-activated cell sorting. *Analyst* **140**: 6163–6174.
- Zhang T, Tremblay PL, Chaurasia AK, Smith JA, Bain TS, Lovley DR. (2013). Anaerobic benzene oxidation via phenol in *Geobacter metallireducens*. *Appl Environ Microbiol* **79**: 7800–7806.

This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>

© The Author(s) 2017