A New Cadinane Sesquiterpene with Significant Anti-HIV-1 Activity from the Cultures of the Basidiomycete *Tyromyces chioneus*

Dong-Ze Liu, Fei Wang, Liu-Meng Yang, Yong-Tang Zheng, Ji-Kai Liu

Received: February 12, 2007 / Accepted: May 16, 2007

© Japan Antibiotics Research Association

Abstract A new cadinane sesquiterpene (4 β ,14-dihydroxy-6 α ,7 β H-1(10)-cadinene, **1**) was isolated from the cultures of the basidiomycete, *Tyromyces chioneus*. Its structure was established on the basis of spectral measurements (MS, IR, 1D and 2D NMR experiments). **1** showed significant anti-HIV-1 activity with EC₅₀=3.0 μ g/ml (SI=25.4).

Keywords *Tyromyces chioneus*, culture broth, cadinane sesquiterpene, anti-HIV-1

of secondary metabolites. *Tyromyces chioneus* belonging to order Polyporales of the family Polyporaceae is an inedible fungi and mainly occurs on wood as a substrate. So far, little work has been done on the chemical constituents of *T. chioneus*. As one part of our search for naturally occurring bioactive metabolites from the higher fungi in Yunnan province [1~6], a new cadinane sesquiterpene (**1**) was isolated from the cultures of the basidiomycete *T. chioneus*. In this paper, we describe the isolation and structure elucidation of **1** as well as its anti-HIV-1 activity.

Introduction

Yunnan Province, southwest of China, is one of the areas with the richest and most diverse bioresources in the world. Among these bioresources, fungi produce a broad variety

Materials and Methods

General

Optical rotation was measured on an Horiba SEPA-300 polarimeter. IR spectra were obtained on a Bruker Tensor 27 instrument as KBr pellets. NMR spectra were recorded on Bruker AV-400 and Bruker DRX-500 spectrometers in CD₃OD and CDCl₃ solvents with TMS as an internal standard. EI-MS was recorded with a VG Autospec-3000 spectrometer. HRESI-MS was recorded with an API QSTAR Pulsar 1 spectrometer.

Silica gel (200~300 mesh, Qingdao Marine Chemical Inc., China) and Sephadex LH-20 (Amersham Biosciences, Sweden) were used for column chromatography. Fractions were monitored by TLC and spots were visualized by heating silica gel plates sprayed with 10% H₂SO₄ in ethanol.

4 β ,14-dihydroxy-6 α ,7 β H-1(10)-cadinene(**1**) 1(10),4-cadinadiene(**2**)

Fig. 1 Structures of **1** and **2**, and key HMBC correlations of **1**.

J.-K. Liu (Corresponding author), **D.-Z. Liu**, **F. Wang**: State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, China, E-mail: jkliu@mail.kib.ac.cn

D.-Z. Liu, **F. Wang**: Graduate University of Chinese Academy of Sciences, Beijing 100049, China

L.-M. Yang, **Y.-T. Zheng**: Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650223, China

Fungal Material

The basidiomycete *T. chioneus* was collected at Ailao Mountain of Yunnan Province, China, in July 2003 and identified by Prof. Mu Zang, Kunming Institute of Botany, Chinese Academy of Sciences (CAS). The voucher specimen was deposited at the Herbarium of Kunming Institute of Botany, CAS.

Fermentation and Isolation

The culture medium consisted of potato (peeled) 200 g, glucose 20 g, KH_2PO_4 3 g, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 1.5 g, citric acid 0.1 g and thiamine hydrochloride 10 mg in 1 liter of deionized water. Reagent bottles were used as shaker flasks (size: 500 ml; volume of media: 300 ml). The pH was adjusted to 6.5 before autoclaving. Fermentation was carried out on a shaker at 22°C and 150 rpm for 10 days.

The whole culture broth of *T. chioneus* (12 liters) was filtered and then extracted twice with EtOAc. The organic layer was concentrated *in vacuo* to give an oily residue (1.3 g) that was applied to a silica gel column, which was eluted stepwise with a CHCl_3 -MeOH solvent system. Fr. III (19 mg), eluted with CHCl_3 -MeOH (100:1, v/v), was further purified by repeated chromatography on a Sephadex LH-20 column eluted with CHCl_3 -MeOH (1:1, v/v) to yield **1** (8.5 mg).

Physico-chemical Properties

4 β ,14-dihydroxy-6 α ,7 β H-1(10)-cadinene (**1**). Colorless Oil; $[\alpha]_D^{29.8} -20.3$ (*c* 0.8, MeOH); IR ν_{max} (KBr) cm^{-1} : 3440, 2959, 2927, 2870, 1630, 1063; HRESI-MS (+), *m/z* 261.1834 $[\text{M}+\text{Na}]^+$ (calcd. for $\text{C}_{15}\text{H}_{26}\text{O}_2\text{Na}$, 261.1830); EI-MS (70 eV) *m/z* (rel. int.): 220 (12) $[\text{M}-\text{H}_2\text{O}]^+$, 202 (13), 189 (13), 177 (100), 159 (45), 147 (27), 133 (25), 119 (40); ^1H -NMR (CD_3OD) and ^{13}C -NMR (CD_3OD and CDCl_3): see Table 1.

Reagents and Cell Culture

AZT (=3'-azido-3'-deoxythymidine) and MTT (=3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) were purchased from Sigma. Cells were donated by the Medical Research Council (MRC), AIDS Reagent Project, UK, and grown in RPMI-1640 medium supplemented with 10% heat-inactivated fetal calf serum (Gibco), 2 mM L-glutamine, 10 mM HEPES (=4-(2-hydroxyethyl) piperazine-1-ethanesulfonic acid), 50 μM 2-mercaptoethanol, 100,000 IU/ml penicillin, and 100 $\mu\text{g}/\text{ml}$ streptomycin sulfate. All cells and virus were stored and resuscitated by common methods.

Cytotoxicity of Compound **1** on C8166 Cells

C8166 was one of the host cells for HIV-1. On a microtiter

plate, 100 μl of $4 \times 10^5/\text{ml}$ cells were seeded. Then 100 μl of various concentrations of **1** was added and the cells were incubated at 37°C in a humidified atmosphere of 5% CO_2 for 72 hours. The cellular toxicity was assessed by an MTT colorimetric assay, the plates were read with a Bio-Tek ELx 800 ELISA reader at 595/630 nm, and the 50% inhibitory concentration (IC_{50}) was calculated.

Cytopathic-Effect (CPE) Inhibition Assay of **1**

In the presence of 100 μl of various concentrations of **1**, C8166 cells ($4 \times 10^5/\text{ml}$) were infected with HIV-1_{IIIB} at a multiplicity of infection (M.O.I) of 0.06. The final volume was 200 μl . The plates were incubated in a humidified incubator at 37°C and 5% CO_2 . AZT (Sigma Chemical Co.) was used for drug control. After 3 days of culture, the cytopathic effect was measured by counting the number of syncytia (multinucleated giant cells) in each well under an inverted microscope, and the 50% effective concentration (EC_{50}) was calculated.

Results and Discussion

1 was obtained as an oil. Based on positive ion HRESI-MS ($[\text{M}+\text{Na}]^+$ *m/z* 261.1834, calcd. for 261.1830) and NMR spectra data, the molecular formula was determined to be $\text{C}_{15}\text{H}_{26}\text{O}_2$, corresponding to 3 degrees of unsaturation. The IR spectrum indicated the presence of hydroxyl group at 3440 cm^{-1} and C=C double bond at 1630 cm^{-1} . The ^1H -NMR spectrum (Table 1) exhibited one methylene proton bearing a hydroxyl group [δ_{H} 3.98 (1H, d, $J=11.6 \text{ Hz}$), 4.11 (1H, d, $J=11.6 \text{ Hz}$) and three methyl signals due to one tertiary methyl group [δ_{H} 1.31 (3H, s)] and two secondary methyl groups [δ_{H} 0.97 (3H, d, $J=6.8 \text{ Hz}$), 0.81 (3H, d, $J=6.8 \text{ Hz}$)].

The ^{13}C -NMR and DEPT spectra (Table 1, in CD_3OD) displayed 15 carbon resonances, including three methine carbons (δ_{C} 47.7, 39.7, 28.7), six methylene carbons (δ_{C} 62.4, 48.3, 41.6, 28.7, 27.4, 22.4), two olefinic carbons (δ_{C} 137.0, 130.3), one quaternary carbon (δ_{C} 71.3) and three methyl carbons (δ_{C} 25.6, 22.1, 17.1). These observations, in combination with the molecular formula, indicated the presence of two OH groups and two rings. The similarity of ^{13}C -NMR data of partial structure for **1** [C-1 (δ_{C} 129.5), C-2 (δ_{C} 26.6), C-6 (δ_{C} 38.5), C-7 (δ_{C} 46.0), C-8 (δ_{C} 21.7), C-12 (δ_{C} 16.7), and C-13 (δ_{C} 21.3)] with those of 1(10),4-cadinadiene (**2**) [7] [C-1 (δ_{C} 129.9), C-2 (δ_{C} 26.7), C-6 (δ_{C} 39.5), C-7 (δ_{C} 45.4), C-8 (δ_{C} 21.2), C-12 (δ_{C} 15.6), and C-13 (δ_{C} 21.7)] suggested that **1** might be a cadinane-type sesquiterpene. This hypothesis was confirmed by careful analysis of ^1H - ^1H

Table 1 NMR spectral data for **1** and **2**. (δ in ppm, J in Hz)

Position	1 (CD ₃ OD)		1 (CDCl ₃)	2 (CDCl ₃)
	δ (H)	δ (C)	δ (C)	δ (C)
1	—	130.3 (s)	129.5	129.9
2	2.78 (1H, ddd, 14.0, 3.2, 2.4) 1.78 (1H, m)	27.4 (t)	26.6	26.7
3	1.41 (1H, m), 1.71 (1H, m)	41.6 (t)	41.0	31.9
4	—	71.3 (s)	70.9	133.9
5	1.94 (1H, m), 1.12 (1H, m)	48.3 (t)	47.5	124.6
6	1.94 (1H, m)	39.7 (d)	38.5	39.5
7	1.03 (1H, m)	47.7 (d)	46.0	45.4
8	1.65 (1H, m), 1.19 (1H, m)	22.4 (t)	21.7	21.2
9	2.03 (1H, m), 2.19 (1H, br d, 16.9)	28.7 (t)	27.9	32.3
10	—	137.0 (s)	136.3	124.1
11	1.84 (1H, m)	28.7 (d)	27.5	26.7
12	0.81 (3H, d, 6.8)	17.1 (q)	16.7	15.6
13	0.97 (3H, d, 6.8)	22.1 (q)	21.3	21.7
14	3.98 (1H, d, 11.6), 4.11 (1H, d, 11.6)	62.4 (t)	62.7	18.4
15	1.31 (3H, s)	25.6 (q)	25.6	23.5

COSY, HSQC, and HMBC spectrum.

Analysis of ¹H-¹H COSY and HSQC spectrum led to the identification of the partial structures CH₂(2)–CH₂(3), CH₂(8)–CH₂(9) and CH₃(12)–CH(11)–CH₃(13). The HMBC correlations of Me-15 (δ_{H} 1.31) with C-3, C-4 and C-5, and H-14 with C-1, C-9 and C-10 indicated that C-3, C-5, Me-15 and C-1, C-9, C-14 were connected to the quaternary carbons C-4 and C-10, respectively. Additional HMBC experiments showed the critical correlations from H-2 to C-1, C-3, C-6 and C-10, from H-7 to C-6 and C-9, from H-6 to C-4, and from both Me-12 (δ_{H} 0.81) and Me-13 (δ_{H} 0.97) to C-7 and C-11. The above analysis determined the planar structure of **1**. The relative configurations of **1** were deduced from analysis of the ROESY spectrum, which shows cross-peaks of H-6 with Me-12 and Me-15, H-8 α (δ_{H} 1.65) with Me-13, and H-7 with H-9 β (δ_{H} 2.19) indicating that H-6 and Me-15 were α -oriented, while H-7 was β -oriented. In the light of the evidence mentioned above, the structure of **1** was determined as 4 β ,14-dihydroxy-6 α ,7 β H-1(10)-cadinene.

1 showed cytotoxicity against C8166 cells (IC₅₀=76.9 μ g/ml) and significant anti-HIV-1 activity with EC₅₀=3.0 μ g/ml and SI (selectivity index) 25.4.

Acknowledgements This project was supported by Chinese Academy of Sciences (KSCX1-YW-R-24; KSCX2-YW-G-025).

References

1. Liu JK. Natural Terphenyls: Developments since 1887. *N*-Containing compounds of macromycetes. *Chem Rev* 106: 2209–2223 (2006)
2. Liu JK. *N*-Containing compounds of macromycetes. *Chem Rev* 105: 2723–2744 (2005)
3. Liu JK. Biologically active substances from mushrooms in Yunnan, China. *Heterocycles* 57: 157–167 (2002)
4. Wang XN, Tan RX, Liu JK. Xylactam, a new nitrogen-containing compound from the fruiting bodies of ascomycete *Xylaria euglossa*. *J Antibiot* 58: 268–270 (2005)
5. Luo DQ, Wang F, Bian XY, Liu JK. Rufuslactone, a new antifungal sesquiterpene from the fruiting bodies of the basidiomycete *Lactarius rufus*. *J Antibiot* 58: 456–459 (2005)
6. Shao HJ, Qing C, Wang F, Luo DQ, Liu JK. A new cytotoxic lanostane triterpenoid from the basidiomycete *Hebeloma versipelle*. *J Antibiot* 58: 828–831 (2005)
7. Barrfro AF, Sanchez JF, Oltra JE, Altarejos J, Ferrol N, Barragan A. *Phytochemistry* 30: 1151–1154 (1991)