

NOTE

Antibiotic R2, a new angucyclinone compound from *Streptosporangium* sp. Sg3

Hadjira Boudjella¹, Abdelghani Zitouni¹, Yannick Coppel², Florence Mathieu³, Marie-Carmen Monje³, Nasseridine Sabaou¹ and Ahmed Lebrihi³

The Journal of Antibiotics (2010) 63, 709–711; doi:10.1038/ja.2010.111; published online 13 October 2010

Keywords: actinomycete; new angucyclinone; *Streptosporangium*; structure elucidation

Angucyclinones are a subclass of angucyclines, which are defined as microbial quinone natural products related to tetracyclines and anthracyclines and bearing, as characteristic structural feature, a tetracyclic benz[*a*]anthracene ring system assembled in an angular manner.¹ This structural moiety is biosynthetically derived from a decaketide chain formed via the polyketide biosynthetic pathway. The term ‘angucycline’ includes molecules with hydrolysable sugar moieties, whereas ‘angucyclinone’ refers to a sugarless compound or a compound with a C-glycosidic linked sugar moiety.^{1,2} Angucyclines are isolated from the fermentation broth of actinomycetes, mostly from the genus *Streptomyces*, but also from *Actinomadura*, *Nocardia* and *Streptosporangium* genera.¹ Angucyclines show a broad spectrum of biological activities including antitumor,^{1,2} antibacterial, antifungal, antiviral,^{3,4} enzyme inhibitory^{5,6} and platelet aggregation inhibitory properties.^{7,8} In a continuous search for new bioactive compounds from actinomycetes other than the genus *Streptomyces*, several Saharan soil samples collected in arid ecosystems were explored.^{9–11} Among the isolates, we were interested by a new actinomycete strain belonging to the genus *Streptosporangium* and producing a new compound (**R2**) identified as a new angucyclinone. This paper describes the isolation, structure elucidation and antimicrobial activities of compound **R2**.

Details on the isolation and the taxonomy of the producing organism *Streptosporangium* sp. Sg3 were described in a previous paper.¹¹ To isolate compound **R2**, *Streptosporangium* sp. Sg3 was cultivated at 30 °C for 9 days in Erlenmeyer flasks (500 ml) containing 100 ml of ISP2 broth (yeast extract 4 g, malt extract 10 g, glucose 4 g, in 1 l distilled water, pH 7.2) on a rotary shaker (250 r.p.m.). The cultures (8 l) were centrifuged and filtered to remove mycelium. The culture filtrate was extracted with an equal volume of *n*-butanol to generate a crude extract (1.9 g). The latter was fractionated by size exclusion chromatography on Sephadex LH-20 (75% MeOH in H₂O), resulting in five fractions, I–V. Fraction IV including **R2**, was of

red color and exhibited an antibacterial activity. It was subjected to semipreparative reversed-phase HPLC using an Interchim UP5ODB column (250×7.8 mm) (Interchim, Montluçon, France) and developed using a continuous grade from 20 to 75% MeOH in H₂O (UV detection at 220 nm), yielding three active fractions (**1**, **2**, **3**). The major peak **2** with the main antibacterial activity was subjected to repeated HPLC to yield 3.5 mg of compound **R2**.

Compound **R2** was obtained as a red powder (optical rotation $[\alpha]_D^{25} +56$ (*c* 0.11, MeOH)). The ESI-MS spectrum contained an ion peak at *m/z* 475.1 [M–H][–], and its molecular formula was determined by HRESI-MS analysis as C₂₆H₁₉O₉ (calcd 475.43 for (M–H)[–], found 475.43). The UV absorption maximum at 535 nm was because of the red color and suggested a quinone chromophore. Absorptions at 3259, 2949 and 1721 cm^{–1} in the IR spectrum of **R2** were characteristic of hydroxy, methyl and carbonyl groups, respectively. **R2** was soluble in MeOH and DMSO, and insoluble in chloroform, *n*-hexane and H₂O.

The structure of **R2** (Figure 1) was determined by ¹H and ¹³C NMR spectroscopy and by using ¹H–¹H COSY45, ¹H–¹³C HMQC and ¹H–¹³C HMBC experiments. The ¹H and ¹³C chemical shifts of compound **R2** are given in Table 1. The ¹³C and Heteronuclear Single-Quantum Correlation spectra showed 26 carbon signals with a large number of quaternary carbons (17 out of 26). From the ¹³C data, it was possible to discern two keto-carbonyl groups (δ_C 188.4 and 186.5), one carboxylic acid group (δ_C 174.1), 19 sp²-hybridized carbons (δ_C from 160.2 to 114.7), two sp³-hybridized carbons bearing an electro-negative heteroatom (δ_C 73.0 and 72.5), one sp³-hybridized carbon (δ_C 22.6) and two methyl groups (δ_C 22.4 and 12.4). In CD₃OD, the ¹H NMR spectrum revealed two ortho-coupled aromatic protons (δ_H 7.82 and 7.66, 2H, *J*=7.5 Hz), two aromatic protons (δ_H 8.24 and 7.02, 2H, *s*), an AB system (δ_H 4.45 and 4.36, *J*_{AB}=10.7 Hz), an ethyl group (δ_H 2.83 and 1.32, 5H, *J*=7.4 Hz) and a methyl group (δ_H 2.70, 3H, *s*). In DMSO, NMR signals were significantly broader but the

¹Laboratoire de Recherche sur les Produits Bioactifs et la Valorisation de la Biomasse, Ecole Normale Supérieure de Kouba, Alger, Algérie; ²CNRS, LCC (Laboratoire de Chimie de Coordination), Toulouse, France and ³Département ‘Bioprocédés & Systèmes Microbiens’, ENSAT/INPT, Université de Toulouse, Laboratoire de Génie Chimique, UMR 5503 (CNRS/INPT/UPS), Castanet-Tolosan, France

Correspondence: Professor A Lebrihi, Département ‘Bioprocédés & Systèmes microbiens’, ENSAT/INPT, Université de Toulouse, Laboratoire de Génie Chimique, UMR 5503 (CNRS/INPT/UPS), 1 avenue de l’Agrobiopole, B.P. 32 607, Auzeville-Tolosane, Castanet-Tolosan 31 326, France.

E-mail: lebrihi@ensat.fr

Received 27 March 2010; revised 5 August 2010; accepted 3 September 2010; published online 13 October 2010




Figure 1 Structure of angucyclinone **R2** (a), and HMBC and COSY correlations (b).

Table 1 ¹H and ¹³C NMR data assignments of CD₃OD at 298 K

Position	δ_C , mult.	δ_H (J in Hz)
1	160.2, qC ^a	—
2	118.5, qC	—
3	143.1, qC	—
4	116.9, CH	7.02, s
4a	147.4, qC	—
5	72.5, CH	4.36, d (10.7)
6	73.0, CH	4.45, d (10.7)
6a	141.9, qC	—
7	114.7, CH	8.24, s
7a	131.6, qC	—
8	188.4, qC	—
8a	115.2 qC	—
9	160.1, qC	—
10	139.9, qC	—
11	135.3, CH	7.66, d (7.5)
12	118.7, CH	7.82, d (7.5)
12a	132.0, qC	—
13	186.5, qC	—
13a	117.4, qC	—
14	158.6, qC ^a	—
14a	128.6, qC	—
14b	114.9, qC	—
15	174.1, qC	—
16	22.4, CH ₃	2.70, s
17	22.6, CH ₂	2.83, q (7.4)
18	12.4, CH ₃	1.32, t (7.4)
OH-1 ^{b,c}	—	17.64, s
OH-5 ^c	—	5.69, s
OH-6 ^c	—	5.92, s
OH-9 ^{b,c}	—	13.98, s
OH-14 ^{b,c}	—	13.08, s
COOH ^{b,c}	—	13.20, s

^{a,b}Resonance assignment may be interchangeable.

^cIn DMSO-*d*₆.

Mult. stands for multiplicity. qC, CH, CH₂ and CH₃ indicate respectively quaternary, tertiary, secondary and primary carbons.

¹H NMR spectrum showed additional resonances: four enol or carboxylic acid protons (δ_H 17.64, 13.98, 13.20 and 13.08, 4H, s) and two hydroxyl signals (δ_H 5.92 and 5.69, 2H, s). The 2D ¹H-¹H and ¹H-¹³C experiments and especially the long range ¹H-¹³C couplings observed in the HMBC spectrum permitted to establish the presence of a benzo[*a*]naphthacenequinone skeleton. Only four carbons did not

show any correlation in the HMBC spectrum (C1, C7a, C14 and C15). Their assignment has been confirmed by comparison with NMR data of the pradimicin antibiotic family.¹² The peak of the four enol or carboxylic acid protons was too broad to give HMBC correlations and so could not be exactly assigned. The observed coupling constant of 10.7 Hz between H5 and H6 suggested that the dihedral angles between H5 and H6 are near 180° and is consistent with a *trans*-dial orientation. This value is similar with ones found for related compounds^{3,12} where a 5*S*,6*S* absolute configuration has been deduced.

The structure of **R2** was determined to be a benzo[*a*]naphthacene-2-carboxylic acid, 4,7,11,12-tetrahydro-1,5,6,9,14-pentahydroxy-8,13-dioxo-3-methyl-10-ethyl.

The minimum inhibitory concentrations (MICs) of **R2** were determined by a conventional agar dilution method using nutrient agar medium. Compound **R2** showed strong activity against the Gram-positive bacteria *Micrococcus luteus* ATCC 9314 and *Bacillus subtilis* ATCC 6633 (MICs=0.5 and 1 μg ml⁻¹, respectively), and moderate activity against *Staphylococcus aureus* CIP 7625 (MIC=10 μg ml⁻¹). A weak activity was observed against *Listeria monocytogenes* CIP 82110 and *Mycobacterium smegmatis* ATCC 607 (MICs=40 and 50 μg ml⁻¹, respectively). Compound **R2** showed no activity (MICs > 100 μg ml⁻¹) against the Gram-negative bacteria (*Escherichia coli* ATCC 10536 and *Pseudomonas fluorescens* number 412), the fungi (*Mucor ramannianus* NRRL 1829 and *Aspergillus carbonarius* number M333) and the yeasts (*Saccharomyces cerevisiae* ATCC 4226 and *Candida albicans* number 224).

In conclusion, from the results presented in this paper, we found that strain *Streptosporangium* sp. Sg3 produces a new angucyclinone that displayed antibacterial activities.

ACKNOWLEDGEMENTS

A part of this work was supported by the grant from 'Comité d'Evaluation et de Prospective de Coopération Inter-universitaire Franco-Algérienne' (Programme Hubert Curien Tassili).

- Rohr, J. & Thiericke, R. Angucycline group antibiotics. *Nat. Prod. Rep.* **9**, 103–137 (1992).
- Krohn, K. & Rohr, J. Angucyclines: total syntheses, new structures, and biosynthetic studies of an emerging new class of antibiotics. *Top. Curr. Chem.* **188**, 128–195 (1997).
- Tsunakawa, M. et al. The structures of pradimicins A, B, and C: a novel family of antifungal antibiotics. *J. Org. Chem.* **54**, 2532–2536 (1989).
- Kondo, S., Gomi, S., Ikeda, D., Hamada, M. & Takeuchi, T. Antifungal and antiviral activities of benanomyces and their analogues. *J. Antibiot.* **44**, 1228–1236 (1991).

- 5 Tsurumi, Y. *et al.* WS79089A, B and C, new endothelin converting enzyme inhibitors isolated from *Streptosporangium roseum* No. 79089. Taxonomy, fermentation, isolation, physico-chemical properties and biological activities. *J. Antibiot.* **47**, 619–630 (1994).
- 6 Matsuda, Y. & Kase, H. KS-619-1, a new inhibitor of Ca²⁺ and calmodulin-dependent cyclic nucleotide phosphodiesterase from *Streptomyces californicus*. *J. Antibiot.* **40**, 1104–1110 (1987).
- 7 Kawashima, A., Kishimura, Y., Tamai, M. & Hanada, K. New platelet aggregation inhibitors. *Chem. Pharm. Bull.* **37**, 3429–3431 (1989).
- 8 Kawashima, A. *et al.* PI-083, a new platelet aggregation inhibitor. *J. Antibiot.* **41**, 1913–1914 (1988).
- 9 Sabaou, N. *et al.* Les sols des oasis du Sahara algérien, source d'actinomycètes rares producteurs d'antibiotiques. *Sécheresse* **9**, 147–153 (1998).
- 10 Zitouni, A. *et al.* *Saccharothrix algeriensis* sp. nov., isolated from Saharan soil. *Int. J. Syst. Evol. Microbiol.* **54**, 1377–1381 (2004).
- 11 Boudjella, H. *et al.* Isolation and partial characterization of pigment-like antibiotics produced by a new strain of *Streptosporangium* isolated from an Algerian soil. *J. Appl. Microbiol.* **103**, 228–236 (2007).
- 12 Tsuno, T. *et al.* Biosynthesis of the pradimicin family of antibiotics. II. Fermentation, isolation and structure determination of metabolites associated with the pradimicins biosynthesis. *J. Antibiot.* **46**, 420–429 (1993).