

ORIGINAL ARTICLE

Synthesis and structure–activity relationships of novel lincomycin derivatives. Part 1. Newly generated antibacterial activities against Gram-positive bacteria with *erm* gene by C-7 modification

Yoshinari Wakiyama, Ko Kumura, Eijiro Umemura, Kazutaka Ueda, Satomi Masaki, Megumi Kumura, Hideki Fushimi and Keiichi Ajito

We synthesized 7(*S*)-7-deoxy-7-arylthiolincomycin derivatives possessing a heterocyclic ring at the C-7 position via sulfur atom by either Mitsunobu reaction of 2,3,4-tris-*O*-(trimethylsilyl)lincomycin or S_N2 reaction of 7-*O*-methanesulfonyl-2,3,4-tri-*O*-trimethylsilyllincomycin. As a result, 7(*S*)-7-deoxy-7-arylthiolincomycin derivatives 16, 21 and 27 exhibited antibacterial activities against respiratory infection-related Gram-positive bacteria with *erm* gene, although clindamycin did not have any activities against those pathogens. Furthermore, 7(*S*)-configuration of lincomycin derivatives was found to be necessary for enhancing antibacterial activities from the comparison results of configurations of 16 (*S*-configuration) and 30 (*R*-configuration) at the 7-position.

The Journal of Antibiotics (2016) 69, 368–380; doi:10.1038/ja.2015.119; published online 16 December 2015

INTRODUCTION

Macrolide antibiotics possess broad spectrum of antibacterial activity against Gram-positive bacteria (*Streptococcus pneumoniae*, *Streptococcus pyogenes*, etc.), *Haemophilus influenzae*, *Moraxella catarrhalis*, *Mycoplasma pneumoniae* and *Neisseria gonorrhoeae* and also have a safety profile as oral drugs. Therefore, macrolide antibiotics have been used as chemotherapeutic agents in clinical sites over many years. Resistant bacteria, however, have markedly increased recently,^{1–3} and this phenomenon has caused serious problems in the treatment of bacterial respiratory infections.

Macrolide antibiotics inhibit bacterial protein synthesis. Macrolide antibiotics have potent antibacterial activities through inhibiting elongation of an amino-acid sequence by binding to 23S ribosomal RNA.^{4–6} The mechanisms of action for bacteria to gain resistance are manifold, but in general, these can be characterized as involving drug efflux, alterations in the drug target site or drug inactivation. Resistant mechanisms in macrolides have diversified recently. Notably, there are major bacterial strains in clinical site, with point mutation in the drug target site,^{7,8} with methylase produced by *erm* gene and with efflux pump produced by *mef* gene.

The methylase of bacteria inhibits macrolide binding to 23S ribosomal RNA through either methylation or dimethylation at the N-6 position of the adenine residue (A2058Ec).^{7,9,10} On the other hand, the efflux pump of bacteria is able to transport a variety of compounds, thus conferring resistance to a broad range of antibiotics.

Clarithromycin¹¹ and azithromycin¹² (Figure 1) are not effective enough against resistant bacteria (*S. pneumoniae* and *S. pyogenes*) with *erm* gene and influenced by *S. pneumoniae* with *mef* gene. On the other hand, telithromycin (TEL),¹³ which was launched as the first ketolide antibiotic, is effective against resistant bacteria with *erm* and *mef* genes. X-ray crystallographic^{6,14} and footprinting analysis^{15–17} indicated that TEL would be capable to bind to not only domain V (A2058Ec, A2059Ec) of 23S rRNA but also to domain II (A752). TEL, however, has possibility to cause serious liver damage^{18,19} and loss of consciousness,^{20,21} and it is scarcely used in Japan. No oral antibiotic, which is effective against the resistant bacteria with *erm* and *mef* genes, with no problems in terms of safety or taste, has been launched so far.

Lincomycin (LCM)^{22–25} was isolated as a secondary metabolite from the fermentation broth of *Streptomyces lincolnensis*. Clindamycin (CLDM)²⁶ was synthesized by chemical modification of LCM (Figure 1) and exhibited improved antimicrobial activities and pharmacokinetics compared with LCM. But they are not effective against resistant bacteria (*S. pneumoniae* and *S. pyogenes*) with *erm* gene (Table 1).

Chemical modifications at the C-7 positions of LCM (7-dehydrolincomycin (7-ketolincomycin),²⁷ 7-deoxylincomycin,²⁸ lincomycin-7-acylate²⁹ and lincomycin-7-carbonate,²⁹ 7(*R*)-7-azido-7-deoxylincomycin,²⁵ 7(*R*)-7-amino-7-deoxylincomycin,²⁵ 7(*R*)-7-cyano-7-deoxylincomycin,²⁵ 7(*R*)-7-deoxy-7-thiolincomycin,³⁰ 7(*S*)-7-deoxy-7-thiolincomycin,³⁰ 7(*S*)-7-bromo-7-deoxylincomycin,²⁶ 7(*S*)-7-deoxy-7-iodolincomycin,²⁶ 7(*R*)-7-chloro-7-deoxylincomycin,²⁶




Figure 1 Chemical structures of the representative macrolides, lincomycin (LCM) and clindamycin (CLDM).

Table 1 Antibacterial activities (MIC, $\mu\text{g ml}^{-1}$) of the representative macrolides, lincomycin (LCM) and clindamycin (CLDM)

Test organism ^a	Characteristics ^b	CAM	AZM	LCM	CLDM
<i>Streptococcus pneumoniae</i> DP1 TypeI	susceptible	0.03	0.06	1	0.06
<i>S. pneumoniae</i> -2	susceptible	0.03	0.03	1	0.12
<i>S. pneumoniae</i> -3	susceptible	0.015	0.03	0.25	0.06
<i>S. pneumoniae</i> -4	<i>ermAM</i> methylase(c)	>128	>128	>128	>128
<i>S. pneumoniae</i> -5	<i>ermAM</i> methylase(c)	>128	>128	>128	>128
<i>S. pneumoniae</i> -6	<i>ermAM</i> methylase(c) + <i>mefE</i>	>128	>128	>128	>128
<i>S. pneumoniae</i> -7	<i>ermAM</i> methylase(i)	>128	>128	128	128
<i>S. pneumoniae</i> -8	<i>ermAM</i> methylase(i)	>128	>128	128	128
<i>S. pneumoniae</i> -9	<i>mefE</i> efflux	0.5	0.5	1	0.12
<i>Streptococcus pyogenes</i> Cook	susceptible	0.015	0.06	0.12	0.06
<i>S. pyogenes</i> -2	<i>ermAM</i> methylase(c)	>128	>128	>128	128
<i>S. pyogenes</i> -3	<i>mefE</i> efflux	8	8	0.25	0.12
<i>Haemophilus influenzae</i>	susceptible	2	0.25	8	16
<i>H. influenzae</i> -2	susceptible	4	1	16	8
<i>H. influenzae</i> -3	susceptible	8	2	16	16
<i>H. influenzae</i> -4	Δ acr	0.5	0.5	4	1

Abbreviations: AZM, azithromycin; CAM, clarithromycin.
^aAll strains except standard organisms were clinically isolated.
^b(c): constitutive; (i): inducible.
 Grey shading strains are target strains.

and 7-epilincosamin (7(S)-lincomycin),²⁶ 7-deoxy-7-methylincosamin³¹ and so on) have been investigated so far. In these compounds, 7(S)-7-bromo-7-deoxyincosamin and 7(S)-7-deoxy-7-iodincosamin exhibited enhanced antibacterial activities in the same order of magnitude as CLDM, and 7(R)-7-azido-7-deoxyincosamin had the same antibacterial activities as LCM.

In the case of possessing a chlorine atom at the 7-position, CLDM (7(S)-Cl) had stronger antibacterial activities than 7-epiclidamycin (7(R)-Cl). 7-Epilincosamin (7(S)-lincomycin) had one-half of the potency of the LCM possessing 7(R)-configuration (7(S)-OH < 7(R)-OH).




7(R)-7-O-methylincosamin, of which 7-hydroxyl group was protected, showed improved potency, whereas 7(S)-7-O-methylincosamin had stronger activities of 3.5 times the response of LCM against *Sarcina lutea* (7(S)-OMe > 7(R)-OMe).^{32,33} Unfortunately, both larger alkoxy groups and substituted alkoxy groups resulted in weaker antibacterial activities than those of LCM.

On the other hand, both 7(S)-7-deoxy-7-thiolincosamin and 7(R)-7-deoxy-7-thiolincosamin showed only 10% activity as compared with LCM. 7(S)-7-alkylthio-7-deoxyincosamin and 7(S)-7-substituted alkylthio-7-deoxyincosamin^{34–38} were more active

than LCM against Gram-positive or Gram-negative organisms, and 7(R)-7-deoxy-7-imidazol-2-yl-thiolincosamin, which had been reported by Sztaricskai *et al.*,³⁹ retained the same order of magnitude of antibacterial activities against tested organisms as those of LCM. Antibacterial activities were affected by both configuration and a structure of a substituent at the 7-position. 7(S)-7-deoxy-7-phenylthiolincosamin was reported by Bannister *et al.*³⁸ but no antibacterial activity of this compound was reported.


Thus we were interested in derivatives with a heterocyclic ring introduced via sulfur atom at the C-7 position with 7(S)-configuration. According to the above publications, LCM derivatives exhibited much stronger *in vitro* activities compared with LCM, but no LCM derivatives were effective against resistant bacteria with *erm* gene.

LCM and CLDM inhibited bacterial protein synthesis similar to macrolide antibiotics. The structures of lincomycin derivatives are different from those of macrolide antibiotics, but X-ray crystallographic analysis indicated that their binding sites to rRNA are closely located in a neighboring area.^{4,6} According to the report,^{4,6} there were several major interactions by hydrogen bonding between



Scheme 1 Synthesis of 7(S)-7-arylthio-7-deoxylincomycin derivatives. Conditions: a, TMSCl, HMDS, Py, r.t., 2 h; b, 80% AcOH, MeOH, r.t., 16 h; Method A: c, PPh₃, DEAD, the corresponding HS-Ar, THF or toluene, 0 °C to 50 °C, 16–18 h, d, 1–2 N HCl, MeOH, r.t., 30 min; Method B: c, PBU₃, DEAD, the corresponding Ar-S-S-Ar, THF, 0 °C to r.t., 19–25 h; d, 2 N HCl, r.t., 0.5–3 h; Method C: c, PPh₃, DEAD, *tert*-butyl 5-mercapto-1,3,4-thiadiazol-2-yl(methyl)carbamate or *tert*-butyl 5-mercapto-1,3,4-thiadiazol-2-yl(methyl)carbamate, THF, 0 °C to r.t., 17 h, d, TFA, r.t., 60 min; e, NaOMe, allyl iodide, MeOH, r.t.; f, SnCl₂·H₂O, NaBH₄, EtOH, r.t., 3 h; g, 6-aminobenzo[d]thiazole-2-thiol, K₂CO₃, DMF, 100 °C, 16 h; h, SnCl₂·H₂O, NaBH₄, EtOH, r.t., 3 h; i, MsCl, TEA, DMF, r.t., 20 min; j, PPh₃, DEAD, *tert*-butyl 2-(5-mercapto-1,3,4-thiadiazol-2-ylamino)-2-oxoethylcarbamate, THF, 0 °C to r.t., 17 h, k, TFA, r.t., 0.5 h; l, TMSCl, HMDS, Py, r.t., 2.5 h; m, 2-methoxyacetyl chloride, TEA, THF, 0 °C, 2 h; n, 1 N HCl, MeOH, r.t., 2 h; o, MsCl, TEA, CHCl₃, r.t., 3 h; p, the corresponding HS-Ar, K₂CO₃, DMF, 80 °C, 3–16 h, q, 1 N HCl, MeOH, r.t., 0.5–2 h.

Table 2 Antibacterial activities (MIC, $\mu\text{g ml}^{-1}$) of 7-thio-substituted LCM derivatives



Test organism ^a	Characteristics ^b	CLDM	2	3	4	6	7	8	9	10	11	28
<i>Streptococcus pneumoniae</i> DP1 Type1	susceptible	0.06	0.06	0.25	0.06	0.25	0.12	0.12	0.12	0.12	0.12	0.06
<i>S. pneumoniae</i> -2	susceptible	0.12	0.06	0.25	0.12	0.25	0.25	0.25	0.25	0.25	0.25	0.12
<i>S. pneumoniae</i> -3	susceptible	0.06	0.06	0.25	0.03	0.12	0.12	0.12	0.12	0.12	0.12	0.06
<i>S. pneumoniae</i> -4	<i>ermAM</i> methylase(c)	>128	128	128	>128	128	>128	>128	128	>128	128	>128
<i>S. pneumoniae</i> -5	<i>ermAM</i> methylase(c)	>128	>128	128	>128	128	>128	>128	128	128	128	>128
<i>S. pneumoniae</i> -6	<i>ermAM</i> methylase(c) + <i>mefE</i>	>128	>128	128	>128	128	>128	>128	128	>128	128	>128
<i>S. pneumoniae</i> -7	<i>ermAM</i> methylase(i)	128	>128	128	>128	128	128	>128	128	64	128	128
<i>S. pneumoniae</i> -8	<i>ermAM</i> methylase(i)	128	64	128	>128	128	128	>128	128	64	128	>128
<i>S. pneumoniae</i> -9	<i>mefE</i> efflux	0.12	0.06	0.25	0.03	0.12	0.06	0.12	0.12	0.12	0.12	0.06
<i>Streptococcus pyogenes</i> Cook	susceptible	0.06	0.03	0.12	0.03	0.12	0.12	0.12	0.12	0.06	0.06	0.06
<i>S. pyogenes</i> -2	<i>ermAM</i> methylase(c)	128	64	16	64	128	64	64	64	128	128	>128
<i>S. pyogenes</i> -3	<i>mefE</i> efflux	0.12	0.06	0.25	0.12	0.12	0.25	0.25	0.25	0.12	0.25	0.12
<i>Haemophilus influenzae</i>	susceptible	16	8	64	32	16	32	32	32	64	16	16
<i>H. influenzae</i> -2	susceptible	8	4	16	32	8	8	16	8	32	16	8
<i>H. influenzae</i> -3	susceptible	16	16	64	64	32	32	64	64	128	32	32
<i>H. influenzae</i> -4	Δ acr	1	0.5	2	1	0.5	1	2	0.5	1	1	1

Abbreviations: CLDM, clindamycin; LCM, lincomycin.

^aAll strains except standard organisms were clinically isolated.

^b(c): constitutive; (i): inducible.

Grey shading strains are target strains.

the peptidyl transferase cavities (A2058Ec, A2059Ec and G2520Ec) and hydroxyl groups at the sugar portion of CLDM. These data suggest that it is difficult for us to improve antibacterial activity by chemical modification at the sugar moiety. In fact, 2-deoxylincomycin⁴⁰ has been reported to show only 1% activity compared with LCM.

As we previously described, TEL has a risk of causing serious side effects^{18–21} and it is rarely used in Japan. Furthermore, its production cost seems to be relatively high owing to its complicated structure. On the other hand, CLDM possessing a simple structure has similar antibacterial activity against susceptible strains such as clarithromycin and exhibits acceptable oral absorption in animals and humans. However, CLDM is not effective enough against resistant bacteria of *S. pneumoniae* or *S. pyogenes* with *erm* gene. So a novel oral lincomycin analog, which is effective against resistant bacteria with *erm* gene and/or *mef* gene and does not have any problems in safety, taste or pharmacokinetics, is strongly desired for treatment of respiratory infections in clinical sites.

On the other hand, 7(S)-7-azido-7-deoxylincomycin, CLDM, 7(S)-7-bromo-7-deoxylincomycin and 7(S)-7-alkylthio-7-deoxylincomycin had the same or stronger antibacterial activity compared with LCM. According to the relationships between configuration and a substituent (7(S)-Cl > 7(R)-Cl and 7(S)-OMe > 7(R)-OMe), enhancement of activities in 7(S)-configuration might require a hydrophobic substituent such as a chlorine atom or a methyl group, not as OH or SH (7(S)-OH < 7(R)-OH (LCM), 7(S)-SH \approx 7(R)-SH < LCM). Larger alkoxy groups and substituted alkoxy groups with 7(S)-configuration resulted in weaker antibacterial activities than those of LCM, while 7(S)-7-alkylthio-7-deoxylincomycin was more active than LCM against Gram-positive organisms. As a result, a sulfur atom might be preferable to an oxygen atom at the 7-position for enhancing antibacterial activities. 7(R)-7-deoxy-7-imidazol-2-yl-thiolincomycin

had the same antibacterial activity as LCM. Here we planned synthesis of lincomycin analogs possessing a heterocyclic ring via sulfur atom focusing on the 7(S)-configuration at the 7-position, in order to generate unreported antibacterial activity against respiratory infection-related Gram-positive bacteria with *erm* gene.


RESULTS AND DISCUSSION

Design of LCM derivatives

The results of X-ray crystallographic analysis^{4–6} have already been reported in application of CLDM. According to three-dimensional information around the 7-position of CLDM, we hypothesized that CLDM had enough three-dimensional space around the 7-position, and it might be able to enhance antibacterial activities by filling the space with an appropriate substituent. So we designed 7(S)-LCM derivatives possessing a hydrophobic heterocyclic ring via sulfur atom and practically synthesized 7(S)-7-arylthio-7-deoxylincomycin by Mitsunobu reaction.

Synthesis of 7(S)-7-arylthio-7-deoxylincomycin derivatives

Synthesis of 7(S)-7-arylthio-7-deoxylincomycin derivatives is outlined in Scheme 1. We first prepared a key intermediate **1**⁴¹ derived from LCM in two steps in order to construct the same configuration at the 7-position as CLDM in final target molecules. Our synthesis began with silylation of all OH groups of LCM, and the 7-O-TMS group was selectively deprotected by AcOH to give the key intermediate (**1**) with an excellent yield. LCM derivatives (**2–5**, **7–17** and **21–27**), which possess a heterocyclic ring at the C-7 position via sulfur atom, were synthesized from **1** with the corresponding thiol under the Mitsunobu condition (Methods A and C).^{42,43} Compounds **6**³⁸ and **18–20** possessing a phenyl thiol group were synthesized by Method B in application of the corresponding disulfide. Furthermore, compound **28**³⁸ was prepared by allylation of **5** under the NaOMe

Table 3 Antibacterial activities (MIC, $\mu\text{g ml}^{-1}$) of LCM derivatives possessing a substituted benzothiazol-2-ylthio group at the 7-position


Test organism ^a	Characteristics ^b	12	13	14	15	16	29	30
<i>Streptococcus pneumoniae</i> DPI Type1	susceptible	0.12	0.12	0.25	0.06	0.06	0.12	0.03
<i>S. pneumoniae</i> -2	susceptible	0.12	0.12	0.25	0.06	0.06	0.12	0.03
<i>S. pneumoniae</i> -3	susceptible	0.12	0.12	0.12	0.03	0.03	0.06	0.03
<i>S. pneumoniae</i> -4	<i>ermAM</i> methylase(c)	64	64	128	64	8	32	32
<i>S. pneumoniae</i> -5	<i>ermAM</i> methylase(c)	32	>128	128	64	32	64	32
<i>S. pneumoniae</i> -6	<i>ermAM</i> methylase(c) + <i>mefE</i>	64	>128	128	64	64	128	64
<i>S. pneumoniae</i> -7	<i>ermAM</i> methylase(i)	64	128	64	32	16	32	32
<i>S. pneumoniae</i> -8	<i>ermAM</i> methylase(i)	32	128	64	32	8	64	32
<i>S. pneumoniae</i> -9	<i>mefE</i> efflux	0.12	0.12	0.12	0.06	0.03	0.06	0.03
<i>Streptococcus pyogenes</i> Cook	susceptible	0.12	0.06	0.12	0.06	0.03	0.06	0.03
<i>S. pyogenes</i> -2	<i>ermAM</i> methylase(c)	16	32	16	16	4	32	16
<i>S. pyogenes</i> -3	<i>mefE</i> efflux	0.25	0.12	0.25	0.06	0.06	0.12	0.06
<i>Haemophilus influenzae</i>	susceptible	32	32	64	32	8	32	16
<i>H. influenzae</i> -2	susceptible	32	16	16	16	4	8	16
<i>H. influenzae</i> -3	susceptible	64	64	64	64	32	32	32
<i>H. influenzae</i> -4	Δ acr	2	1	1	0.25	0.25	0.5	0.5

Abbreviation: LCM, lincomycin.

^aAll strains except standard organisms were clinically isolated.

^b(c): constitutive; (i): inducible.

Grey shading strains are target strains.

condition to investigate structure–activity relationship (SAR) of an allyl group. As a part of chemical modification of the benzothiazole group, **29** was also prepared by reduction of the nitro group of **17** using $\text{SnCl}_2 \cdot \text{H}_2\text{O} \cdot \text{NaBH}_4$. Next, **30** was synthesized by $\text{S}_{\text{N}}2$ reaction of CLDM with the corresponding thiol under the basic condition to confirm the effect on antibacterial activity by configuration at the C-7 position. On the other hand, the nitrophenyl derivatives (**18–20**), which were synthesized under the Mitsunobu condition, were reduced to give the corresponding aminophenyl derivatives (**31–33**). Because compound **33** was the most potent compared with **31** or **32**, **33** was converted to compound **34**. A Boc group and three trimethylsilyl (TMS) groups of **35** prepared by the Mitsunobu condition were simultaneously deprotected to give a desired compound **36**. TMS protection of **21** gave **37**, which was reacted with 2-methoxyacetyl chloride with triethylamine, and then deprotected to provide a methoxyacetyl derivative (**38**).

On the other hand, Mitsunobu conditions still had problems, and thus compounds **18**, **20** and **40** were synthesized in only low yields. So compounds **20**, **40** and **41** were synthesized by $\text{S}_{\text{N}}2$ reactions with the corresponding thiol in good yields under the basic conditions in application of a methanesulfonate **39** synthesized from **1**. SAR of novel LCM derivatives are shown in Tables 2–5.

SAR analysis of LCM derivatives possessing a heterocyclic ring at the C-7 position via sulfur atom


Antibacterial activity of LCM derivatives, which possessed a heterocyclic ring, the benzene ring or an allyl group at the C-7 position via

sulfur atom, is shown in Table 2. Compound **2** was shown to be slightly more potent against *H. influenzae* than CLDM. Compound **3** had weak activity against *S. pyogenes* with *erm* gene, although CLDM did not have any activities against the pathogen. So we pursued further optimization at the 7-position. As a result, derivatives possessing a phenyl, thienyl or 1,3,4-thiadiazolyl group did not exhibit remarkable activity against *S. pyogenes* with *erm* gene.

Filling a space around the C-7 position might enhance antibacterial activities against resistant *S. pyogenes* with *erm* gene. Thus we selected **3** as a fundamental bicycle framework for further optimization. Additionally, **6** and **11** were selected as general examples of an isolated 6-membered ring and an isolated 5-membered ring, respectively. Next we tried to enhance antibacterial activities by introducing a substitute to the selected three frameworks.

SAR analysis of LCM derivatives possessing a benzothiazol-2-ylthio group at the 7-position

LCM derivatives having a benzothiazoyl or a thiazoropyridyl group at the 7-position were synthesized and their antibacterial activities are shown in Table 3. Notably, conversion of **3** to **12**, **15** or **16** at the 7-position improved antibacterial activity against *S. pneumoniae* with *erm* gene. Especially, antibacterial activity of **16** was significantly enhanced. An amino group might have a role to enhance antibacterial activity and may interact with a certain binding site on 23S rRNA. Compound **29** possessing an amino group at the 5-position in the benzothiazole ring generally exhibited improved antibacterial activity against *S. pneumoniae* with *erm* gene compared with **3**. On the other hand, we prepared **30** possessing the (*R*)-configuration at the

Table 4 Antibacterial activities (MIC, $\mu\text{g ml}^{-1}$) of LCM derivatives possessing a substituted phenylthio group at the 7-position


Test organism ^a	Characteristics ^b	18	19	20	31	32	33	34	40	41
<i>Streptococcus pneumoniae</i> DP1 Type1	susceptible	1	0.25	0.25	0.5	0.25	0.12	0.03	0.25	0.12
<i>S. pneumoniae</i> -2	susceptible	1	0.25	0.25	1	0.25	0.25	0.03	0.25	0.25
<i>S. pneumoniae</i> -3	susceptible	0.5	0.25	0.12	0.5	0.25	0.12	0.03	0.06	0.06
<i>S. pneumoniae</i> -4	<i>ermAM</i> methylase(c)	>128	128	128	>128	>128	>128	64	32	128
<i>S. pneumoniae</i> -5	<i>ermAM</i> methylase(c)	>128	128	128	>128	>128	>128	128	64	128
<i>S. pneumoniae</i> -6	<i>ermAM</i> methylase(c) + <i>mefE</i>	128	128	>128	>128	>128	>128	128	128	128
<i>S. pneumoniae</i> -7	<i>ermAM</i> methylase(i)	128	128	128	>128	>128	128	16	32	128
<i>S. pneumoniae</i> -8	<i>ermAM</i> methylase(i)	128	128	128	>128	>128	128	16	32	128
<i>S. pneumoniae</i> -9	<i>mefE</i> efflux	0.5	0.25	0.12	0.25	0.25	0.06	0.03	0.06	0.06
<i>Streptococcus pyogenes</i> Cook	susceptible	0.5	0.12	0.12	0.12	0.25	0.06	0.03	0.25	0.06
<i>S. pyogenes</i> -2	<i>ermAM</i> methylase(c)	>128	128	128	>128	>128	>128	8	32	32
<i>S. pyogenes</i> -3	<i>mefE</i> efflux	1	0.25	0.25	0.25	0.25	0.12	0.03	0.25	0.25
<i>Haemophilus influenzae</i>	susceptible	>128	128	128	128	32	16	4	64	64
<i>H. influenzae</i> -2	susceptible	128	64	32	32	8	8	16	32	64
<i>H. influenzae</i> -3	susceptible	>128	128	>128	>128	32	32	32	128	128
<i>H. influenzae</i> -4	Δ lacr	64	1	1	2	1	0.5	0.25	1	1

Abbreviation: LCM, lincomycin.

^aAll strains except standard organisms were clinically isolated.^b(c): constitutive; (i): inducible.

Grey shading strains are target strains.

7-position to confirm its potency. Based on comparison of antibacterial activities of compound **30** to those of compound **16**, we found that (*S*)-configuration at the 7-position was important for potent antibacterial activities.

SAR analysis of LCM derivatives possessing a phenylthio group at the 7-position


Next we consequently pursued modification of LCM derivatives possessing a substituted phenyl group at the C-7 position via sulfur atom and their antibacterial activities are shown in Table 4. The enhanced antibacterial activities by introducing an amino (**16**) or a nitro group (**15**) as shown in Table 3 encouraged us to further explore SAR through introducing an amino or a nitro group at the *o*-, *m*- and *p*-position on the phenyl ring. Unexpectedly, compounds **18–20** and **31–33** showed weaker antibacterial activities against resistant bacteria with *erm* gene than **15** or **16**. Among nitrophenyl and aminophenyl derivatives in Table 4, the *para*-amino derivative (**33**) exhibited relatively stronger activities against susceptible *S. pneumoniae* and *H. influenzae*. Conversion of the amino group to a methoxycarbonyl group (**33** to **40**) at the *p*-position on the phenyl ring improved antibacterial activities against bacteria with *erm* gene. Based on SAR analysis of compounds **18–20**, **31–33**, **40** and **41**, *para*-substituted analogs exhibited relatively stronger antibacterial activities against resistant bacteria than *o*-substituted or *m*-substituted analogs. Furthermore, **34** possessing a methanesulfonyl group at the amino group of **33** exhibited the most potent antibacterial activity in the phenylthio derivatives, except some strains belonging to *S. pneumoniae* with *erm* gene. These results suggest that (i) it is important for a substituent to keep a specific size, length and three-dimensional direction for appropriate binding to rRNA, and (ii) there are several important hydrogen bondings with some functional moieties, such as C=O, N=O, S=O or NH₂.

SAR analysis of LCM derivatives possessing a 1,3,4-thiadiazolylthio group at the 7-position

Next we pursued chemical modification of LCM derivatives possessing a substituted 1,3,4-thiadiazole at the C-7 position via sulfur atom and their antibacterial activities are shown in Table 5. We could find several important functional moieties for improvement of activity so far. Then we first introduced an amino group on the 1,3,4-thiadiazole ring. Compound **21** exhibited improved activities against both resistant bacteria with *erm* gene and *H. influenzae* compared with **11**. We synthesized several compounds **22**, **23**, **36** and **38**, which have an additional group introduced to the amino group of **21**. Compound **36** possessing a glyceryl moiety as an aliphatic amine had weaker antibacterial activities than **21**. Conversion of the glyceryl moiety to a methoxyacetyl moiety (**36** → **38**) improved antibacterial activities. Furthermore, derivatives **22** and **23** were synthesized and a difluoromethylthiophenyl analog (**23**) showed stronger activities against *S. pneumoniae* with *erm* gene than **21**. These results suggest that the phenyl moiety of **23** is an important group and that it enhances antibacterial activities against resistant bacteria with *erm* gene. We synthesized **27** in order to confirm this hypothesis. As expected, **27** showed relatively stronger antibacterial activities against bacteria with *erm* gene compared with **21**. Finally, we were interested in conversion of a NH-CO bonding to a CO-NH bonding (**36** and **38** → **24–26**). Compound **26** also exhibited effective antibacterial activities against bacteria with *erm* gene. These results suggest that filling a space around the 7-position of LCM has an important role to enhance antibacterial activities by hydrogen bonding, π - π stacking or CH- π interaction to undefined binding site on 23S rRNA.

CONCLUSION

With a purpose of generating new effective templates against bacteria, we first prepared the key intermediate **1**⁴¹ derived from LCM with two

Table 5 Antibacterial activities (MIC, $\mu\text{g ml}^{-1}$) of LCM derivatives possessing a substituted 1, 3, 4-thiadiazolythio group at the 7-position


Test organism ^a	Characteristics ^b	21	22	23	24	25	26	27	36	38
<i>Streptococcus pneumoniae</i> DP1 TypeI	susceptible	0.03	0.06	0.12	0.25	0.12	0.12	0.06	0.25	0.06
<i>S. pneumoniae</i> -2	susceptible	0.06	0.06	0.12	0.25	0.12	0.12	0.06	0.5	0.03
<i>S. pneumoniae</i> -3	susceptible	0.06	0.03	0.12	0.12	0.06	0.12	0.03	0.25	0.12
<i>S. pneumoniae</i> -4	<i>ermAM</i> methylase(c)	16	64	16	128	64	16	8	128	64
<i>S. pneumoniae</i> -5	<i>ermAM</i> methylase(c)	64	128	16	>128	64	16	8	>128	64
<i>S. pneumoniae</i> -6	<i>ermAM</i> methylase(c) + <i>mefE</i>	128	N.D.	32	>128	N.D.	128	64	>128	64
<i>S. pneumoniae</i> -7	<i>ermAM</i> methylase(i)	16	32	8	128	32	8	8	128	8
<i>S. pneumoniae</i> -8	<i>ermAM</i> methylase(i)	16	16	2	>128	32	16	8	128	8
<i>S. pneumoniae</i> -9	<i>mefE</i> efflux	0.03	0.03	0.12	0.25	0.12	0.12	0.06	0.25	0.06
<i>Streptococcus pyogenes</i> Cook	susceptible	0.03	0.06	0.12	0.12	0.12	0.12	0.06	0.25	0.06
<i>S. pyogenes</i> -2	<i>ermAM</i> methylase(c)	8	32	8	128	16	16	2	64	8
<i>S. pyogenes</i> -3	<i>mefE</i> efflux	0.06	0.06	0.12	0.25	0.12	0.25	0.12	1	0.06
<i>Haemophilus influenzae</i>	susceptible	8	8	16	64	8	32	16	64	32
<i>H. influenzae</i> -2	susceptible	4	8	8	64	8	64	8	64	16
<i>H. influenzae</i> -3	susceptible	8	16	32	64	32	128	32	128	32
<i>H. influenzae</i> -4	Δ <i>lacR</i>	0.25	0.5	0.25	2	0.5	2	0.5	4	0.5

Abbreviations: LCM, lincomycin; ND, not detected.

^aAll strains except standard strains were clinically isolated.

^b(c): constitutive; (i): inducible.

Grey shading strains are target strains.

steps to synthesize LCM analogs possessing the same configuration at the 7-position as CLDM in final target molecules. Our LCM derivatives were generally synthesized under Mitsunobu condition with the corresponding thiol from the key intermediate (1).

Compounds 16, 21 and 27 exhibited antibacterial activities against respiratory infection-related Gram-positive bacteria with *erm* gene, although CLDM did not have any activities against those pathogens. Furthermore, we confirmed that 7(S)-configuration was necessary for enhancing antibacterial activities on the basis of comparison results of configurations of 16 and 30. This work suggests that LCM derivatives may overcome resistant bacteria. SAR analysis, as indicated in this paper, would be useful for further medicinal chemistry in application of LCM derivatives. Our synthetic research in LCM derivatives is in progress.

EXPERIMENTAL PROCEDURE

General methods

¹H NMR spectra were measured with a BRUKER Ascend 400 NMR spectrometer (BRUKER corporation, Coventry, UK) for 400 MHz, JEOL JNM-GSX 400 NMR spectrometer (JEOL Ltd., Tokyo, Japan) for 400 MHz or a Varian Gemini 300 NMR spectrometer (Varian Inc., Palo Alto, CA, USA) for 300 MHz in CDCl₃ or CD₃OD. TMS (0 p.p.m.) in CDCl₃ or CD₃OD was used as an internal reference standard. Mass spectra (MS) were obtained on a JEOL JMS-700 mass spectrometer (JEOL Ltd.) or Agilent Technologies 6530-Q-TOF LC/MS mass spectrometer (Agilent Technologies, Santa Clara, CA, USA). The optical rotations were recorded with Jasco P-2300 digital polarimeter (Jasco corporation, Tokyo, Japan). Column chromatography was performed with silica gel (Wakogel C200, Wako Pure Chemical Industries Ltd., Osaka, Japan). Preparative thin layer chromatography was performed with silica gel (Merck, Darmstadt, Germany; TLC plates Silica gel 60 F254). All organic extracts were dried over anhydrous MgSO₄, and the solvent was removed with a rotary evaporator under reduced pressure.

2,3,4-Tris-O-(trimethylsilyl)lincomycin (1). To a solution of lincomycin (50 g, 123 mmol) in pyridine (200 ml) were added trimethylchlorosilane (90 ml, 704 mmol) and hexamethyldisilazane (65 ml, 310 mmol) and stirred at room temperature (RT) for 2 h, and then it was concentrated under reduced pressure. The residue was diluted with water, then extracted with hexane, washed with water and concentrated under reduced pressure. To a solution of the resulting residue in methanol (150 ml) was added 80% aqueous acetic acid (22.5 ml) stirred at RT for 16 h. The mixture was diluted with saturated aqueous NaHCO₃ (30 ml) and concentrated under reduced pressure. The residue was diluted with water and hexane, then extracted with hexane, washed with water, dried over MgSO₄ and concentrated under reduced pressure. The title compound was obtained as a colorless solid (69.5 g, 91%). ESI-MS (*m/z*) 623 (M+H)⁺ as C₂₇H₅₈N₂O₆Si₃; ¹H NMR (400 MHz, chloroform-*d*) δ 0.14 (s, 18 H), 0.18 (s, 9 H), 0.85–0.93 (m, 3 H), 1.14 (d, *J* = 6.4 Hz, 3 H), 1.22–1.35 (m, 4 H), 1.79–1.90 (m, 1 H), 1.92–2.07 (m, 3 H), 2.09 (s, 3 H), 2.38 (s, 3 H), 3.00 (dd, *J* = 10.9, 3.9 Hz, 1 H), 3.07 (br d, *J* = 1.6 Hz, 1 H), 3.12–3.21 (m, 1 H), 3.59 (dd, *J* = 9.5, 2.5 Hz, 1 H), 3.80 (br d, *J* = 2.5 Hz, 1 H), 4.00 (d, *J* = 9.5 Hz, 1 H), 4.04–4.13 (m, 1 H), 4.15 (dd, *J* = 9.5, 5.6 Hz, 1 H), 4.27–4.33 (m, 1 H), 5.21 (d, *J* = 5.6 Hz, 1 H), 7.42 (d, *J* = 9.8 Hz, 1 H).

7(S)-7-(Benzo[d]oxazol-2-ylthio)-7-deoxylincomycin (2). To a solution of compound 1 (240 mg, 0.39 mmol) in tetrahydrofuran (THF; 5 ml) at 0 °C were added triphenylphosphine (150 mg, 0.57 mmol), diethylazodicarboxylate (0.10 ml, 0.55 mmol) and benzo[d]oxazole-2-thiol (85 mg, 0.56 ml) and stirred at 0 °C for 1 h. The mixture was stirred at RT for 16 h. The mixture was diluted with 2 N HCl (1 ml)-MeOH (1 ml) and then stirred at RT for 30 min and concentrated under reduced pressure. The resulting residue was dissolved by water and washed with diethyl ether. The mixture was added to NaHCO₃ (150 mg), then extracted with ethyl acetate, washed with water, dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by preparative TLC (CHCl₃/CH₃OH/28% aq NH₄OH = 20/1/0.1) to obtain the title compound as a colorless solid (147.6 mg, 71%). [α]_D²⁶ +77.6° (c 0.77, MeOH); ESI-MS (*m/z*) 540 (M+H)⁺ as C₂₅H₃₇N₃O₆S₂; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₅H₃₇N₃O₆S₂: 540.2202, found: 540.2204; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.85–0.97 (m, 3 H), 1.26–1.40 (m, 4 H), 1.61

(d, $J=6.8$ Hz, 3 H), 1.77–1.85 (m, 1 H), 1.87 (s, 3 H), 1.97–2.09 (m, 2 H), 2.12–2.28 (m, 1 H), 2.35 (s, 3 H), 2.98 (dd, $J=10.5$, 5.2 Hz, 1 H), 3.21 (dd, $J=8.5$, 6.2 Hz, 1 H), 3.56 (dd, $J=10.2$, 3.2 Hz, 1 H), 3.83 (br d, $J=3.2$ Hz, 1 H), 4.10 (dd, $J=10.2$, 5.6 Hz, 1 H), 4.43 (br d, $J=9.8$ Hz, 1 H), 4.45 (dq, $J=6.8$, 3.2 Hz, 1 H), 4.64 (dd, $J=9.8$, 3.2 Hz, 1 H), 5.24 (d, $J=5.6$ Hz, 1 H), 7.27–7.36 (m, 2 H), 7.49–7.60 (m, 2 H).

7(S)-7-(Benzo[d]thiazol-2-ylthio)-7-deoxylincomycin (3). Compound **1** (320 mg, 0.51 mmol) and benzo[d]thiazole-2-thiol (250 mg, 1.49 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **3** (225.5 mg, 79%) as a colorless solid. $[\alpha]_D^{26}$ 74.1° (c 1.05, MeOH); ESI-MS (m/z) 556 (M+H)⁺ as C₂₅H₃₇N₃O₅S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₅H₃₇N₃O₅S₃: 556.1974, found: 556.1975; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.85–0.98 (m, 3 H), 1.24–1.40 (m, 4 H), 1.59 (d, $J=6.8$ Hz, 3 H), 1.78–1.90 (m, 1 H), 1.83 (s, 3 H), 1.98–2.10 (m, 2 H), 2.12–2.27 (m, 1 H), 2.34 (s, 3 H), 3.01 (dd, $J=10.4$, 5.3 Hz, 1 H), 3.19 (dd, $J=8.6$, 6.1 Hz, 1 H), 3.58 (dd, $J=10.3$, 3.3 Hz, 1 H), 3.82 (br dd, $J=3.3$, 0.6 Hz, 1 H), 4.11 (dd, $J=10.3$, 5.6 Hz, 1 H), 4.43 (br dd, $J=9.7$, 0.6 Hz, 1 H), 4.52 (dq, $J=6.8$, 3.1 Hz, 1 H), 4.62 (dd, $J=9.7$, 3.1 Hz, 1 H), 5.25 (d, $J=5.6$ Hz, 1 H), 7.35 (ddd, $J=8.1$, 7.2, 1.2 Hz, 1 H), 7.45 (ddd, $J=8.1$, 7.2, 1.2 Hz, 1 H), 7.81–7.89 (m, 2 H).

7(S)-7-(1H-Benzo[d]imidazol-2-ylthio)-7-deoxylincomycin (4). Compound **1** (240 mg, 0.39 mmol) and 1H-benzo[d]imidazole-2-thiol (87.9 mg, 0.59 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **4** (157.7 mg, 76%) as a colorless solid. $[\alpha]_D^{27}$ +91.9° (c 1.06, MeOH); ESI-MS (m/z) 539 (M+H)⁺ as C₂₅H₃₈N₄O₅S₂; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₅H₃₈N₄O₅S₂: 539.2362, found: 539.2361; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.87–0.97 (m, 3 H), 1.25–1.38 (m, 4 H), 1.47 (d, $J=7.1$ Hz, 3 H), 1.72–1.83 (m, 1 H), 1.90–2.05 (m, 2 H), 1.93 (s, 3 H), 2.14–2.28 (m, 1 H), 2.24 (s, 3 H), 3.01 (dd, $J=10.3$, 5.4 Hz, 1 H), 3.13 (dd, $J=8.6$, 6.2 Hz, 1 H), 3.60 (dd, $J=10.3$, 3.4 Hz, 1 H), 3.84 (br d, $J=3.4$ Hz, 1 H), 4.12 (dd, $J=10.3$, 5.6 Hz, 1 H), 4.16 (dq, $J=7.1$, 3.2 Hz, 1 H), 4.43–4.49 (m, 1 H), 4.51 (dd, $J=9.5$, 3.2 Hz, 1 H), 5.24 (d, $J=5.6$ Hz, 1 H), 7.19–7.26 (m, 2 H), 7.50 (br s, 2 H).

7(S)-S-(7-Deoxylincomycin-7-yl)benzothioate (5). To a solution of compound **1** (500 mg, 0.8 mmol) in toluene (5 ml) at 0 °C were added triphenylphosphine (316 mg, 1.20 mmol), diethylazodicarboxylate (0.22 ml, 1.20 mmol) and benzothioic S-acid (172 mg, 1.24 mmol) and stirred at RT for 3 h. The mixture was diluted with 2 N HCl (2 ml) and concentrated under reduced pressure. The resulting residue was dissolved by water and washed with diethyl ether. The mixture was added NaHCO₃ (150 mg), then extracted with ethyl acetate, washed with water, dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by preparative TLC (CHCl₃/CH₃OH/28% aq NH₄OH=9/1/0.1) to obtain the title compound as a colorless solid (100 mg, 24%). ESI-MS (m/z) 527 (M+H)⁺ as C₂₅H₃₈N₂O₆S₂; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.85–0.98 (m, 3 H), 1.23–1.39 (m, 4 H), 1.44 (d, $J=6.8$ Hz, 3 H), 1.83–1.92 (m, 1 H), 1.88 (s, 3 H), 1.98–2.07 (m, 1 H), 2.07–2.15 (m, 1 H), 2.15–2.28 (m, 1 H), 2.43 (s, 3 H), 3.07 (dd, $J=10.5$, 5.1 Hz, 1 H), 3.27 (dd, $J=8.3$, 5.8 Hz, 1 H), 3.55 (dd, $J=10.2$, 3.3 Hz, 1 H), 3.78–3.85 (m, 1 H), 4.11 (dd, $J=10.2$, 5.7 Hz, 1 H), 4.23–4.32 (m, 2 H), 4.58 (dd, $J=9.7$, 3.2 Hz, 1 H), 5.25 (d, $J=5.7$ Hz, 1 H), 7.42–7.55 (m, 2 H), 7.59–7.68 (m, 1 H), 7.91–8.00 (m, 2 H).

7(S)-7-Deoxy-7-phenylthiolincomycin (6). To a solution of compound **1** (1.0 g, 1.6 mmol) in THF (15 ml) at 0 °C were added tributylphosphine (971 mg, 4.8 mmol), diethylazodicarboxylate (0.59 ml, 3.2 mmol) and 1,2-diphenyldisulfide (530 mg, 2.4 mmol) and stirred at RT for 24 h. The mixture was diluted with 2 N HCl (1 ml) and stirred at RT for 30 min and then concentrated under reduced pressure. The resulting residue was dissolved by water and washed with diethyl ether. The mixture was added to NaHCO₃ (150 mg), then extracted with ethyl acetate, washed with water, dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (CHCl₃/CH₃OH/28% aq NH₄OH=20/1/0.1 → 9/1/0.1) to obtain the title compound as a colorless solid (724.0 mg, 91%). $[\alpha]_D^{26}$ +111.1° (c 0.63, MeOH); ESI-MS (m/z) 499 (M+H)⁺ as C₂₄H₃₈N₂O₅S₂; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₄H₃₈N₂O₅S₂: 499.2300, found: 499.2304; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.88–0.97 (m, 3 H), 1.29

(d, $J=6.9$ Hz, 3 H), 1.31–1.41 (m, 4 H), 1.80–1.90 (m, 1 H), 1.95–2.04 (m, 1 H), 2.00 (s, 3 H), 2.05–2.25 (m, 2 H), 2.39 (s, 3 H), 2.98 (dd, $J=10.7$, 4.6 Hz, 1 H), 3.24 (dd, $J=8.3$, 5.9 Hz, 1 H), 3.58 (dd, $J=10.2$, 3.3 Hz, 1 H), 3.74 (br d, $J=3.3$ Hz, 1 H), 3.86 (qd, $J=6.9$, 2.6 Hz, 1 H), 4.10 (dd, $J=10.2$, 5.6 Hz, 1 H), 4.35 (dd, $J=9.7$, 0.6 Hz, 1 H), 4.41 (dd, $J=9.7$, 2.6 Hz, 1 H), 5.26 (d, $J=5.6$ Hz, 1 H), 7.22–7.28 (m, 1 H), 7.29–7.36 (m, 2 H), 7.40–7.46 (m, 2 H).

7(S)-7-Deoxy-7-(pyridin-4-ylthio)lincomycin (7). Compound **1** (100 mg, 0.16 mmol) and pyridine-4-thiol (27.6 mg, 0.25 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **7** (99.8 mg, 40%) as a colorless solid. $[\alpha]_D^{26}$ +88.3° (c 0.94, MeOH); ESI-MS (m/z) 500 (M+H)⁺ as C₂₃H₃₇N₃O₅S₂; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₃H₃₇N₃O₅S₂: 500.2253, found: 500.2259; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.87–0.98 (m, 3 H), 1.28–1.40 (m, 4 H), 1.46 (d, $J=7.0$ Hz, 3 H), 1.78 (s, 3 H), 1.79–1.89 (m, 1 H), 1.96–2.11 (m, 2 H), 2.13–2.25 (m, 1 H), 2.39 (s, 3 H), 2.98 (dd, $J=10.5$, 4.9 Hz, 1 H), 3.24 (dd, $J=8.4$, 5.9 Hz, 1 H), 3.57 (dd, $J=10.2$, 3.2 Hz, 1 H), 3.80 (br dd, $J=3.2$, 0.8 Hz, 1 H), 4.09 (dd, $J=10.2$, 5.6 Hz, 1 H), 4.11 (dq, $J=7.0$, 2.9 Hz, 1 H), 4.37 (br dd, $J=9.6$, 0.8 Hz, 1 H), 4.59 (dd, $J=9.6$, 2.9 Hz, 1 H), 5.23 (d, $J=5.6$ Hz, 1 H), 7.33–7.39 (m, 2 H), 8.29–8.35 (m, 2 H).

7(S)-7-Deoxy-7-(pyrimidin-4-ylthio)lincomycin (8). Compound **1** (100 mg, 0.16 mmol) and pyrimidine-4-thiol (27.9 mg, 0.25 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **8** (124 mg, 50%) as a colorless solid. $[\alpha]_D^{27}$ +85.0° (c 1.53, MeOH); ESI-MS (m/z) 501 (M+H)⁺ as C₂₂H₃₆N₄O₅S₂; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₂H₃₆N₄O₅S₂: 501.2205, found: 501.2208; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.87–0.98 (m, 3 H), 1.26–1.41 (m, 4 H), 1.48 (d, $J=6.7$ Hz, 3 H), 1.79 (s, 3 H), 1.79–1.90 (m, 1 H), 1.97–2.12 (m, 2 H), 2.13–2.26 (m, 1 H), 2.37 (s, 3 H), 3.01 (dd, $J=10.5$, 5.1 Hz, 1 H), 3.24 (dd, $J=8.4$, 6.0 Hz, 1 H), 3.55 (dd, $J=10.2$, 3.2 Hz, 1 H), 3.80 (br dd, $J=3.2$, 0.7 Hz, 1 H), 4.10 (dd, $J=10.2$, 5.6 Hz, 1 H), 4.34 (br dd, $J=9.5$ Hz, 0.7, 1 H), 4.49–4.60 (m, 2 H), 5.23 (d, $J=5.6$ Hz, 1 H), 7.40 (dd, $J=5.6$, 1.5 Hz, 1 H), 8.36 (br dd, $J=5.6$, 0.4 Hz, 1 H), 8.87 (br dd, $J=1.5$, 0.4 Hz, 1 H).

7(S)-7-Deoxy-7-(thiophen-2-ylthio)lincomycin (9). Compound **1** (240 mg, 0.39 mmol) and thiophene-2-thiol (100 mg, 0.86 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **9** (19.4 mg, 10%) as a colorless solid. $[\alpha]_D^{26}$ +140.7° (c 0.47, MeOH); ESI-MS (m/z) 505 (M+H)⁺ as C₂₂H₃₆N₂O₅S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₂H₃₆N₂O₅S₃: 505.1865, found: 505.1863; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.89–0.98 (m, 3 H), 1.27 (d, $J=7.1$ Hz, 3H), 1.30–1.41 (m, 4 H), 1.78–1.89 (m, 1 H), 1.92–2.00 (m, 1 H), 2.01–2.08 (m, 1 H), 2.08–2.19 (m, 1 H), 2.21 (s, 3 H), 2.33 (s, 3 H), 2.96 (dd, $J=10.7$, 4.6 Hz, 1 H), 3.20 (dd, $J=8.1$, 5.6 Hz, 1 H), 3.54–3.65 (m, 2 H), 3.73 (br d, $J=2.8$ Hz, 1 H), 4.11 (dd, $J=10.3$, 5.6 Hz, 1 H), 4.34 (dd, $J=9.8$, 2.9 Hz, 1 H), 4.37–4.43 (m, 1 H), 5.29 (d, $J=5.6$ Hz, 1 H), 7.06 (dd, $J=5.4$, 3.5 Hz, 1H), 7.22 (dd, $J=3.5$, 1.2 Hz, 1 H), 7.55 (dd, $J=5.4$, 1.2 Hz, 1 H).

7(S)-7-Deoxy-7-(thiazol-2-ylthio)lincomycin (10). Compound **1** (240 mg, 0.39 mmol) and thiazole-2-thiol (43.0 mg, 0.37 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **10** (13.6 mg, 7%) as a colorless solid. $[\alpha]_D^{26}$ +109.5° (c 0.67, MeOH); ESI-MS (m/z) 506 (M+H)⁺ as C₂₁H₃₅N₃O₅S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₁H₃₅N₃O₅S₃: 506.1817, found: 506.1802; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.88–0.97 (m, 3 H), 1.27–1.40 (m, 4 H), 1.61 (d, $J=7.0$ Hz, 3 H), 1.79–1.92 (m, 1H), 1.95–2.14 (m, 2H), 2.01 (s, 3H), 2.15–2.29 (m, 1 H), 2.35 (s, 3 H), 3.01 (dd, $J=10.4$, 5.3 Hz, 1 H), 3.26 (dd, $J=8.6$, 6.1 Hz, 1 H), 3.57 (dd, $J=10.3$, 3.3 Hz, 1 H), 3.78 (br d, $J=3.3$ Hz, 1 H), 4.10 (dd, $J=10.3$, 5.6 Hz, 1 H), 4.13 (dq, $J=7.0$, 3.1 Hz, 1 H), 4.39 (br d, $J=9.8$ Hz, 1 H), 4.51 (dd, $J=9.8$, 3.1 Hz, 1 H), 5.25 (d, $J=5.6$ Hz, 1 H), 7.54 (d, $J=3.5$ Hz, 1 H), 7.73 (d, $J=3.5$ Hz, 1 H).

7(S)-7-Deoxy-7-(1,3,4-thiadiazol-2-ylthio)lincomycin (11). Compound **1** (240 mg, 0.39 mmol) and 1,3,4-thiadiazole-2-thiol (80.0 mg, 0.68 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **11** (121.0 mg, 62%) as a colorless solid. $[\alpha]_D^{27}$ +100.2° (c 2.03,

MeOH); ESI-MS (m/z) 507 (M+H)⁺ as C₂₀H₃₄N₄O₅S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₀H₃₄N₄O₅S₃: 507.1770, found: 507.1773; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.87–0.98 (m, 3 H), 1.26–1.42 (m, 4 H), 1.53 (d, J = 6.8 Hz, 3 H), 1.80–1.90 (m, 1 H), 1.93 (s, 3 H), 1.97–2.12 (m, 2 H), 2.15–2.28 (m, 1 H), 2.38 (s, 3 H), 3.03 (dd, J = 10.5, 5.1 Hz, 1 H), 3.27 (dd, J = 8.4, 6.1 Hz, 1 H), 3.57 (dd, J = 10.2, 3.2 Hz, 1 H), 3.81 (br dd, J = 3.2, 0.8 Hz, 1 H), 4.11 (dd, J = 10.2, 5.6 Hz, 1 H), 4.37–4.46 (m, 2 H), 4.60 (dd, J = 9.8, 3.2 Hz, 1 H), 5.26 (d, J = 5.6 Hz, 1 H), 9.37 (s, 1H).

7(S)-7-(5-Chlorobenzo[d]thiazol-2-ylthio)-7-deoxylincomycin (12). Compound **1** (160 mg, 0.26 mmol) and 5-chlorobenzo[d]thiazole-2-thiol (160 mg, 0.79 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **12** (110.6 mg, 73%) as a colorless solid. [α]_D²⁶ +52.8° (c 0.25, MeOH); ESI-MS (m/z) 590 (M+H)⁺ as C₂₅H₃₆ClN₃O₅S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₅H₃₆ClN₃O₅S₃: 590.1584, found: 590.1581; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.87–0.96 (m, 3 H), 1.27–1.39 (m, 4 H), 1.59 (d, J = 7.0 Hz, 3 H), 1.78–1.88 (m, 1 H), 1.86 (s, 3 H), 1.97–2.08 (m, 2 H), 2.13–2.27 (m, 1 H), 2.32 (s, 3 H), 2.98 (dd, J = 10.5, 5.2 Hz, 1 H), 3.20 (dd, J = 8.5, 6.2 Hz, 1 H), 3.56 (dd, J = 10.2, 3.2 Hz, 1 H), 3.82 (br dd, J = 3.2, 0.9 Hz, 1 H), 4.10 (dd, J = 10.2, 5.6 Hz, 1 H), 4.41 (br dd, J = 9.7, 0.9 Hz, 1 H), 4.53 (dq, J = 7.0, 3.3 Hz, 1 H), 4.61 (dd, J = 9.7, 3.3 Hz, 1 H), 5.24 (d, J = 5.6 Hz, 1 H), 7.35 (dd, J = 8.6, 2.1 Hz, 1 H), 7.81–7.87 (m, 2 H).

7(S)-7-Deoxyl-7-(thiazolo[5,4-*c*]pyridin-2-ylthio)lincomycin (13). Compound **1** (320 mg, 0.51 mmol) and thiazolo[5,4-*c*]pyridine-2-thiol (250 mg, 1.49 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **13** (205.9 mg, 72%) as a colorless solid. [α]_D²⁶ +67.1° (c 0.50, MeOH); ESI-MS (m/z) 505 (M+H)⁺ as C₂₄H₃₆N₄O₅S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₄H₃₆N₄O₅S₃: 557.1926, found: 557.1924; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.87–0.98 (m, 3 H), 1.27–1.41 (m, 4 H), 1.62 (d, J = 6.7 Hz, 3 H), 1.82 (s, 3 H), 1.80–1.91 (m, 1 H), 1.98–2.13 (m, 2 H), 2.15–2.28 (m, 1 H), 2.38 (s, 3 H), 3.03 (dd, J = 10.4, 5.3 Hz, 1 H), 3.24 (dd, J = 8.6, 6.2 Hz, 1 H), 3.56 (dd, J = 10.3, 3.2 Hz, 1 H), 3.83 (br dd, J = 3.2, 0.73 Hz, 1 H), 4.11 (dd, J = 10.3, 5.6 Hz, 1 H), 4.43 (br dd, J = 9.5, 0.73 Hz, 1 H), 4.63–4.72 (m, 2 H), 5.24 (d, J = 5.6 Hz, 1 H), 7.82 (dd, J = 5.6, 0.9 Hz, 1 H), 8.51 (d, J = 5.6 Hz, 1 H), 9.06 (d, J = 0.9 Hz, 1 H).

7(S)-7-(6-Cyanobenzo[d]thiazol-2-ylthio)-7-deoxylincomycin (14). Compound **1** (160 mg, 0.26 mmol) and 6-cyanobenzo[d]thiazole-2-thiol (55 mg, 0.29 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **14** (98.4 mg, 66%) as a colorless solid. [α]_D²⁶ +73.0° (c 1.10, MeOH); ESI-MS (m/z) 581 (M+H)⁺ as C₂₆H₃₆N₄O₅S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₆H₃₆N₄O₅S₃: 581.1926, found: 581.1926; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.87–0.97 (m, 3 H), 1.25–1.41 (m, 4 H), 1.61 (d, J = 6.6 Hz, 3 H), 1.78–1.89 (m, 1 H), 1.82 (s, 3 H), 1.98–2.09 (m, 2 H), 2.14–2.26 (m, 1 H), 2.35 (s, 3 H), 2.98 (dd, J = 10.4, 5.1 Hz, 1 H), 3.21 (dd, J = 8.5, 6.2 Hz, 1 H), 3.56 (dd, J = 10.2, 3.2 Hz, 1 H), 3.82 (br dd, J = 3.2, 0.8 Hz, 1 H), 4.10 (dd, J = 10.2, 5.7 Hz, 1 H), 4.42 (br dd, J = 9.5, 0.8 Hz, 1 H), 4.60–4.70 (m, 2 H), 5.24 (d, J = 5.7 Hz, 1 H), 7.76 (dd, J = 8.5, 1.7 Hz, 1 H), 7.94 (br dd, J = 8.5, 0.6 Hz, 1 H), 8.32 (br dd, J = 1.7, 0.6 Hz, 1 H).

7(S)-7-Deoxy-7-(6-nitrobenzo[d]thiazol-2-ylthio)lincomycin (15). Compound **1** (240 mg, 0.39 mmol) and 6-nitrobenzo[d]thiazole-2-thiol (180 mg, 0.85 mmol) were treated in toluene (5 ml) according to the similar procedure as described for the preparation of **2** to afford **15** (152.7 mg, 66%) as a colorless solid. [α]_D²⁹ +67.6° (c 0.60, MeOH); ESI-MS (m/z) 601 (M+H)⁺ as C₂₅H₃₆N₄O₇S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₅H₃₆N₄O₇S₃: 601.1824, found: 601.1827; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.87–0.98 (m, 3 H), 1.28–1.40 (m, 4 H), 1.62 (d, J = 6.7 Hz, 3 H), 1.77–1.89 (m, 1 H), 1.83 (s, 3 H), 1.98–2.09 (m, 2 H), 2.14–2.27 (m, 1 H), 2.36 (s, 3 H), 2.99 (dd, J = 10.5, 5.1 Hz, 1 H), 3.22 (dd, J = 8.5, 6.2 Hz, 1 H), 3.56 (dd, J = 10.1, 3.2 Hz, 1 H), 3.83 (br dd, J = 3.2, 0.8 Hz, 1 H), 4.10 (dd, J = 10.1, 5.6 Hz, 1 H), 4.43 (br dd, J = 9.5, 0.8 Hz, 1 H), 4.62–4.71 (m, 2 H), 5.24 (d, J = 5.6 Hz, 1 H), 7.95 (d, J = 9.0, 1 H), 8.32 (dd, J = 9.0, 2.3, Hz, 1 H), 8.85 (d, J = 2.3 Hz, 1 H).

7(S)-7-(6-Aminobenzo[d]thiazol-2-ylthio)-7-deoxylincomycin (16). Compound **1** (630 mg, 1.01 mmol) and 6-aminobenzo[d]thiazole-2-thiol (300 mg, 1.65

mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **16** (386.7 mg, 67%) as a colorless solid. [α]_D²⁶ +89.0° (c 1.11, MeOH); ESI-MS (m/z) 571 (M+H)⁺ as C₂₅H₃₈N₄O₅S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₅H₃₈N₄O₅S₃: 571.2083, found: 571.2075; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.86–0.97 (m, 3 H), 1.26–1.38 (m, 4 H), 1.52 (d, J = 7.0 Hz, 3 H), 1.75–1.89 (m, 1 H), 1.94 (s, 3 H), 1.95–2.11 (m, 2 H), 2.12–2.25 (m, 1 H), 2.32 (s, 3 H), 3.03 (dd, J = 10.4, 5.3 Hz, 1 H), 3.17 (dd, J = 8.6, 6.2 Hz, 1 H), 3.58 (dd, J = 10.2, 3.2 Hz, 1 H), 3.81 (br dd, J = 3.2, 0.7 Hz, 1 H), 4.10 (dd, J = 10.2, 5.5 Hz, 1 H), 4.27 (dq, J = 7.0, 3.1 Hz, 1 H), 4.43 (br dd, J = 9.8, 0.7 Hz, 1 H), 4.55 (dd, J = 9.8, 3.1 Hz, 1 H), 5.25 (d, J = 5.5 Hz, 1 H), 6.85 (dd, J = 8.7, 2.3 Hz, 1 H), 7.08 (dd, J = 2.3, 0.25 Hz, 1 H), 7.59 (dd, J = 8.7, 0.25 Hz, 1 H).

7(S)-7-Deoxy-7-(5-nitrobenzo[d]thiazol-2-ylthio)lincomycin (17). Compound **1** (320 mg, 0.51 mmol) and 5-nitrobenzo[d]thiazole-2-thiol (120 mg, 0.57 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **17** (138.8 mg, 45%) as a colorless solid. [α]_D²⁶ +58.2° (c 0.81, MeOH); ESI-MS (m/z) 601 (M+H)⁺ as C₂₅H₃₆N₄O₇S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₅H₃₆N₄O₇S₃: 601.1824, found: 601.1826; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.88–0.97 (m, 3 H), 1.30–1.40 (m, 4 H), 1.62 (d, J = 6.8 Hz, 3 H), 1.80–1.90 (m, 1 H), 1.85 (s, 3 H), 1.99–2.10 (m, 2 H), 2.16–2.30 (m, 1 H), 2.36 (s, 3 H), 3.01 (dd, J = 10.3, 5.3 Hz, 1 H), 3.25 (dd, J = 8.7, 6.2 Hz, 1 H), 3.57 (dd, J = 10.3, 3.2 Hz, 1 H), 3.84 (br dd, J = 3.2, 0.8 Hz, 1 H), 4.11 (dd, J = 10.3, 5.7 Hz, 1 H), 4.42 (br dd, J = 9.6, 0.8 Hz, 1 H), 4.61 (dd, J = 6.8, 3.3 Hz, 1 H), 4.65 (dd, J = 9.6, 3.3 Hz, 1 H), 5.25 (d, J = 5.7 Hz, 1 H), 8.08 (d, J = 8.8, 1 H), 8.21 (dd, J = 8.8, 2.2, Hz, 1 H), 8.64 (br dd, J = 2.2, 0.24 Hz, 1 H).

7(S)-7-Deoxy-7-(2-nitrophenylthio)lincomycin (18). Compound **1** (200 mg, 0.32 mmol) and 1,2-bis(2-nitrophenyl)disulfide (148.5 mg, 0.48 mmol) were treated according to the similar procedure as described for the preparation of **6** to afford **18** (40.1 mg, 23%) as a colorless solid. [α]_D²⁷ +69.9° (c 0.91, MeOH); ESI-MS (m/z) 544 (M+H)⁺ as C₂₄H₃₇N₃O₇S₂; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₄H₃₇N₃O₇S₂: 544.2151, found: 544.2157; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.86–0.99 (m, 3 H), 1.29–1.38 (m, 4 H), 1.38 (d, J = 6.8 Hz, 3 H), 1.79 (s, 3 H), 1.81–1.92 (m, 1 H), 1.98–2.14 (m, 2 H), 2.14–2.26 (m, 1 H), 2.42 (s, 3 H), 3.01 (dd, J = 10.6, 4.8 Hz, 1 H), 3.28 (dd, J = 8.7, 6.1 Hz, 1 H), 3.57 (dd, J = 10.3, 3.2 Hz, 1 H), 3.79 (br dd, J = 3.2, 0.8 Hz, 1 H), 4.06 (dq, J = 6.8, 2.7 Hz, 1 H), 4.08 (dd, J = 10.3, 5.6 Hz, 1 H), 4.37 (br dd, J = 9.8, 0.8 Hz, 1 H), 4.57 (dd, J = 9.8, 2.7 Hz, 1 H), 5.21 (d, J = 5.6 Hz, 1 H), 7.38 (ddd, J = 8.3, 7.2, 1.2 Hz, 1 H), 7.64 (ddd, J = 8.2, 7.2, 1.4 Hz, 1 H), 7.70 (br dd, J = 8.2, 1.2 Hz, 1 H), 8.05 (dd, J = 8.3, 1.4 Hz, 1 H).

7(S)-7-Deoxy-7-(3-nitrophenylthio)lincomycin (19). Compound **1** (1.0 g, 1.6 mmol) and 1,2-bis(3-nitrophenyl)disulfide (742 mg, 2.4 mmol) were treated according to the similar procedure as described for the preparation of **6** to afford **19** (366.5 mg, 42%) as a colorless solid. [α]_D²⁹ +75.2° (c 0.48, MeOH); ESI-MS (m/z) 544 (M+H)⁺ as C₂₄H₃₇N₃O₇S₂; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₄H₃₇N₃O₇S₂: 544.2151, found: 544.2148; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.89–0.97 (m, 3 H), 1.30–1.37 (m, 4 H), 1.38 (d, J = 6.8 Hz, 3 H), 1.80–1.91 (m, 1 H), 1.93 (s, 3 H), 1.97–2.12 (m, 2 H), 2.13–2.24 (m, 1 H), 2.40 (s, 3 H), 2.99 (dd, J = 10.6, 4.8 Hz, 1 H), 3.24 (dd, J = 8.3, 5.9 Hz, 1 H), 3.58 (dd, J = 10.2, 3.3 Hz, 1 H), 3.79 (br dd, J = 3.3, 0.8 Hz, 1 H), 3.99 (dq, J = 6.8, 2.8 Hz, 1 H), 4.10 (dd, J = 10.2, 5.6 Hz, 1 H), 4.36 (br dd, J = 9.5, 0.8 Hz, 1 H), 4.52 (dd, J = 9.5, 2.8 Hz, 1 H), 5.25 (d, J = 5.6 Hz, 1 H), 7.55–7.61 (m, 1 H), 7.81 (ddd, J = 7.9, 1.8, 0.9 Hz, 1 H), 8.09 (br ddd, J = 8.2, 2.2, 0.9 Hz, 1 H), 8.21–8.23 (m, 1 H).

7(S)-7-Deoxy-7-(4-nitrophenylthio)lincomycin (20). Compound **1** (200 mg, 0.32 mmol) and 1,2-bis(4-nitrophenyl)disulfide (148.5 mg, 0.48 mmol) were treated according to the similar procedure as described for the preparation of **6** to afford **20** (29.7 mg, 17%) as a colorless solid. [α]_D²⁶ +67.1° (c 0.29, MeOH); ESI-MS (m/z) 544 (M+H)⁺ as C₂₄H₃₇N₃O₇S₂; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₄H₃₇N₃O₇S₂: 544.2151, found: 544.2151; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.87–0.98 (m, 3 H), 1.30–1.39 (m, 4 H), 1.44 (d, J = 6.8 Hz, 3 H), 1.79–1.90 (m, 1 H), 1.81 (s, 3 H), 1.97–2.12 (m, 2 H), 2.13–2.26 (m, 1 H), 2.40 (s, 3 H), 3.00 (dd, J = 10.6, 5.0 Hz, 1 H), 3.25 (dd, J = 8.4, 6.1 Hz, 1 H), 3.57 (dd, J = 10.2, 3.2 Hz, 1 H), 3.80 (br dd, J = 3.2, 0.8 Hz, 1 H), 4.09 (dd, J = 10.2, 5.6 Hz, 1 H),

4.03–4.16 (m, 1 H), 4.37 (br dd, $J=9.7, 0.8$ Hz, 1 H), 4.58 (dd, $J=9.7, 2.9$ Hz, 1 H), 5.24 (d, $J=5.6$ Hz, 1 H), 7.51–7.57 (m, 2 H), 8.13–8.19 (m, 2 H).

7(S)-7-(5-Amino-1,3,4-thiadiazol-2-ylthio)-7-deoxylincomycin (21). To a solution of compound **1** (240 mg, 0.39 mmol) in THF (5 ml) at 0 °C were added triphenylphosphine (150 mg, 0.57 mmol), diethylazodicarboxylate (0.1 ml, 0.55 mmol) and *t*-butyl (5-mercapto-1,3,4-thiadiazol-2-yl)carbamate (130 mg, 0.56 mmol) and stirred at RT for 17 h. The mixture was concentrated under reduced pressure and diluted with saturated aqueous NaHCO₃ (10 ml), extracted with ethyl acetate, washed with water, dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate) to obtain 7(S)-7-(5-amino-1,3,4-thiadiazol-2-ylthio)-7-deoxy-2,3,4-tris-*O*-(trimethylsilyl)lincomycin as a colorless solid (271.3 mg, 84%). 7(S)-7-(5-Amino-1,3,4-thiadiazol-2-ylthio)-7-deoxy-2,3,4-tris-*O*-(trimethylsilyl)lincomycin (271.3 mg, 0.32 mmol) in 90% aqueous trifluoroacetic acid (5 ml) was kept at RT for 1 h. The mixture was concentrated under reduced pressure and diluted with saturated aqueous NaHCO₃ (10 ml) and then extracted with ethyl acetate, washed with water, dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by preparative TLC (CHCl₃/CH₃OH/28% aq NH₄OH = 5/1/0.1) to obtain the title compound as a colorless solid (111.4 mg, 66%). [α]_D²⁶ +139.8° (c 0.28, MeOH); ESI-MS (m/z) 520 (M+H)⁺ as C₂₀H₃₅N₅O₅S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₀H₃₅N₅O₅S₃: 522.1879, found: 522.1877; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.87–0.97 (m, 3 H), 1.27–1.38 (m, 4 H), 1.41 (d, $J=7.0$ Hz, 3 H), 1.76–1.90 (m, 1 H), 1.92–2.07 (m, 2 H), 2.11 (s, 3 H), 2.14–2.26 (m, 1 H), 2.33 (s, 3 H), 2.97 (dd, $J=10.5, 4.9$ Hz, 1 H), 3.26 (dd, $J=8.6, 6.2$ Hz, 1 H), 3.57 (dd, $J=10.3, 3.2$ Hz, 1 H), 3.74–3.78 (m, 1 H), 3.94 (dq, $J=7.0, 2.9$ Hz, 1 H), 4.10 (dd, $J=10.3, 5.6$ Hz, 1 H), 4.39 (br dd, $J=9.8, 0.5$ Hz, 1 H), 4.45 (dd, $J=9.8, 2.9$ Hz, 1 H), 5.26 (d, $J=5.6$ Hz, 1 H). For the qualified analytical purpose, the above colorless solid was further purified by reverse-phase column chromatography (YMC triart C18, 20 × 250 mm, RT, 18.9 ml min⁻¹, 50 mM AcONH₄/CH₃CN = 70/30) and precipitated (MeOH/ethyl acetate) to obtain the highly purified title compound as a colorless solid.

7(S)-7-Deoxy-7-(5-methylamino-1,3,4-thiadiazol-2-ylthio)lincomycin (22). Compound **1** (240 mg, 0.39 mmol) and *t*-butyl (5-mercapto-1,3,4-thiadiazol-2-yl)(methyl)carbamate (100 mg, 0.40 mmol) were treated according to the similar procedure as described for the preparation of **21** to afford **22** (147.1 mg, 71% (two steps)) as a colorless solid. [α]_D²⁶ +118.4° (c 0.27, MeOH); ESI-MS (m/z) 534 (M+H)⁺ as C₂₁H₃₇N₅O₅S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₁H₃₇N₅O₅S₃: 536.2035, found: 536.2039; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.87–0.98 (m, 3 H), 1.27–1.38 (m, 4 H), 1.41 (d, $J=7.0$ Hz, 3 H), 1.77–1.88 (m, 1 H), 1.93–2.01 (m, 1 H), 2.04 (dd, $J=10.1, 8.7$ Hz, 1 H), 2.12 (s, 3 H), 2.14–2.26 (m, 1 H), 2.33 (s, 3H), 2.93–3.01 (m, 1 H), 2.97 (s, 3 H), 3.25 (dd, $J=8.4, 6.2$ Hz, 1 H), 3.57 (dd, $J=10.3, 3.2$ Hz, 1 H), 3.78 (br dd, $J=3.2, 0.7$ Hz, 1 H), 3.94 (dq, $J=7.0, 2.9$ Hz, 1 H), 4.10 (dd, $J=10.3, 5.6$ Hz, 1 H), 4.38 (br dd, $J=9.8, 0.7$ Hz, 1 H), 4.46 (dd, $J=9.8, 2.9$ Hz, 1 H), 5.26 (d, $J=5.6$ Hz, 1 H).

7(S)-7-Deoxy-7-(5-(4-(difluoromethylthio)phenylamino)-1,3,4-thiadiazol-2-ylthio)lincomycin (23). Compound **1** (240 mg, 0.39 mmol) and 5-(4-(difluoromethylthio)phenylamino)-1,3,4-thiadiazole-2-thiol (120 mg, 0.41 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **23** (99.5 mg, 38%) as a colorless solid. [α]_D²⁴ +96.7° (c 0.78, MeOH); ESI-MS (m/z) 680 (M+H)⁺ as C₂₇H₃₉F₂N₅O₅S₄; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₇H₃₉F₂N₅O₅S₄: 680.1880, found: 680.1876; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.85–0.95 (m, 3 H), 1.26–1.39 (m, 4 H), 1.46 (d, $J=7.0$ Hz, 3 H), 1.78–1.88 (m, 1 H), 1.93–2.06 (m, 2 H), 2.12 (s, 3 H), 2.14–2.28 (m, 1 H), 2.33 (s, 3 H), 2.95–3.02 (m, 1 H), 3.26 (dd, $J=8.4, 6.2$ Hz, 1 H), 3.59 (dd, $J=10.3, 3.3$ Hz, 1 H), 3.79 (br d, $J=3.3$ Hz, 1 H), 4.02–4.10 (m, 1 H), 4.12 (dd, $J=10.3, 5.6$ Hz, 1 H), 4.41 (br d, $J=9.8$ Hz, 1 H), 4.49 (dd, $J=9.8, 3.6$ Hz, 1 H), 5.28 (d, $J=5.6$ Hz, 1 H), 7.00 (t, $J=56.8$ Hz, 1 H), 7.52–7.58 (m, 2 H), 7.64–7.70 (m, 2 H).

7(S)-7-(5-Carbamoyl-1,3,4-thiadiazol-2-ylthio)-7-deoxylincomycin (24). Compound **1** (240 mg, 0.39 mmol), and 5-mercapto-1,3,4-thiadiazole-2-carboxamide (95 mg, 0.59 mmol) were treated at 50 °C according to the similar

procedure as described for the preparation of **2** to afford **24** (67.8 mg, 32%) as a colorless solid. [α]_D²⁶ +83.6° (c 0.69, MeOH); ESI-MS (m/z) 550 (M+H)⁺ as C₂₁H₃₅N₅O₆S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₁H₃₅N₅O₆S₃: 550.1828, found: 550.1829; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.86–0.97 (m, 3 H), 1.30–1.41 (m, 4 H), 1.57 (d, $J=7.0$ Hz, 3 H), 1.78–1.89 (m, 1 H), 1.94 (s, 3 H), 1.97–2.09 (m, 2 H), 2.13–2.26 (m, 1 H), 2.37 (s, 3 H), 2.99 (dd, $J=10.5, 5.1$ Hz, 1 H), 3.24 (dd, $J=8.5, 6.2$ Hz, 1 H), 3.56 (dd, $J=10.3, 3.2$ Hz, 1 H), 3.81 (br dd, $J=3.2, 0.8$ Hz, 1 H), 4.10 (dd, $J=10.3, 5.6$ Hz, 1 H), 4.40 (br dd, $J=9.7, 0.8$ Hz, 1 H), 4.49 (dq, $J=7.0, 3.1$ Hz, 1 H), 4.61 (dd, $J=9.7, 3.1$ Hz, 1 H), 5.25 (d, $J=5.6$ Hz, 1 H). For the qualified analytical purpose, the above colorless solid was further purified by reverse-phase column chromatography (YMC triart C18, 20 × 250 mm, RT, 18.9 ml min⁻¹, 50 mM AcONH₄/CH₃CN = 50/50) to obtain the highly purified title compound as a colorless solid.

7(S)-7-Deoxy-7-(5-methylcarbamoyl-1,3,4-thiadiazol-2-ylthio)lincomycin (25). Compound **1** (240 mg, 0.39 mmol) and 5-mercapto-*N*-methyl-1,3,4-thiadiazole-2-carboxamide (100 mg, 0.57 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **25** (26.1 mg, 12%) as a colorless solid. [α]_D²⁶ +81.8° (c 1.29, MeOH); ESI-MS (m/z) 564 (M+H)⁺ as C₂₂H₃₇N₅O₆S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₂H₃₇N₅O₆S₃: 564.1984, found: 564.1977; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.85–0.98 (m, 3 H), 1.26–1.42 (m, 4 H), 1.56 (d, $J=6.8$ Hz, 3 H), 1.78–1.89 (m, 1 H), 1.94 (s, 3 H), 1.97–2.12 (m, 2 H), 2.13–2.27 (m, 1 H), 2.38 (s, 3 H), 2.94 (s, 3 H), 3.00 (dd, $J=10.5, 5.1$ Hz, 1 H), 3.25 (dd, $J=8.6, 6.1$ Hz, 1 H), 3.56 (dd, $J=10.2, 3.2$ Hz, 1 H), 3.79–3.84 (m, 1 H), 4.10 (dd, $J=10.2, 5.6$ Hz, 1 H), 4.40 (br d, $J=9.8$ Hz, 1 H), 4.48 (dq, $J=6.8, 3.1$ Hz, 1 H), 4.62 (dd, $J=9.8, 3.1$ Hz, 1 H), 5.25 (d, $J=5.6$ Hz, 1 H). For the qualified analytical purpose, the above colorless solid was further purified by reverse-phase column chromatography (YMC triart C18, 20 × 250 mm, RT, 11.3 ml min⁻¹, 0.1% aq. TFA/CH₃CN = 60/40) to obtain the highly purified title compound as a colorless solid.

7(S)-7-Deoxy-7-(5-(2-(pyrrolidin-1-yl)ethylcarbamoyl)-1,3,4-thiadiazol-2-ylthio)lincomycin (26). Compound **1** (240 mg, 0.39 mmol) and 5-(2-(pyrrolidin-1-yl)ethylcarbamoyl)-1,3,4-thiadiazole-2-thiol (130 mg, 0.50 mmol) were treated according to the similar procedure as described for the preparation of **2** to afford **26** (69.8 mg, 28%) as a colorless solid. [α]_D²⁷ +78.0° (c 0.54, MeOH); ESI-MS (m/z) 647 (M+H)⁺ as C₂₇H₄₆N₆O₆S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₇H₄₆N₆O₆S₃: 647.2719, found: 647.2716; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.87–0.97 (m, 3 H), 1.28–1.39 (m, 4 H), 1.56 (d, $J=7.0$ Hz, 3 H), 1.77–1.90 (m, 5 H), 1.94 (s, 3 H), 1.97–2.12 (m, 2 H), 2.13–2.26 (m, 1 H), 2.37 (s, 3 H), 2.57–2.67 (m, 4 H), 2.73 (t, $J=6.7$ Hz, 2 H), 2.98 (dd, $J=10.5, 5.1$ Hz, 1 H), 3.24 (dd, $J=8.5, 6.2$ Hz, 1 H), 3.51–3.60 (m, 3 H), 3.81 (br dd, $J=3.2, 0.8$ Hz, 1 H), 4.10 (dd, $J=10.2, 5.6$ Hz, 1 H), 4.40 (br dd, $J=9.7, 0.8$ Hz, 1 H), 4.48 (dq, $J=7.0, 3.1$ Hz, 1 H), 4.62 (dd, $J=9.7, 3.1$ Hz, 1 H), 5.25 (d, $J=5.6$ Hz, 1 H).

7(S)-7-Deoxy-7-(5-phenyl-1,3,4-thiadiazol-2-ylthio)lincomycin (27). Compound **1** (240 mg, 0.39 mmol) and 5-phenyl-1,3,4-thiadiazole-2-thiol (100 mg, 0.51 mmol) in toluene (5 ml) were treated according to the similar procedure as described for the preparation of **2** to afford **27** (46.5 mg, 21%) as a colorless solid. [α]_D²⁷ +157.2° (c 1.47, CHCl₃); ESI-MS (m/z) 583 (M+H)⁺ as C₂₆H₃₈N₄O₅S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₆H₃₈N₄O₅S₃: 583.2083, found: 583.2086; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.86–0.97 (m, 3 H), 1.26–1.38 (m, 4 H), 1.57 (d, $J=7.0$ Hz, 3 H), 1.79–1.91 (m, 1 H), 1.96–2.11 (m, 2 H), 2.03 (s, 3 H), 2.15–2.28 (m, 1 H), 2.38 (s, 3 H), 3.03 (dd, $J=10.5, 5.1$ Hz, 1 H), 3.27 (dd, $J=8.5, 6.1$ Hz, 1 H), 3.58 (dd, $J=10.3, 3.2$ Hz, 1 H), 3.83 (br dd, $J=3.2, 0.7$ Hz, 1 H), 4.12 (dd, $J=10.3, 5.6$ Hz, 1 H), 4.36–4.46 (m, 2 H), 4.60 (dd, $J=9.7, 3.2$ Hz, 1 H), 5.27 (d, $J=5.6$ Hz, 1 H), 7.49–7.58 (m, 3 H), 7.89–7.95 (m, 2 H).

7(S)-7-Allylthio-7-deoxylincomycin (28). To a solution of compound **5** (83.2 mg, 0.16 mmol) in methanol (1 ml) were added allyl iodide (26.5 mg, 0.16 mmol) and 28% sodium methoxide in methanol (0.56 ml) and stirred at RT for 14 h. The mixture was diluted with 1 N HCl (1 ml) and concentrated under reduced pressure. The resulting residue was dissolved by water and washed with diethyl ether. The mixture was added to NaHCO₃ (150 mg)

and then extracted with ethyl acetate, washed with water, dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by preparative TLC ($\text{CHCl}_3/\text{CH}_3\text{OH}/28\%$ aq $\text{NH}_4\text{OH}=20/1/0.1$) to obtain the title compound as a colorless solid (22.0 mg, 30%). $[\alpha]_{\text{D}}^{25} +115.3^\circ$ (*c* 0.34, MeOH); ESI-MS (*m/z*) 463 ($\text{M}+\text{H}^+$) as $\text{C}_{21}\text{H}_{38}\text{N}_2\text{O}_5\text{S}_2$; TOF-ESI-HRMS ($\text{M}+\text{H}^+$)⁺ calcd for $\text{C}_{21}\text{H}_{38}\text{N}_2\text{O}_5\text{S}_2$: 463.2300, found: 463.2295; ^1H NMR (400 MHz, methanol- d_4) δ 0.89–0.96 (m, 3 H), 1.27–1.40 (m, 4 H), 1.31 (d, *J* = 7.0 Hz, 3 H), 1.82–1.92 (m, 1 H), 2.00 (ddd, *J* = 12.5, 7.6, 4.7 Hz, 1 H), 2.08–2.27 (m, 2 H), 2.19 (s, 3 H), 2.44 (s, 3 H), 3.05 (dd, *J* = 10.5, 4.4 Hz, 1 H), 3.23–3.39 (m, 4 H), 3.56 (dd, *J* = 10.2, 3.3 Hz, 1 H), 3.67–3.74 (m, 1 H), 4.09 (dd, *J* = 10.2, 5.6 Hz, 1 H), 4.21 (br d, *J* = 9.8 Hz, 1 H), 4.27 (dd, *J* = 9.8, 2.4 Hz, 1 H), 5.04–5.11 (m, 1 H), 5.19 (dq, *J* = 17.0, 1.4 Hz, 1 H), 5.25 (d, *J* = 5.6 Hz, 1 H), 5.85 (ddt, *J* = 17.0, 10.7, 1.0 Hz, 1 H).

7(S)-7-(5-Aminobenzo[d]thiazol-2-ylthio)-7-deoxylincomycin (29). To a solution of compound **17** (75.0 mg, 0.12 mmol) in ethanol (3.0 ml) were added $\text{SnCl}_2\cdot\text{H}_2\text{O}$ (140 mg, 0.62 mmol) and NaBH_4 (6.5 mg, 0.17 mmol) and stirred at RT for 3 h. The mixture was concentrated under reduced pressure. The resulting residue was dissolved by ethyl acetate, washed with water, dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by preparative TLC ($\text{CHCl}_3/\text{CH}_3\text{OH}/28\%$ aq $\text{NH}_4\text{OH}=20/1/0.1$) to obtain the title compound as a colorless solid (34.2 mg, 48%). $[\alpha]_{\text{D}}^{25} +59.4^\circ$ (*c* 0.58, MeOH); ESI-MS (*m/z*) 571 ($\text{M}+\text{H}^+$)⁺ as $\text{C}_{25}\text{H}_{38}\text{N}_4\text{O}_5\text{S}_3$; TOF-ESI-HRMS ($\text{M}+\text{H}^+$)⁺ calcd for $\text{C}_{25}\text{H}_{38}\text{N}_4\text{O}_5\text{S}_3$: 571.2083, found: 571.2090; ^1H NMR (400 MHz, methanol- d_4) δ 0.87–0.96 (m, 3 H), 1.25–1.39 (m, 4 H), 1.55 (d, *J* = 7.0 Hz, 3 H), 1.76–1.86 (m, 1 H), 1.88 (s, 3 H), 1.96–2.08 (m, 2 H), 2.12–2.24 (m, 1 H), 2.32 (s, 3 H), 2.99 (dd, *J* = 10.5, 5.2 Hz, 1 H), 3.18 (dd, *J* = 8.5, 6.2 Hz, 1 H), 3.57 (dd, *J* = 10.1, 3.2 Hz, 1H), 3.81 (br dd, *J* = 3.2, 0.7 Hz, 1 H), 4.10 (dd, *J* = 10.1, 5.6 Hz, 1 H), 4.39 (dq, *J* = 7.0, 3.1 Hz, 1 H), 4.42 (br dd, *J* = 9.7, 0.7 Hz, 1 H), 4.58 (dd, *J* = 9.7, 3.1 Hz, 1 H), 5.24 (d, *J* = 5.6 Hz, 1 H), 6.81 (dd, *J* = 8.6, 2.2 Hz, 1H), 7.16–7.18 (m, 1 H), 7.52 (dd, *J* = 8.6, 0.2 Hz, 1 H).

7(R)-7-(6-Aminobenzo[d]thiazol-2-ylthio)-7-deoxylincomycin (30). To a solution of clindamycin hydrochloride (460 mg, 1.0 mmol) in DMF (5 ml) were added K_2CO_3 (152 mg, 1.1 mmol) and 6-aminobenzo[d]thiazole-2-thiol (200 mg, 1.1 mmol) and stirred at 100 °C for 16 h. Then it was concentrated under reduced pressure. The residue was diluted with water and then extracted with ethyl acetate, washed with water, dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by preparative TLC ($\text{CHCl}_3/\text{CH}_3\text{OH}/28\%$ aq $\text{NH}_4\text{OH}=20/1/0.1$) to obtain the title compound as a colorless solid (233.3 mg, 41%). $[\alpha]_{\text{D}}^{27} +129.5^\circ$ (*c* 0.45, MeOH); ESI-MS (*m/z*) 571 ($\text{M}+\text{H}^+$)⁺ as $\text{C}_{25}\text{H}_{38}\text{N}_4\text{O}_5\text{S}_3$; TOF-ESI-HRMS ($\text{M}+\text{H}^+$)⁺ calcd for $\text{C}_{25}\text{H}_{38}\text{N}_4\text{O}_5\text{S}_3$: 571.2083, found: 571.2079; ^1H NMR (400 MHz, methanol- d_4) δ 0.87–0.96 (m, 3 H), 1.27–1.39 (m, 4 H), 1.50 (d, *J* = 7.1 Hz, 3 H), 1.75–1.87 (m, 1 H), 1.95–2.01 (m, 1 H), 2.05 (dd, *J* = 10.2, 8.9 Hz, 1 H), 2.17 (s, 3 H), 2.13–2.26 (m, 1 H), 2.39 (s, 3 H), 2.98 (dd, *J* = 10.4, 5.0 Hz, 1 H), 3.21 (dd, *J* = 8.5, 6.2 Hz, 1 H), 3.54 (dd, *J* = 10.2, 3.3 Hz, 1 H), 3.80–3.84 (m, 1 H), 4.11 (dd, *J* = 10.2, 5.6 Hz, 1 H), 4.28 (br dd, *J* = 9.2, 0.6 Hz, 1 H), 4.30 (dq, *J* = 7.1, 3.9 Hz, 1 H), 4.65 (dd, *J* = 9.2, 3.9 Hz, 1 H), 5.29 (d, *J* = 5.6 Hz, 1 H), 6.84 (dd, *J* = 8.7, 2.3 Hz, 1 H), 7.08 (dd, *J* = 2.3, 0.4 Hz, 1 H), 7.57 (dd, *J* = 8.7, 0.4 Hz, 1 H). For the qualified analytical purpose, the above colorless solid was further purified by reverse-phase column chromatography (YMC triart C18, 20 × 250 mm, RT, 18.9 ml min⁻¹, 50 mM $\text{AcONH}_4/\text{CH}_3\text{CN}=35/65$) to obtain the highly purified title compound as a colorless solid.

7(S)-7-(2-Aminophenylthio)-7-deoxylincomycin (31). To a solution of compound **18** (39.6 mg, 0.07 mmol) in ethanol (2 ml) were added $\text{SnCl}_2\cdot\text{H}_2\text{O}$ (82.1 mg, 0.36 mmol) and NaBH_4 (13.7 mg, 5.0 mmol) and stirred at RT for 3 h. To the mixture was added aq NaHCO_3 and then extracted with ethyl acetate, washed with water, dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by preparative TLC ($\text{CHCl}_3/\text{CH}_3\text{OH}/28\%$ aq $\text{NH}_4\text{OH}=9/1/0.1$) to obtain the title compound as a colorless solid (10.9 mg, 29%). $[\alpha]_{\text{D}}^{28} +74.5^\circ$ (*c* 0.33, MeOH); ESI-MS (*m/z*) 514 ($\text{M}+\text{H}^+$)⁺ as $\text{C}_{24}\text{H}_{39}\text{N}_3\text{O}_5\text{S}_2$; TOF-ESI-HRMS ($\text{M}+\text{H}^+$)⁺ calcd for $\text{C}_{24}\text{H}_{39}\text{N}_3\text{O}_5\text{S}_2$: 514.2409, found: 514.2412; ^1H NMR (400 MHz, methanol- d_4) δ 0.89–0.97

(m, 3 H), 1.17 (d, *J* = 7.1 Hz, 3 H), 1.28–1.40 (m, 4 H), 1.75–1.85 (m, 1 H), 1.89–1.98 (m, 1 H), 2.04–2.12 (m, 1 H), 2.14–2.23 (m, 1 H), 2.19 (s, 3 H), 2.32 (s, 3 H), 3.01 (dd, *J* = 10.6, 5.0 Hz, 1 H), 3.25 (dd, *J* = 8.3, 5.9 Hz, 1 H), 3.58 (dd, *J* = 10.2, 3.2 Hz, 1 H), 3.67 (dq, *J* = 7.1, 2.8 Hz, 1 H), 3.74 (br dd, *J* = 3.2, 0.5 Hz, 1 H), 4.11 (dd, *J* = 10.2, 5.6 Hz, 1 H), 4.31 (dd, *J* = 9.8, 2.8 Hz, 1 H), 4.38 (br dd, *J* = 9.8, 0.5 Hz, 1 H), 5.27 (d, *J* = 5.6 Hz, 1 H), 6.61–6.67 (m, 1 H), 6.81 (dd, *J* = 8.1, 1.1 Hz, 1 H), 7.13 (ddd, *J* = 8.8, 7.2, 1.5 Hz, 1 H), 7.34 (dd, *J* = 7.7, 1.5 Hz, 1 H).

7(S)-7-(3-Aminophenylthio)-7-deoxylincomycin (32). Compound **19** (238.2 mg, 0.44 mmol), $\text{SnCl}_2\cdot\text{H}_2\text{O}$ (494.4 mg, 2.19 mmol) and NaBH_4 (82.9 mg, 2.19 mmol) were treated according to the similar procedure as described for the preparation of **31** to afford **32** (36.0 mg, 16%) as a colorless solid. $[\alpha]_{\text{D}}^{29} +91.0^\circ$ (*c* 0.60, MeOH); ESI-MS (*m/z*) 514 ($\text{M}+\text{H}^+$)⁺ as $\text{C}_{24}\text{H}_{39}\text{N}_3\text{O}_5\text{S}_2$; TOF-ESI-HRMS ($\text{M}+\text{H}^+$)⁺ calcd for $\text{C}_{24}\text{H}_{39}\text{N}_3\text{O}_5\text{S}_2$: 514.2409, found: 514.2408; ^1H NMR (400 MHz, methanol- d_4) δ 0.89–0.98 (m, 3 H), 1.31 (d, *J* = 7.0 Hz, 3 H), 1.30–1.40 (m, 4 H), 1.79–1.92 (m, 1 H), 1.93–2.10 (m, 2 H), 2.02 (s, 3 H), 2.11–2.23 (m, 1 H), 2.38 (s, 3 H), 2.97 (dd, *J* = 10.7, 4.6 Hz, 1 H), 3.25 (dd, *J* = 8.1, 5.6 Hz, 1 H), 3.59 (dd, *J* = 10.2, 3.3 Hz, 1 H), 3.72 (br d, *J* = 3.3 Hz, 1 H), 3.75–3.83 (m, 1 H), 4.10 (dd, *J* = 10.2, 5.5 Hz, 1 H), 4.33–4.40 (m, 2 H), 5.26 (d, *J* = 5.5 Hz, 1 H), 6.59 (ddd, *J* = 8.0, 2.2, 0.9 Hz, 1 H), 6.71 (ddd, *J* = 7.7, 1.7, 0.9 Hz, 1 H), 6.78–6.82 (m, 1 H), 6.99–7.06 (m, 1 H).

7(S)-7-(4-Aminophenylthio)-7-deoxylincomycin (33). Compound **20** (29.3 mg, 0.054 mmol), $\text{SnCl}_2\cdot\text{H}_2\text{O}$ (60.8 mg, 0.27 mmol) and NaBH_4 (10.0 mg, 0.26 mmol) were treated according to the similar procedure as described for the preparation of **31** to afford **33** (9.7 mg, 35%) as a colorless solid. $[\alpha]_{\text{D}}^{28} +142.0^\circ$ (*c* 0.51, MeOH); ESI-MS (*m/z*) 514 ($\text{M}+\text{H}^+$)⁺ as $\text{C}_{24}\text{H}_{39}\text{N}_3\text{O}_5\text{S}_2$; TOF-ESI-HRMS ($\text{M}+\text{H}^+$)⁺ calcd for $\text{C}_{24}\text{H}_{39}\text{N}_3\text{O}_5\text{S}_2$: 514.2409, found: 514.2411; ^1H NMR (400 MHz, methanol- d_4) δ 0.89–0.98 (m, 3 H), 1.20 (d, *J* = 7.1 Hz, 3 H), 1.30–1.41 (m, 4 H), 1.80–1.90 (m, 1 H), 1.92–2.00 (m, 1 H), 2.04–2.21 (m, 2 H), 2.17 (s, 3 H), 2.34 (s, 3 H), 2.98 (dd, *J* = 10.6, 4.6 Hz, 1 H), 3.24 (dd, *J* = 8.2, 5.6 Hz, 1 H), 3.53 (dq, *J* = 7.1, 2.8 Hz, 1 H), 3.60 (dd, *J* = 10.3, 3.3 Hz, 1 H), 3.68–3.72 (m, 1 H), 4.10 (dd, *J* = 10.3, 5.6 Hz, 1 H), 4.25 (dd, *J* = 9.9, 2.8 Hz, 1 H), 4.38 (br d, *J* = 9.9 Hz, 1 H), 5.27 (d, *J* = 5.6 Hz, 1 H), 6.62–6.68 (m, 2 H), 7.20–7.26 (m, 2 H).

7(S)-7-(4-Methanesulfonamidophenylthio)-7-deoxylincomycin (34). To a solution of compound **33** (83.8 mg, 0.16 mmol) in DMF (1 ml) were added triethylamine (0.027 ml, 0.2 mmol) and methanesulfonyl chloride (0.015 ml, 0.2 mmol) and stirred at RT for 20 min. To the mixture was added saturated aqueous NaHCO_3 (10 ml) and then extracted with ethyl acetate, dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by preparative TLC ($\text{CHCl}_3/\text{CH}_3\text{OH}/28\%$ aq $\text{NH}_4\text{OH}=9/2/0.2$) to obtain the title compound as a colorless solid (28.0 mg, 29%). $[\alpha]_{\text{D}}^{26} +94.2^\circ$ (*c* 1.01, MeOH); ESI-MS (*m/z*) 592 ($\text{M}+\text{H}^+$)⁺ as $\text{C}_{25}\text{H}_{41}\text{N}_5\text{O}_7\text{S}_3$; TOF-ESI-HRMS ($\text{M}+\text{H}^+$)⁺ calcd for $\text{C}_{25}\text{H}_{41}\text{N}_5\text{O}_7\text{S}_3$: 592.2185, found: 592.2180; ^1H NMR (400 MHz, methanol- d_4) δ 0.89–0.98 (m, 3 H), 1.28 (d, *J* = 7.0 Hz, 3 H), 1.31–1.41 (m, 4 H), 1.78–1.92 (m, 1 H), 1.99 (ddd, *J* = 12.9, 7.9, 5.0 Hz, 1 H), 2.04 (s, 3 H), 2.05–2.11 (m, 1 H), 2.11–2.23 (m, 1 H), 2.38 (s, 3 H), 2.94–3.03 (m, 1 H), 2.97 (s, 3 H), 3.25 (dd, *J* = 8.1, 5.6 Hz, 1 H), 3.58 (dd, *J* = 10.2, 3.2 Hz, 1 H), 3.74 (br d, *J* = 3.2 Hz, 1 H), 3.78 (dq, *J* = 7.0, 2.5 Hz, 1 H), 4.10 (dd, *J* = 10.2, 5.6 Hz, 1 H), 4.31–4.36 (m, 1 H), 4.38 (dd, *J* = 9.7, 2.5 Hz, 1 H), 5.26 (d, *J* = 5.6 Hz, 1 H), 7.18–7.25 (m, 2 H), 7.40–7.48 (m, 2 H).

7(S)-7-(5-(2-Aminoacetamido)-1,3,4-thiadiazol-2-ylthio)-7-deoxylincomycin (36). To a solution of compound **1** (240 mg, 0.39 mmol) in THF (5 ml) at 0 °C were added triphenylphosphine (150 mg, 0.57 mmol), diethylazodicarboxylate (0.1 ml, 0.55 mmol) and *t*-butyl 2-((5-mercapto-1,3,4-thiadiazol-2-yl)amino)-2-oxoethyl)carbamate (150 mg, 0.52 mmol) and stirred RT for 17 h. The mixture was concentrated under reduced pressure, and the resulting residue (compound **35**) in 90% aqueous trifluoroacetic acid (5 ml) was stirred at RT for 30 min. The mixture was concentrated under reduced pressure. The resulting residue was purified by preparative reverse-phase column chromatography (YMC triart C18, 20 × 250 mm², RT, 18.9 ml min⁻¹, 0.1% aq TFA/ $\text{CH}_3\text{CN}=85/15$). This trifluoroacetate was desalted by preparative reverse-phase column chromatography (YMC triart C18, 20 × 250 mm², RT, 18.9 ml min⁻¹, $\text{H}_2\text{O}/\text{MeOH}=100/0$ (15 min) → 0/100 (15–40 min)) to obtain the

highly purified title compound as a colorless solid (54.0 mg, 24%). $[\alpha]_{\text{D}}^{25} +100.4^{\circ}$ (*c* 0.29, MeOH); ESI-MS (*m/z*) 578 (M+H)⁺ as C₂₂H₃₈N₆O₆S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₂H₃₈N₆O₆S₃: 579.2093, found: 579.2095; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.86–0.97 (m, 3 H), 1.29–1.39 (m, 4 H), 1.42 (d, *J* = 7.0 Hz, 3 H), 1.77–1.87 (m, 1 H), 1.93–2.09 (m, 2 H), 2.06 (s, 3 H), 2.13–2.25 (m, 1 H), 2.32 (s, 3 H), 2.98 (dd, *J* = 10.5, 5.0 Hz, 1 H), 3.26 (dd, *J* = 8.5, 6.2 Hz, 1 H), 3.57 (dd, *J* = 10.2, 3.2 Hz, 1 H), 3.54–3.62 (m, 2 H), 3.78 (br dd, *J* = 3.2, 0.5 Hz, 1 H), 4.08 (dq, *J* = 7.0, 2.8 Hz, 1 H), 4.10 (dd, *J* = 10.2, 5.6 Hz, 1 H), 4.40 (br dd, *J* = 9.8, 0.5 Hz, 1 H), 4.47 (dd, *J* = 9.8, 2.8 Hz, 1 H), 5.26 (d, *J* = 5.6 Hz, 1 H).

7(S)-7-(5-Amino-1,3,4-thiadiazol-2-ylthio)-7-deoxy-2,3,4-tris-O-(trimethylsilyl)lincomycin (37). To a solution of compound **21** (403 mg, 0.77 mmol) in pyridine (8 ml) were added trimethylchlorosilane (0.61 ml, 4.8 mmol) and hexamethyldisilazane (1.05 ml, 5.0 mmol) and stirred at RT for 2.5 h, and then it was concentrated under reduced pressure. The residue was diluted with saturated aqueous NH₄Cl and extracted with ethyl acetate, washed with brine, dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 1/1) to obtain the title compound as a colorless solid (467 mg, 82%). ESI-MS (*m/z*) 738 (M+H)⁺ as C₂₉H₅₉N₅O₅S₃Si₃; ¹H NMR (400 MHz, chloroform-*d*) δ 0.13 (s, 18H), 0.17 (s, 9 H), 0.79–0.97 (m, 3 H), 1.15–1.37 (m, 4 H), 1.42 (d, *J* = 6.9 Hz, 3 H), 1.71–1.90 (m, 2 H), 1.91–2.06 (m, 2 H), 2.08 (s, 3 H), 2.40 (s, 3 H), 2.99 (dd, *J* = 10.7, 3.6 Hz, 1 H), 3.12–3.23 (m, 1 H), 3.61 (dd, *J* = 9.6, 2.5 Hz, 1 H), 3.74 (br d, *J* = 2.2 Hz, 1 H), 4.04–4.24 (m, 3 H), 4.60–4.70 (m, 1 H), 5.23 (d, *J* = 5.5 Hz, 1 H), 5.50 (s, 2 H), 7.59 (d, *J* = 10.7 Hz, 1 H).

7(S)-7-Deoxy-7-(5-(2-methoxyacetamido)-1,3,4-thiadiazol-2-ylthio)lincomycin (38). To a solution of compound **37** (104 mg, 0.14 mmol) in THF (3 ml) were added triethylamine (0.06 ml, 0.43 mmol) and 2-methoxyacetyl chloride (0.02 ml, 0.22 mmol) and stirred at 0 °C for 2 h. The mixture was diluted with saturated aqueous NH₄Cl and extracted with ethyl acetate, washed with brine, dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by preparative TLC (CHCl₃/CH₃OH = 19/1) to obtain 7(S)-7-deoxy-7-(5-(2-methoxyacetamido)-1,3,4-thiadiazol-2-ylthio)-2,3,4-tris-O-(trimethylsilyl)lincomycin as a colorless solid (109.8 mg, 96%). **7(S)-7-Deoxy-7-(5-(2-methoxyacetamido)-1,3,4-thiadiazol-2-ylthio)-2,3,4-tris-O-(trimethylsilyl)lincomycin** (109.8 mg, 0.14 mmol) in 1 N HCl (1 ml)–MeOH (1 ml) was stirred at RT for 2 h. The mixture was concentrated under reduced pressure. The resulting residue was dissolved by water and washed with diethyl ether. The mixture was added NaHCO₃ (150 mg) and then extracted with ethyl acetate, dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by preparative TLC (CHCl₃/CH₃OH/28% aq NH₄OH = 5/1/0.1) to obtain the title compound as a colorless solid (37.8 mg, 47%). $[\alpha]_{\text{D}}^{26} +98.3^{\circ}$ (*c* 0.65, MeOH); ESI-MS (*m/z*) 594 (M+H)⁺ as C₂₃H₃₉N₅O₇S₃; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₃H₃₉N₅O₇S₃: 594.2090, found: 594.2095; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.88–0.97 (m, 3 H), 1.30–1.41 (m, 4 H), 1.46 (d, *J* = 7.0 Hz, 3 H), 1.82–1.92 (m, 1 H), 1.97–2.08 (m, 1 H), 2.04 (s, 3 H), 2.08–2.15 (m, 1 H), 2.15–2.27 (m, 1 H), 2.39 (s, 3 H), 3.08 (dd, *J* = 10.4, 5.0 Hz, 1 H), 3.25–3.34 (m, 1 H), 3.48 (s, 3 H), 3.57 (dd, *J* = 10.3, 3.2 Hz, 1 H), 3.77–3.81 (m, 1 H), 4.10 (dd, *J* = 10.3, 5.6 Hz, 1 H), 4.16–4.24 (m, 1 H), 4.21 (s, 2 H), 4.39 (br dd, *J* = 9.8, 0.6 Hz, 1 H), 4.53 (dd, *J* = 9.8, 3.1 Hz, 1 H), 5.26 (d, *J* = 5.6 Hz, 1 H).

7-O-Methanesulfonyl-2,3,4-tris-O-(trimethylsilyl)lincomycin (39). To a solution of compound **1** (5.0 g, 8.0 mmol) in chloroform (25 ml) were added triethylamine (2.79 ml, 20.1 mmol) and methanesulfonyl chloride (1.24 ml, 16.1 mmol) and stirred at RT for 3 h. The mixture was dissolved by chloroform (60 ml), washed with saturated aqueous NaHCO₃ (50 ml), dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 3/1) to obtain the title compound as a colorless solid (5.50 g, 98%). ESI-MS (*m/z*) 701 (M+H)⁺ as C₂₈H₆₀N₂O₈S₂Si₃; ¹H NMR (400 MHz, chloroform-*d*) δ 0.13 (s, 9 H), 0.14 (s, 9 H), 0.17 (s, 9 H), 0.89 (br t, *J* = 6.9 Hz, 3 H), 1.21–1.36 (m, 4 H), 1.40 (d, *J* = 6.6 Hz, 3 H), 1.79–1.89 (m, 1 H), 1.92–2.09 (m, 3 H), 2.11 (s, 3 H), 2.40 (s, 3 H), 2.99 (dd, *J* = 10.7, 3.7 Hz, 1 H), 3.09 (s, 3 H), 3.14–3.21 (m, 1 H), 3.52

(dd, *J* = 9.5, 2.4 Hz, 1 H), 3.75 (br d, *J* = 2.4 Hz, 1 H), 3.90 (d, *J* = 9.7 Hz, 1 H), 4.15 (dd, *J* = 9.5, 5.6 Hz, 1 H), 4.70–4.78 (m, 1 H), 5.09–5.15 (m, 1 H), 5.16 (d, *J* = 5.6 Hz, 1 H), 7.61 (d, *J* = 10.7 Hz, 1 H).

7(S)-7-Deoxy-7-(4-methoxycarbonylphenylthio)lincomycin (40). To a solution of compound **39** (100 mg, 0.14 mmol) in DMF (1.0 ml) were added K₂CO₃ (59.8 mg, 0.43 mmol) and methyl 4-mercaptobenzoate (48.9 mg, 0.29 mmol) and stirred at 80 °C for 3 h. The mixture was cooled down to RT, diluted with 1 N HCl (2 ml)–MeOH (1 ml), stirred at RT for 2 h and concentrated under reduced pressure. The resulting residue was dissolved by water, washed with diethyl ether. To the mixture was added NaHCO₃ (150 mg) and then extracted with ethyl acetate, washed with water, dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by preparative TLC (CHCl₃/CH₃OH/28% aq NH₄OH = 5/1/0.1) to obtain the title compound as a colorless solid (58.0 mg, 73%). $[\alpha]_{\text{D}}^{24} +84.6^{\circ}$ (*c* 0.97, MeOH); ESI-MS (*m/z*) 557 (M+H)⁺ as C₂₆H₄₀N₂O₇S₂; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₆H₄₀N₂O₇S₂: 557.2355, found: 557.2359; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.88–0.96 (m, 3 H), 1.29–1.37 (m, 4 H), 1.40 (d, *J* = 6.8 Hz, 3 H), 1.79–1.91 (m, 1 H), 1.84 (s, 3 H), 1.96–2.05 (m, 1 H), 2.05–2.12 (m, 1 H), 2.12–2.25 (m, 1 H), 2.40 (s, 3 H), 3.00 (dd, *J* = 10.6, 4.8 Hz, 1 H), 3.24 (dd, *J* = 8.2, 5.7 Hz, 1 H), 3.58 (dd, *J* = 10.2, 3.2 Hz, 1 H), 3.78 (br dd, *J* = 3.2, 0.7 Hz, 1 H), 3.89 (s, 3 H), 4.03 (dq, *J* = 6.8, 2.8 Hz, 1 H), 4.10 (dd, *J* = 10.2, 5.6 Hz, 1 H), 4.37 (br dd, *J* = 9.7, 0.7 Hz, 1 H), 4.52 (dd, *J* = 9.7, 2.8 Hz, 1 H), 5.24 (d, *J* = 5.6 Hz, 1 H), 7.42–7.49 (m, 2 H), 7.90–7.97 (m, 2 H).

7(S)-7-Deoxy-7-(3-methoxycarbonylphenylthio)lincomycin (41). Compound **39** (200 mg, 0.29 mmol) and methyl 3-mercaptobenzoate (95.9 mg, 0.57 mmol) were treated according to the similar procedure as described for the preparation of **40** to afford **41** (101.6 mg, 64%) as a colorless solid. $[\alpha]_{\text{D}}^{29} +93.6^{\circ}$ (*c* 2.37, MeOH); ESI-MS (*m/z*) 557 (M+H)⁺ as C₂₅H₃₇N₃O₆S₂; TOF-ESI-HRMS (M+H)⁺ calcd for C₂₅H₃₇N₃O₆S₂: 557.2355, found: 557.2355; ¹H NMR (400 MHz, methanol-*d*₄) δ 0.88–0.97 (m, 3 H), 1.29–1.40 (m, 4 H), 1.33 (d, *J* = 7.0 Hz, 3 H), 1.78–1.91 (m, 1 H), 1.95–2.04 (m, 1 H), 1.97 (s, 3 H), 2.04–2.11 (m, 1 H), 2.11–2.23 (m, 1 H), 2.39 (s, 3 H), 2.99 (dd, *J* = 10.7, 4.7 Hz, 1 H), 3.23 (dd, *J* = 8.2, 5.7 Hz, 1 H), 3.58 (dd, *J* = 10.2, 3.3 Hz, 1 H), 3.74–3.79 (m, 1 H), 3.87–3.96 (m, 1 H), 3.91 (s, 3 H), 4.11 (dd, *J* = 10.2, 5.6 Hz, 1 H), 4.36 (br dd, *J* = 9.7, 0.6 Hz, 1 H), 4.48 (dd, *J* = 9.7, 2.8 Hz, 1 H), 5.27 (d, *J* = 5.6 Hz, 1 H), 7.42–7.49 (m, 1 H), 7.67 (ddd, *J* = 7.8, 1.8, 1.1 Hz, 1 H), 7.87–7.92 (m, 1 H), 8.02–8.06 (m, 1 H).

In vitro antibacterial activity

MIC, $\mu\text{g ml}^{-1}$ was determined by the agar dilution method, which was described in Clinical and Laboratory Standards Institute (M7-A5 in 2000). Test strains of *S. pneumoniae* and *S. pyogenes* were subjected to seed culture using brain heart infusion agar (Becton Dickinson and Company, Tokyo, Japan) and 5% defibrinated horse blood. Test strains of *H. influenzae* were subjected to seed culture using sensitivity disk agar-N 'Nissui' (SDA; Nissui, Tokyo, Japan), 5% defibrinated horse blood, 5 $\mu\text{g ml}^{-1}$ Hemin and 15 $\mu\text{g ml}^{-1}$ NAD. A 5- μl portion of cell suspension of the test strains having about 10⁶ CFU per ml was inoculated into SDA supplemented with 5% defibrinated horse blood, 5 $\mu\text{g ml}^{-1}$ Hemin and 15 $\mu\text{g ml}^{-1}$ NAD and incubated at 37 °C for 18–22 h. Then MIC was measured.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank Mr A Tamura, Dr E Shitara, Dr T Okutomi, Dr T Yoshida, and Mr M Yamamoto for encouragement and valuable discussion. We are grateful to Professor Emeritus Dr M Konno for supervision through our in-house drug discovery program in lincomycin field. We also thank Ms M Ishii for direction in intellectual properties; Mr T Watanabe for contribution toward computational chemistry; Ms S Miki, Ms T Miyara and Ms K Kaneda for analytical and synthetic chemistry; Mr Y Takayama, Ms Y Inoue, Mr T Matsuhira and Ms K Yamada for biological studies; and Ms M Takagi for English manuscript.

- 1 Reinert, R. R., van der Linden, M. & Al-Lahham, A. Molecular characterization of the first telithromycin-resistant *Streptococcus pneumoniae* isolate in Germany. *Antimicrob. Agents Chemother.* **49**, 3520–3522 (2005).
- 2 Kim, S. H. *et al.* Changing trends in antimicrobial resistance and serotypes of *Streptococcus pneumoniae* isolates in Asian countries: an Asian Network for Surveillance of Resistant Pathogens (ANSORP) study. *Antimicrob. Agents Chemother.* **56**, 1418–1426 (2012).
- 3 Ajito, K., Miura, T., Furuuchi, T. & Tamura, A. Sixteen-membered macrolides: chemical modifications and future applications. *Heterocycles* **89**, 281–352 (2014).
- 4 Schlünzen, F. *et al.* Structural basis for the interaction of antibiotics with the peptidyl transferase centre in eubacteria. *Nature* **413**, 814–821 (2001).
- 5 Poehlsaard, J. & Douthwaite, S. The macrolide binding site on the bacterial ribosome. *Curr. Drug Targets Infect. Disord.* **2**, 67–78 (2002).
- 6 Tu, D., Blaha, G., Moore, P. B. & Steitz, T. A. Structures of MLS_GK antibiotics bound to mutated large ribosomal subunits provide a structural explanation for resistance. *Cell* **121**, 257–270 (2005).
- 7 Welsblum, B. Erythromycin resistance by ribosome modification. *Antimicrob. Agents Chemother.* **39**, 577–585 (1995).
- 8 Vester, B. & Douthwaite, S. Macrolide resistance conferred by base substitutions in 23 S rRNA. *Antimicrob. Agents Chemother.* **45**, 1–12 (2001).
- 9 O'hara, K. Up-to-date report on the resistance of macrolide antibiotics and future macrolides. *Jpn. J. Chemother.* **48**, 169–190 (2000).
- 10 Syrogiannopoulos, G. A. *et al.* Antimicrobial susceptibility and macrolide resistance inducibility of *Streptococcus pneumoniae* carrying *erm* (A), *erm* (B), or *mef* (A). *Antimicrob. Agents Chemother.* **47**, 2699–2702 (2003).
- 11 Morimoto, S., Takahashi, Y., Watanabe, Y. & Omura, S. Chemical modification of erythromycins. I. Synthesis and antibacterial activity of 6-O-methylerythromycins A. *J. Antibiot.* **37**, 187–189 (1984).
- 12 Slobodan, D. *et al.* Erythromycin series. Part 13. Synthesis and structure elucidation of 10-dihydro-10-deoxy-11-methyl-11-azaerythromycin A. *J. Chem. Res. Synop* **1988**, 152–153 (1988).
- 13 Denis, A. *et al.* Synthesis and antibacterial activity of HMR 3647 a new ketolide highly potent against erythromycin-resistant and susceptible pathogens. *Bioorg. Med. Chem. Lett.* **9**, 3075–3080 (1999).
- 14 Berisio, R. *et al.* Structural insight into the antibiotic action of telithromycin against resistant mutants. *J. Bacteriol.* **185**, 4276–4279 (2003).
- 15 Douthwaite, S., Hansen, L. H. & Mauvais, P. Macrolide-ketolide inhibition of MLS-resistant ribosomes is improved by alternative drug interaction with domain II of 23 S rRNA. *Mol. Microbiol.* **36**, 183–193 (2000).
- 16 Hansen, L. H., Mauvais, P. & Douthwaite, S. The macrolide-ketolide antibiotic binding site is formed by structures in domains II and V of 23 S ribosomal RNA. *Mol. Microbiol.* **31**, 623–631 (1999).
- 17 Xiong, L., Shah, S., Mauvais, P. & Mankin, A. S. A ketolide resistance mutation in domain II of 23 S rRNA reveals the proximity of hairpin 35 to the peptidyl transferase centre. *Mol. Microbiol.* **31**, 633–639 (1999).
- 18 Clay, K. D. *et al.* Severe hepatotoxicity of telithromycin: three case reports and literature review. *Ann. Intern. Med.* **144**, 415–420 (2006).
- 19 Ross, D. B. The FDA and the case of ketek. *N. Engl. J. Med.* **356**, 1601–1604 (2007).
- 20 Gleason, P. P., Walters, C., Heaton, A. H. & Schafer, J. A. Telithromycin: the perils of hasty adoption and persistence of off-label prescribing. *J. Manag. Care Pharm.* **13**, 20–25 (2007).
- 21 Department of Health and Human Services. Telithromycin (marketed as ketek) information. Available at <http://www.fda.gov/drugs/drugsafety/postmarketdrugsafetyinformationforpatientsandproviders/ucm107824.htm>.
- 22 Mason, D. J., Dietz, A. & Deboer, C. Lincomycin, a new antibiotic I. Discovery and biological properties. *Antimicrob. Agents Chemother.* 554–559 (1962).
- 23 Magerlein, B. J. & Lincomycin, X. The chemical synthesis of lincomycin. *Tetrahedron Lett.* **1**, 33–36 (1970).
- 24 Howarth, G. B., Szarek, W. A. & Jones, J. K. N. The synthesis of lincomycin. *J. Chem. Soc. (c)* **16**, 2218–2224 (1970).
- 25 Perlman, D. *Structure-Activity Relationships Among the Semisynthetic Antibiotics*, (Academic Press, New York, San Francisco, London, A Subsidiary of Harcourt Brace Jovanovich, Publishers, 600–651, 1977).
- 26 Birkenmeyer, R. D. & Kagan, F. Lincomycin. XI. Synthesis and structure of clindamycin. A potent antibacterial agent. *J. Med. Chem.* **13**, 616–619 (1970).
- 27 Hoeksema, H. Octoses from antibiotics. The Upjohn Company, Kalamazoo, Mich., *Abstr. Pap. Division of Carbohydrate Chemistry, 149th Meet. Am. Chem. Soc. Detroit, Mich.*, p. 9C (1965).
- 28 Magerlein, B. J., Birkenmeyer, R. D. & Kagan, F. Chemical modification of lincomycin. *Antimicrob. Agents Chemother.* 727–736 (1966).
- 29 Sinkula, A. A., Morozowich, W., Lewis, C. & Mackellar, F. A. Synthesis and bioactivity of lincomycin-7-monoesters. *J. Pharm. Sci.* **58**, 1389–1392 (1969).
- 30 Magerlein, B. J. & Kagan, F. Lincomycin. IX. 7-Thiol and thioamido analogs of lincomycin. *J. Med. Chem.* **12**, 974–977 (1969).
- 31 Lewis, J. G. *et al.* Novel Antimicrobial 7-methyl Lincosamides: Prolamide Analogs. 43rd Interscience Conference on Antimicrobial Agents and Chemotherapy. Poster F-1388. (Washington, DC, USA, 2004).
- 32 Bannister, B. Modifications of lincomycin involving the carbohydrate portion. Part III. The 7-O-methyl and 6-de-(1-hydroxyethyl) analogues. *J. Chem. Soc. Perkin Trans. I* **16**, 1676–1682 (1973).
- 33 Bannister, B. Modifications of lincomycin involving the carbohydrate portion. Part IV. (7S)-7-alkoxy-7-deoxy-analogues. *J. Chem. Soc. Perkin Trans. I* 360–369 (1974).
- 34 Bannister, B. & Mydlow, P. K. The S-alkylation of sulphides by an activated carbohydrate epimine under acidic catalysis: The formation of α -acetamido-sulphides. Part 5. The introduction of functionality into the sulphide substituent. *J. Chem. Res. (S)* **1989**, 90–91 (1989).
- 35 Bannister, B. The S-alkylation of sulphides by an activated carbohydrate epimine under acidic catalysis: The formation of α -acetamido-sulphides. Part 4. Reaction with dithioacetals and monothioacetals. *J. Chem. Soc., Perkin. Trans. I* **1980**, 540–552 (1980).
- 36 Bannister, B. (7S)-7-deoxy-7-substituted-alkylthio-lincomycin. S-Alkylation of sulphides by an activated epimine under acidic catalysis: formation of α -acetamido-sulphides. *Tetrahedron* **40**, 1633–1660 (1984).
- 37 Bannister, B. (The Upjohn company). Derivatives of lincomycin and its analogs and process. US Patent US3915954 A, (1973).
- 38 Bannister, B. (The Upjohn company). Derivatives of lincomycin and its analogs and process. Canadian Patent CA-971956 A1 (1972).
- 39 Sztaricskai, F. *et al.* Semisynthetic modification of antibiotic lincomycin. *J. Antibiot.* **49**, 941–943 (1996).
- 40 Bannister, B. Modifications of lincomycin involving the carbohydrate portion. Part I. The 2-O-methyl and 2-deoxy-analogues. *J. Chem. Soc., Perkin Trans. I* **1972**, 3025–3030 (1972).
- 41 Houtman, R. L. & Mich, P. (The Upjohn company). Trimethylsilyl ethers of lincomycin and its compounds. US Patent US3418414 (1966).
- 42 Umemura, E. *et al.* Synthesis of novel lincomycin derivatives and their in vitro antibacterial activities. *J. Antibiot.* **66**, 195–198 (2013).
- 43 Umemura, E. *et al.* Lincomycin derivative and antimicrobial agent containing the same as active ingredient. WO 2007/066805 A1 (2007).