

## ORIGINAL ARTICLE

# Novel autophagy inducers lentztrehaloses A, B and C

Shun-ichi Wada, Yumiko Kubota, Ryuichi Sawa, Maya Umekita, Masaki Hatano, Shun-ichi Ohba, Chigusa Hayashi, Masayuki Igarashi and Akio Nomoto<sup>\*‡</sup>

Trehalose has widespread use as a sweetener, humectant and stabilizer, and is now attracting attention as a promising candidate for the treatment of neurodegenerative diseases as it is an autophagy inducer and chemical chaperone. However, the bioavailability of trehalose is low because it is digested by the hydrolyzing enzyme trehalase, expressed in the intestine and kidney. Enzyme-stable analogs of trehalose would potentially solve this problem. We have previously reported an enzyme-stable analog of trehalose, lentztrehalose, and herein report two new analogs. The original lentztrehalose has been renamed lentztrehalose A and the analogs named lentztrehaloses B and C. Lentztrehalose B is a di-dehydroxylated analog and lentztrehalose C is a cyclized analog of lentztrehalose A. All the lentztrehaloses are only minimally hydrolyzed by mammalian trehalase. The production of the lentztrehaloses is high in rather dry conditions and low in wet conditions. Lentztrehalose B shows a moderate antioxidative activity. These facts suggest that the lentztrehaloses are produced as humectants or protectants for the producer microorganism under severe environmental conditions. All the lentztrehaloses induce autophagy in human cancer cells at a comparable level to trehalose. Considering the enzyme-stability, these lentztrehaloses can be regarded as promising new drug candidates for the treatment of neurodegenerative diseases and other autophagy-related diseases, such as diabetes, arteriosclerosis, cancer and heart disease.

*The Journal of Antibiotics* (2015) 68, 521–529; doi:10.1038/ja.2015.23; published online 11 March 2015

## INTRODUCTION

Trehalose is a unique disaccharide formed from two molecules of glucose linked by an  $\alpha,\alpha$ -1,1-glucoside bond. It is prevalently observed in plants, microorganisms and animals.<sup>1</sup> Industrially, trehalose is widely used in foods, drugs and cosmetics as a sweetener, humectant or stabilizer.<sup>2</sup> Trehalose is known to have several biological activities, including bone reinforcement,<sup>3</sup> anti-tumor<sup>4</sup> and anti-obesity<sup>5</sup> activities. As it is a chemical chaperone, a neuroprotective effect is also expected and the application of trehalose in the treatment of neurodegenerative diseases has also been investigated. Decrease of aggregates, improved motor dysfunction and extended life span were reported in mice administered trehalose in a model of Huntington's disease.<sup>6</sup> In addition, the recent discovery of the autophagy-inducing activity of trehalose in cultured cells is of considerable interest.<sup>7</sup>

Autophagy is a catabolic process that involves degradation of unnecessary or dysfunctional cellular components through the action of lysosomes.<sup>8</sup> The accumulation of abnormal proteins in neural cells is a common feature in neurodegenerative diseases, such as Alzheimer's disease, amyotrophic lateral sclerosis, Parkinson's disease, Huntington's disease and prion diseases, and which causes the progression of such diseases.<sup>9</sup> The accumulated proteins can be decreased by autophagy, which leads to an increased survival of neural cells and improvement in the disease condition. Thus, autophagy-inducing agents are attracting attention as new drug candidates for such diseases. In the past few years, several groups

have reported alleviation of disease progression after trehalose administration in mouse neurodegenerative disease models of prion, Alzheimer's, amyotrophic lateral sclerosis and Parkinson's diseases.<sup>10–13</sup> Unlike other autophagy inducers, such as rapamycin and verapamil, trehalose is a food constituent and thus attracts special attention as a safe candidate for the treatment of neurodegenerative diseases. There are, however, several problems with the therapeutic use of trehalose. Orally taken trehalose is digested by the hydrolyzing enzyme trehalase expressed in the intestine and kidney,<sup>14</sup> making its bioavailability low. Hence, a large amount of trehalose would be needed for the treatment of disease. Applying the administration protocol used in mouse experiments to human use, more than 50 g of trehalose would be needed to be taken daily. The burst release of the resultant product, glucose, is also problematic and may cause, or exacerbate, diabetes, obesity and vascular disorders. The development of enzyme-stable analogs of trehalose would be helpful in solving this problem.

We previously reported such an enzyme-stable analog, lentztrehalose, isolated from a rare actinomycete *Lentzea* sp. ML457-mF8.<sup>15</sup> Lentztrehalose is only minimally hydrolyzed by a porcine kidney trehalase and thought to be stable in the mammalian body. This stability is likely the cause of the better or comparable levels of antitumor, bone reinforcement and suppression of obesity activities in mice administered lentztrehalose compared with trehalose, when lentztrehalose was administered at 1/4–1/2 the amount of trehalose.


Institute of Microbial Chemistry (BIKAKEN), Tokyo, Japan

Correspondence: Dr S Wada, Institute of Microbial Chemistry (BIKAKEN), 3-14-23, Kamiosaki, Shinagawa-ku, Tokyo 141-0021, Japan.

E-mail: wadas@bikaken.or.jp

<sup>\*</sup>Deceased on 13th November 2014.

Received 10 December 2014; revised 3 February 2015; accepted 6 February 2015; published online 11 March 2015



**Figure 1** Structure of lentsztrehaloses. (a) Structure of lentsztrehaloses. TRH: trehalose, LTA, B, C: lentsztrehaloses A, B, C. (b) <sup>1</sup>H-<sup>1</sup>H COSY and HMBC correlations of lentsztrehaloses B and C.

In this study, we describe two new lentsztrehalose analogs from the same actinomycete strain. Their structures, physicochemical properties and biological activities, including autophagy-inducing activities are reported herein.

## RESULTS AND DISCUSSION

### Isolation and properties of lentsztrehaloses B and C

In this study, we describe two new analogs of lentsztrehalose. Since the new analogs were isolated from the same actinomycete strain *Lentzea* sp. ML457-mF8 and highly relevant to lentsztrehalose, we renamed lentsztrehalose as lentsztrehalose A and named the new analogs lentsztrehaloses B and C (Figure 1a). The producing strain was previously cultured on steamed flattened barley and impurities hid the existence of lentsztrehaloses B and C in the isolation process. The media was changed to steamed germinated brown rice, which has less impurities, and the new analogs were found in the more hydrophobic fractions during the isolation of lentsztrehalose A. To obtain pure (>95%) lentsztrehaloses, the culture extract was sequentially subjected to chromatography with DIAION HP20, octadecyl silica (HPLC), polyamine (HPLC), silica gel, activated carbon and Sephadex LH-20 columns. Approximately 1.0, 1.0 and 0.6 g of lentsztrehaloses A, B and C, respectively were purified from media made from 1.0 kg of germinated brown rice grain.

The physicochemical properties of lentsztrehaloses B and C are shown in Table 1. The optical rotations of lentsztrehaloses B and C were +177° and +160°, respectively. The molecular formulas of lentsztrehaloses B and C were determined to be C<sub>17</sub>H<sub>30</sub>O<sub>11</sub> and C<sub>17</sub>H<sub>30</sub>O<sub>12</sub> by high resolution electrospray ionization (HRESI)-MS, elemental analysis and NMR data (Table 1 and Table 2, and Supplementary Figures 1–19). The sweetness of the lentsztrehaloses was comparable to that of trehalose (Table 1) with lentsztrehalose C the

sweetest. Lentsztrehalose B had some sharpness of taste, which made it less sweet. None of the lentsztrehaloses showed toxicity against 36 strains of microbes from 16 species up to 128 μg ml<sup>-1</sup> and 53 strains of human and mouse cancer cell lines and three mouse splanchnic primary culture cells up to 200 μg ml<sup>-1</sup>. Acute toxicity was not observed in the ICR mouse up to 500 mg kg<sup>-1</sup> po administration and i.v. injection (data not shown).

### Structure elucidation of lentsztrehalose B

The structure determination was mainly carried out by comparison with the published data for lentsztrehalose A.<sup>15</sup> The <sup>1</sup>H-<sup>1</sup>H COSY and TOCSY spectra indicated a two spin-spin coupling system resulting from two hexoses between H-1' (δ<sub>H</sub> 5.11) and H-6' (δ<sub>H</sub> 3.65, 3.77), and H-1'' (δ<sub>H</sub> 5.09) and H-6'' (δ<sub>H</sub> 3.67, 3.79) as shown in Figure 1 and Supplementary Figures 4 and 7. The hexoses are connected with a 1,1-glycosidic linkage as determined by HMBC correlations, from H-1' to C-1'' (δ<sub>C</sub> 95.1) and H-1'' to C-1' (δ<sub>C</sub> 95.0). These chemical shifts (Table 2) and the characteristic <sup>1</sup>H coupling constants (all large except for <sup>3</sup>J<sub>H1'-H2</sub> = 3.7 Hz and <sup>3</sup>J<sub>H1''-H2'</sub> = 3.7 Hz), NOEs (Supplementary Figures 8 and 9) and optical rotation listed above revealed a trehalose moiety. In lentsztrehalose B, another <sup>1</sup>H-<sup>1</sup>H COSY correlation, from the oxymethylene at δ<sub>H</sub> 4.17 and 4.37 (H-1) to the olefinic proton at δ<sub>H</sub> 5.37 (H-2) was observed. Further connectivity was established by HMBC (Supplementary Figure 6). Long-range couplings were observed from H-5 (δ<sub>H</sub> 1.74) and H-4 (δ<sub>H</sub> 1.69) to C-3 (δ<sub>C</sub> 137.6) and C-2 (δ<sub>C</sub> 122.7); from H-5 to C-4 (δ<sub>C</sub> 18.2); from H-4 to C-5 (δ<sub>C</sub> 25.9); from H-1 to C-4' (δ<sub>C</sub> 79.0); and from H-4' (δ<sub>H</sub> 3.25) to C-1 (δ<sub>C</sub> 70.1) established that lentsztrehalose B possesses a 3-methyl-2-butenyl side chain attached to the C-4' of trehalose. The structure of lentsztrehalose B was elucidated to be 4-O-[(3-methyl-2-butenyl)] trehalose as shown in Figure 1.

**Table 1** Physicochemical properties of lentztrehaloses B and C

	Lentztrehalose B	Lentztrehalose C
Appearance	White solid	White solid
Optical rotation	$[\alpha]_D^{25} +177$ (c 1.06, MeOH)	$[\alpha]_D^{25} +160$ (c 1.03, MeOH)
UV $\lambda_{max}$ (MeOH)	End absorption	End absorption
IR (KBr) $\nu_{max}$ (cm <sup>-1</sup> )	3410, 2931, 1674, 1449, 1378, 1203, 1146, 1105, 1053, 995, 839, 801, 613	3422, 2935, 1635, 1457, 1191, 1153, 1054, 996, 941, 845, 803, 637
Molecular formula	C <sub>17</sub> H <sub>30</sub> O <sub>11</sub>	C <sub>17</sub> H <sub>30</sub> O <sub>12</sub>
<i>HRESI-MS</i> ( <i>m/z</i> )		
Calculated	433.1680 (C <sub>17</sub> H <sub>30</sub> O <sub>11</sub> +Na) <sup>+</sup>	449.1629 (C <sub>17</sub> H <sub>30</sub> O <sub>12</sub> +Na) <sup>+</sup>
Found	433.1683 (M+Na) <sup>+</sup>	449.1629 (M+Na) <sup>+</sup>
<i>Elemental analysis</i>		
Calculated	C 47.7% H 7.5% (C <sub>17</sub> H <sub>30</sub> O <sub>11</sub> ·H <sub>2</sub> O)	C 45.9% H 7.3% (C <sub>17</sub> H <sub>30</sub> O <sub>12</sub> ·H <sub>2</sub> O)
Found	C 47.7% H 7.8%	C 45.7% H 7.3%
ORAC value (μmol TE g <sup>-1</sup> )	207.3	8.1
Sweetness score <sup>a</sup>	5.5 ± 4.3	8.2 ± 3.9

Abbreviation: ORAC, oxygen radical absorbance capacity.

<sup>a</sup>Arbitrary score measured by a sensory test.Ten panelists tasted 10 μl of 500 mg ml<sup>-1</sup> blinded solutions of the lentztrehaloses, sucrose, trehalose and water. The sweetness of the lentztrehaloses and trehalose was scored on a scale where water and sucrose were scored as 0 and 10, respectively. The scores of lentztrehalose A and trehalose measured at the same time were 6.2 ± 2.5 and 6.8 ± 2.2, respectively. Mean ± s.d.

### Structure elucidation of lentztrehalose C

The <sup>1</sup>H–<sup>1</sup>H COSY and TOCSY spectra also indicated a two spin–spin coupling system from H-1' (δ<sub>H</sub> 5.14) to H-6' (δ<sub>H</sub> 3.64, 3.73) and from H-1'' (δ<sub>H</sub> 5.10) to H-6'' (δ<sub>H</sub> 3.67, 3.79) derived from two hexoses coupled via a 1,1-glycosidic linkage as determined by HMBC in the same manner as for lentztrehalose B (Supplementary Figure 13 and 15–17). The chemical shifts of lentztrehalose C were slightly different from those of lentztrehalose A in C-2' to C-4', but the <sup>1</sup>H coupling constants (Table 2), which were closely related to those of A, and the optical rotation indicated that lentztrehalose C also possesses a trehalose moiety. In the HMBC spectrum, cross-peaks were observed from H-5 (δ<sub>H</sub> 1.37) and H-4 (δ<sub>H</sub> 1.16) to C-3 (δ<sub>C</sub> 78.7) and C-2 (δ<sub>C</sub> 79.1); from H-5 to C-4 (δ<sub>C</sub> 23.4); from H-4 to C-5 (δ<sub>C</sub> 25.0); from H-1 (δ<sub>H</sub> 3.63, 3.86) to C-4' (δ<sub>C</sub> 79.0); and from H-4' (δ<sub>H</sub> 3.17) to C-1 (δ<sub>C</sub> 72.4) established that lentztrehalose C also possesses a 2, 3-dihydroxy-3-methylbutoxy chain attached to the C-4' of trehalose, the same as lentztrehalose A. Newly appeared long-range correlations were observed from the H-3' (δ<sub>H</sub> 3.92) of trehalose to the oxymethine at C-2 (δ<sub>C</sub> 79.1) and from H-2 (δ<sub>H</sub> 3.79) to C-3' (δ<sub>C</sub> 76.1). These correlations established the presence of a six-membered ring structure between C-2 in the 2,3-dihydroxy-3-methylbutoxy moiety and the C-3' of trehalose with an ether linkage as shown in Figure 1. The NOEs observed between H-2 and H-3' showed the 2-hydroxy-2-propyl group of C-2 was in an equatorial orientation in the additional six-membered ring. The structure of lentztrehalose C was elucidated to be 3, 4-O-[(1*R*)-1-(2-hydroxy-2-propyl)-1, 2-ethanediyl] trehalose as shown in Figure 1. Although ~60–70 natural compounds other than lentztrehalose A are known to have a 2, 3-dihydroxy-3-methylbutoxy moiety, none of these compounds are reported to have a cyclized analog as observed for lentztrehalose C.

### Production of lentztrehaloses

The side chain of lentztrehalose B is likely to be derived from dimethylallyl pyrophosphate, which is made in most organisms by the mevalonate and non-mevalonate pathways. Because both trehalose and the side chain moiety are prevalent in many organisms, we considered that lentztrehaloses might be common compounds and searched for other lentztrehalose producing actinomycetes. Although more than 50 species of actinomycetes were cultured under the same conditions and the metabolites analyzed, no other species produced any of the lentztrehaloses A, B or C (data not shown). We theorized that there is a unique enzyme required to transfer dimethylallyl to the C-4 of trehalose that is found in the producing strain.

During the exploration of the optimal culture conditions for a high yield of lentztrehaloses, it was observed that the producing strain preferred rather dry conditions and excess humidity suppressed the production of lentztrehaloses (Figure 2). In a germinated brown rice medium (rice grain:water = 3:5), trehalose, and lentztrehaloses A, B and C were produced at approximately 5.0-, 2.0-, 1.5- and 0.8-mg per 1.0 g rice grain, respectively after 4 weeks' culture (Figure 2, 1 ×). When the water content was reduced by half, the culture seemed dry at 4 weeks and the yields of trehalose and the lentztrehaloses were not much changed (Figure 2, 1/2 ×). Contrarily, in the porridge-like medium, the yields of trehalose and the lentztrehaloses were greatly reduced (Figure 2, 2 ×). We next examined the addition of trehalose to the culture, because trehalose might be the direct material used for the synthesis of the lentztrehaloses and thus its addition increases the yield (Figure 2, 3% trehalose bars). The addition of trehalose slightly increased the yield of lentztrehalose A and decreased the yield of lentztrehalose B in 1 × and 1/2 × cultures. Thus, trehalose does not appear to be an efficient material for the synthesis of lentztrehaloses under these conditions. In 2 × culture, however, additional trehalose increased the yield of lentztrehalose A to a level comparable to drier conditions. The biosynthesis of the trehalose moiety appears to be the rate limiting step for lentztrehalose production in wet environmental conditions and the transfer of the side chain component seems independent of it. The higher yield in dry conditions suggests that the lentztrehaloses, as well as trehalose, are produced as humectants or protectants for the producing microorganisms.

### Enzyme stability of lentztrehaloses B and C

Unlike trehalose, all of the lentztrehaloses were minimally hydrolyzed by porcine kidney trehalase (Figure 3a). In our previous report,<sup>15</sup> lentztrehalose A slightly released glucose when a large amount was reacted with porcine kidney trehalase. This effect may have been caused by an impurity in the previous lot of lentztrehalose A, which might have contained trehalose or other oligo- or poly-saccharides. These impurities released glucose after reaction with trehalase or other glycosidases present as contaminants. In the more recent purification process, several more chromatography steps using silica, polyamine and activated carbon columns were employed to reduce contamination with other sugars. Therefore, it is concluded that pure lentztrehaloses are trehalase-stable. While lentztrehaloses A and C did not suppress the activity of trehalase up to 10 mM, lentztrehalose B weakly suppressed it at a high concentration (Figure 3b). The mechanism, however, is unclear and lentztrehalose B is not considered to be a specialized inhibitor of trehalase.


### Antioxidative activity of lentztrehalose B

As a part of the investigation into the properties of the lentztrehaloses, we examined the antioxidative activity using an oxygen radical absorbance capacity (ORAC) activity assay.<sup>16</sup> In the assay, fluorescein

Table 2 <sup>1</sup>H and <sup>13</sup>C NMR data of lentztrehaloses B, C and A in methanol-d<sub>4</sub>

Lentztrehalose B				Lentztrehalose C				Lentztrehalose A <sup>a</sup>				
Position	$\delta_C$ (multiplicity)	$\delta_H$ ( $J$ , in Hz)	NOE	HMBC	Position	$\delta_C$ (multiplicity)	$\delta_H$ ( $J$ , in Hz)	NOE	HMBC	Position	$\delta_C$ (multiplicity)	$\delta_H$ ( $J$ , in Hz)
1	70.1 (t)	4.17 (1H, dd, 7.1, 11.2)	4'	2, 3, 4'	1	72.4 (t)	3.63 (1H, dd, 1.6, 11.9)		4'	1	74.7 (t)	4.00 (1H, dd, 2.8, 10.6)
2	122.7 (d)	4.37 (1H, dd, 6.9, 11.2)	4'	2, 3, 4'	2	79.1 (d)	3.86 (1H, dd, 8.3, 11.7)		1, 3, 4, 5	2	78.2 (d)	3.54 (1H, dd, 2.8, 8.0)
3	137.6 (s)	5.37 (1H, m)	1, 5	4, 5	3	78.7 (s)	3.79 (1H, dd, 1.6, 8.5)			3	72.8 (s)	3.65 m
4	18.2 (q)	1.69 (3H, s)		2, 3, 5	4	23.4 (q)	1.16 (3H, s)		3, 5	4	26.4 (q)	1.18 (3H, s)
5	25.9 (q)	1.74 (3H, s)		2, 3, 4	5	25.0 (q)	1.37 (3H, s)		2, 3, 4	5	25.4 (q)	1.16 (3H, s)
1'	95.0 (d)	5.11 (1H, d, 3.7)	2', 5'	5', 1''	1'	94.6 (d)	5.14 (1H, d, 3.9)	2', 5', 1''	5', 3', 1''	1'	95.1 (d)	5.10 (1H, d, 3.8)
2'	73.5 (d)	3.48 (1H, dd, 3.7, 9.6)	1', 4'	3'	2'	71.8 (d)	3.44 (1H, dd, 3.9, 9.4)	1', 4'	3'	2'	73.3 (d)	3.50 (1H, dd, 3.8, 9.6)
3'	74.8 (d)	3.88 (1H, t, 9.5)	5'	2', 4'	3'	76.1 (d)	3.92 (1H, dd, 8.5, 9.2)	2, 1'	2, 1', 2', 4'	3'	74.4 (d)	3.90 (1H, t, 9.6)
4'	79.0 (d)	3.25 (1H, dd, 8.9, 10.0)	1, 2'	1, 3', 5', 6'	4'	79.0 (d)	3.17 (1H, dd, 8.3, 9.9)	1, 2', 6'	1, 4', 5', 6'	4'	80.4 (d)	3.28 (1H, t, 9.6)
5'	72.9 (d)	3.81 (1H, ddd, 2.1, 4.8, 10.1)			5'	72.2 (d)	3.81 (1H, ddd, 2.2, 5.2, 10.2)			5'	72.7 (d)	3.85 (1H, br dt, 3.3, 9.6)
6'	62.3 (t)	3.65 (1H, dd, 4.8, 11.9)			6'	62.5 (t)	3.64 (1H, dd, 5.2, 12.1)		4', 5'	6'	62.1 (t)	3.75 (1H, m)
1''	95.1 (d)	5.09 (1H, d, 3.7)	2'', 5''	4'	1''	94.8 (d)	3.73 (1H, dd, 2.2, 12.1)	1', 2'', 5''	4', 5'	1''	95.0 (d)	3.78 (1H, m)
2''	73.3 (d)	3.45 (1H, dd, 3.7, 9.6)	1'', 4''	3''	2''	73.2 (d)	3.48 (1H, dd, 3.7, 9.8)	1'', 4''	3''	2''	73.2 (d)	3.45 (1H, dd, 3.8, 9.6)
3''	74.6 (d)	3.78 (1H, t, 9.4)		2'', 4''	3''	74.5 (d)	3.82 (1H, t, 9.3)		1'', 2'', 4''	3''	74.5 (d)	3.77 (1H, m)
4''	72.0 (d)	3.32 (1H, dd, 9.1, 9.9)	2''	3'', 5'', 6''	4''	72.0 (d)	3.32 (1H, t, 9.2)	2'', 6''	3'', 5'', 6''	4''	71.9 (d)	3.31 (1H, t, 9.5)
5''	73.8 (d)	3.81 (1H, m)		4''	5''	73.8 (d)	3.80 (1H, m)			5''	73.8 (d)	3.79 (1H, m)
6''	62.7 (t)	3.67 (1H, dd, 5.5, 12.1)		5''	6''	62.7 (t)	3.67 (1H, dd, 5.2, 11.8)		4'', 5''	6''	62.6 (t)	3.67 (1H, dd, 4.7, 12.0)
		3.79 (1H, m)					3.79 (1H, dd, 2.2, 12.1)		4'', 5''			3.78 (1H, m)

<sup>1</sup>H (600 MHz) and <sup>13</sup>C (150 MHz). Chemical shifts in p.p.m. from TMS as internal standard.<sup>a</sup>Literature data.



**Figure 2** Effect of water content in the medium on the production of lentsztrehaloses. The lentsztrehalose producing strain ML457-mF8 was cultured in steamed germinated brown rice media at 30 °C for 4 weeks. The ratios of rice grain:water were 3:5 (1×), 3:2.5 (1/2×) and 3:10 (2×) for each medium. In another set of cultures, a water solution containing 3% trehalose was used instead of water. The culture was extracted with an equal amount of EtOH, and the yields of trehalose and lentsztrehaloses A, B and C (TRH and LTA, B and C) were evaluated with an HPLC system by using a hydrophilic interaction chromatography column and evaporative light scattering detector.

is oxidized by peroxy radicals and the attenuation of fluorescence is monitored in the presence of antioxidant samples or a water-soluble analog of Vitamin E Trolox as a control (Figure 4a). Trehalose and lentsztrehalose A did not show antioxidative activity and rather promoted the attenuation of fluorescence. Lentsztrehalose C faintly showed retardation of the attenuation at 100 μM and the ORAC value calculated was 8.1 μmol TE g<sup>-1</sup> (Figure 4a, Table 1). Lentsztrehalose B, however, clearly protected fluorescein from the attack of peroxy radicals and the ORAC value was 207.3 μmol TE g<sup>-1</sup>. Lentsztrehalose B is stable in a water solution at room temperature for 6 days and the sodium adducts at *m/z* 433 ([M+Na]<sup>+</sup>) and *m/z* 455 ([M+2Na-H]<sup>+</sup>) of lentsztrehalose B were only observed in positive ESI/TOF-MS analysis (Figure 4b). In 15% hydrogen peroxide solution, additional peaks at *m/z* 449, 467 and 483 appeared. We consider the double bond of lentsztrehalose B is necessary for the antioxidative activity and oxygen atoms or hydrogen peroxide were spontaneously added there and formed epoxidated (*m/z* 449) or dihydroxylated (*m/z* 467) structures. The dihydroxylated form is probably lentsztrehalose A. Lentsztrehalose B may be able to trap one more oxygen atom and form the structure represented by the peak at *m/z* 483, although where the oxygen atom is bound is as yet unclear.

### Induction of autophagy

Induction of autophagy in mammalian cells is a function of trehalose that has attracted increasing attention in recent years. We examined this function of the lentsztrehaloses. Human cancer cell lines are easy to handle and were employed in this study. Twelve strains of cells were treated with 100-mM trehalose and the expression levels of autophagy marker protein microtubule-associated protein 1 light chain 3 (LC3)

were checked by western blotting in a preliminary experiment (Supplementary Figure 20). Immediately after proLC3 is translated, the C-terminal fragment is cleaved to yield a cytosolic form LC3-I. LC3-I is further converted to an autophagosome-associating form LC-II.<sup>8</sup> Except for two strains whose LC3 expression levels were constitutively high, other strains showed a clear increase in the expression level of LC3-II. We selected two highly sensitive cell lines, melanoma cell Mewo and ovarian cancer cell OVK18, for evaluating lentsztrehaloses. As reported in the previous study,<sup>7</sup> trehalose required a concentration of approximately 100 mM to induce a clear increase in LC3 expression (Figure 5a). The lentsztrehaloses showed a similar pattern and treatment with only 100 mM could clearly increase the expression level of LC3. As seen in Mewo cells treated with 0.1–10-mM lentsztrehalose C, a concentration-dependent increase in LC3 expression was sometimes observed. It was, however, not always reproducible. The induction of autophagy was also evaluated by using another marker Cyto-ID.<sup>17</sup> This green fluorescent reagent turns bright in vesicles produced during autophagy. Although the fluorescence was observed in a few spots in the control cells, it spread over the whole cell, except for the nucleus, in Mewo and OVK18 cells treated with 100-mM trehalose or lentsztrehalose (Figure 5b). These data indicated that lentsztrehalose can induce autophagy in human cells at a comparable level to trehalose. Considering the activity of lentsztrehalose C, the hydroxyl groups on C-3' and C-4' in trehalose can be modified while keeping the autophagy-inducing activity and it should be possible to synthesize trehalose analogs with better activity and bioavailability.

### CONCLUSION


We identified two new analogs of lentsztrehalose and determined their structures. Lentsztrehaloses A and C are probably formed from prenyl ether lentsztrehalose B through an epoxide that is hydrolyzed to give isomer A or intramolecular nucleophilic attack of C'-3-OH results in isomer C. It is not clear if both lentsztrehaloses A and C are derived from a corresponding epoxide since the configuration of lentsztrehalose A at C-2 is not yet determined. The chemical synthesis of lentsztrehalose A is now under development at our institute and the configuration at C-2 will be determined in the near future and the structural relevance of lentsztrehaloses A and C will be discussed in due course. All three lentsztrehaloses, A, B and C, were trehalase-stable and showed a level of autophagy-inducing activity comparable to trehalose. As more stable analogs of trehalose and with comparable autophagy induction activities, lentsztrehaloses are promising drug candidates for the treatment of neurodegenerative diseases and diabetes, arteriosclerosis, cancer and heart disease, which may be cured or alleviated by the induction of autophagy.<sup>8</sup> In addition to its autophagy-inducing activity, lentsztrehalose B has antioxidative activity. Antioxidants are being studied as possible drugs for amyotrophic lateral sclerosis treatment<sup>18</sup> and the curative effects of lentsztrehalose B for the treatment of amyotrophic lateral sclerosis is of particular interest for future studies.

### MATERIALS AND METHODS

#### Isolation and structure determination of lentsztrehaloses B and C

The previously reported methods for the production and isolation of lentsztrehalose A were modified.<sup>15</sup> Strain ML457-mF8 was cultured in a steamed germinated brown rice medium (IRits, Tokyo, Japan, rice grain:water = 3:5, autoclaved at 121 °C for 20 min) for 3 weeks. The whole culture was extracted with twice the volume of MeOH and EtOH, sequentially. The extract was evaporated and separated with a DIAION HP20 (Mitsubishi Chemical, Tokyo, Japan) column in a batch process using water and 50% MeOH. Although





**Figure 3** Enzyme stability of lentztrehaloses. (a) Trehalose or lentztrehaloses were reacted with porcine kidney trehalase and the concentrations of released glucose were determined by a hexokinase assay. Mean  $\pm$  s.e. from triplicate determinations. The graphs of lentztrehaloses are overlapped at the bottom in the same scale with trehalose (upper left) and each graph of lentztrehalose is magnified respectively. TRH: trehalose, LTA, B, C: lentztrehaloses A, B, C. (b) The trehalase inhibitory effects of the lentztrehaloses were evaluated. Trehalase was reacted with 1-mM trehalose in the presence of 10-mM lentztrehaloses A, C, or each concentration of B. Mean  $\pm$  s.e. from triplicate determinations. \*The difference between control and 10-mM lentztrehalose B was significant ( $P < 0.05$ ), whereas the difference between control and 1-mM lentztrehalose B was not significant ( $P = 0.19$ ).

lentztrehalose A was purified from the water fraction, B and C were isolated from the 50% MeOH fraction. Each fraction was separated by sequential chromatography with an octadecyl silica (Hydrosphere C18, YMC, Kyoto, Japan) and polyamine (YMC-Pack Polyamine II, YMC) HPLC, silica gel (Wakosil C-300, Wako Pure Chemical Industries, Ltd, Osaka, Japan), activated carbon (Wako) and Sephadex LH-20 (GE Healthcare Japan, Hino, Tokyo, Japan) open columns. The elution solvents used were water and MeOH for octadecyl silica HPLC, acetonitrile and water for polyamine HPLC, BuOH: MeOH:water = 4:1:2 mixture for the silica column and MeOH for the activated carbon and LH-20 columns. For the purity or yield check, lentztrehaloses were separated by HPLC with a hydrophilic interaction chromatography column (XBridge Amide, Waters, Milford, MA, USA) by using a linear gradient of 90–50% acetonitrile and detected by an evaporative light scattering detector system (ELSD 2000ES, Alltech, Deerfield, IL, USA).

The structure and physicochemical properties of lentztrehaloses B and C were determined by spectrometric analyses performed as previously described<sup>15,19</sup> using the equipment as follows. Optical rotations were recorded on a P-1030 polarimeter (JASCO Inc., Tokyo, Japan). UV spectra were obtained on a U2800 spectrophotometer (Hitachi High-Technologies Corporation, Tokyo, Japan). <sup>1</sup>H (600 MHz) and <sup>13</sup>C (150 MHz) NMR spectra were measured in methanol-*d*<sub>4</sub> with a JNM-ECA600 spectrometer (JEOL Resonance Inc., Tokyo, Japan) using tetramethylsilane as an internal reference. HRESI-MS spectra were obtained on a LTQ Orbitrap XL mass spectrometer (Thermo Fisher Scientific, San Jose, CA, USA) or T-100LC TOF-MS (JEOL Ltd, Akishima, Tokyo, Japan). IR spectra were recorded on a FT/IR-4100 Fourier transform infrared spectrometer (JASCO). The novelty of the lentztrehaloses was confirmed with the databases Dictionary of Natural Products on DVD

(Chapman & Hall/CRC Press, Boca Raton, FL, USA) and SciFinder (Chemical Abstracts Service, Columbus, OH, USA).

#### Trehalase assay

Trehalase digestion experiments were performed as previously described.<sup>15</sup> The pH of 135-mM citrate buffer, however, was misdescribed as 7.5 in the previous report, the correct value is 5.7.

#### ORAC assay


The ORAC assay was performed using a OxiSelect Oxygen Radical Antioxidant Capacity Activity Assay kit (Cell Biolabs, Inc., San Diego, CA, USA) following the manufacturer's protocol. The fluorescence was detected with a fluorescence plate reader ARVO (PerkinElmer, Waltham, MA) using the filters Ex. 485 nm and Em. 535 nm.

#### Detection of autophagy


Human melanoma cell Mewo (JCRB0066) was obtained from the Health Science Research Resources Bank (Sennan, Japan). Human ovarian cancer cell OVK18 was provided by the RIKEN BRC (Tsukuba, Japan) through the National Bio-Resource Project of the MEXT, Japan. The cells were plated on 96-well microplates at  $1.6 \times 10^4$  cells per 0.2 ml per well in a Dulbecco's Modified Eagle Medium (Nissui, Tokyo, Japan) supplemented with 10% fetal calf serum (Sigma-Aldrich, St Louis, MO, USA) and 1% penicillin–streptomycin–glutamine solution (Thermo Fisher Scientific). The cells were cultured at 37°C for 72 h, and a half volume of medium was picked up to dissolve trehalose or the lentztrehaloses and put back in the original well to treat the cells with the sugars. After culturing for 24 h, the medium was removed and

cells were washed with phosphate-buffered saline (DS Pharma Biomedical, Suita, Osaka, Japan) once and lysed with a 2 × SDS-PAGE sample buffer (4% SDS, 0.125 M Tris-HCl (pH 6.8), 10% sucrose). The protein concentration was

measured by BCA assay (Thermo Fisher Scientific) and 2.5-μg protein/sample was applied on western blotting. Western blotting was basically performed as previously described<sup>20–22</sup> using anti-LC3 antibody (Medical & Biological



**Figure 4** Antioxidative activity of lentztrehalose B. (a) Antioxidative activities of the lentztrehaloses were evaluated by oxygen radical absorbance capacity activity assay. TRH: trehalose, LTA, B, C: lentztrehaloses A, B, C. (b) Lentztrehalose B was dissolved at 0.5 mM in water or 15% H<sub>2</sub>O<sub>2</sub>, and allowed to stand for 6 days at room temperature. The mass spectrum was analyzed with a time-of-flight MS.



**Figure 5** Autophagy inducing activity of lentztrehaloses. (a) Human melanoma cell Mewo and ovarian cancer cell OVK18 were treated with each concentration of trehalose (TRH) or lentztrehaloses (LTA, B and C) for 24 h and the expression level of autophagy markers LC3-I and II were observed by western blotting. The expression of  $\beta$ -tubulin was used as a control. The same 100-mM TRH sample was loaded on the different gels as a control. (b) Mewo and OVK18 cells were treated with 100-mM TRH or lentztrehaloses for 24 h and then autophagy marker dye Cyto-ID for 30 min. The vesicles formed during the autophagic process were stained brightly. PC, phase contrast.

Laboratories co., Ltd., Nagoya, Japan), anti- $\beta$  tubulin (Cell Signaling Tech., Danvers, MA, USA) and horseradish peroxidase conjugated anti-rabbit IgG (Cell Signaling Tech) at 1/2000. The signal was detected with Amersham ECL Prime Western Blotting Detection System (GE Healthcare). For the cell staining assay, the cells were plated at  $8 \times 10^3$  per 0.2 ml per well and cultured for 72 h. The cells were treated with trehalose or a lentztrehalose as in the case for western blotting. The autophagy detecting dye Cyto-ID (Enzo Life Sciences, Farmingdale, NY, USA) was diluted with Dulbecco's Modified Eagle Medium at 1/10, and the solution was added to the cell at 4  $\mu$ l per well and incubated at 37°C for 30 min. The excess dye was removed by washing with phosphate-buffered saline once and cells were observed with a fluorescent microscope ECLIPSE-TE2000-U (Nikon, Tokyo, Japan) using the B filter (Ex. 420–490 nm, Em. 520 nm).

#### ACKNOWLEDGEMENTS

We acknowledge the assistance of Dr K Iijima, Dr T Someno, Dr K Yamazaki and Dr Y Takahashi at the Institute of Microbial Chemistry for their support of this study and for helpful discussions. This study was partially supported by the Japan Society for the Promotion of Science.

- 1 Elbein, A. D., Pan, Y. T., Pastuszak, I. & Carroll, D. New insights on trehalose: a multifunctional molecule. *Glycobiology* **13**, 17R–27R (2003).
- 2 Ohtake, S. & Wang, Y. J. Trehalose: current use and future applications. *J. Pharm. Sci.* **100**, 2020–2053 (2011).

- 3 Nishizaki, Y. *et al.* Disaccharide-trehalose inhibits bone resorption in ovariectomized mice. *Nutr. Res.* **20**, 653–664 (2000).
- 4 Ukawa, Y., Gu, Y., Ohtsuki, M., Suzuki, I. & Hisamatsu, M. Antitumor effect of trehalose on sarcoma 180 in ICR mice. *J. Appl. Glycosci.* **52**, 367–368 (2005).
- 5 Arai, C. *et al.* Trehalose prevents adipocyte hypertrophy and mitigates insulin resistance. *Nutr. Res.* **30**, 840–848 (2010).
- 6 Tanaka, M. *et al.* Trehalose alleviates polyglutamine-mediated pathology in a mouse model of Huntington disease. *Nat. Med.* **10**, 148–154 (2004).
- 7 Sarkar, S., Davies, J. E., Huang, Z., Tunnacliffe, A. & Rubinsztein, D. C. Trehalose, a novel mTOR-independent autophagy enhancer, accelerates the clearance of mutant huntingtin and alpha-synuclein. *J. Biol. Chem.* **282**, 5641–5652 (2007).
- 8 Cheng, Y., Ren, X., Hait, W. N. & Yang, J. M. Therapeutic targeting of autophagy in disease: biology and pharmacology. *Pharmacol. Rev.* **65**, 1162–1197 (2013).
- 9 Chiti, F. & Dobson, C. M. Protein misfolding, functional amyloid, and human disease. *Annu. Rev. Biochem.* **75**, 333–366 (2006).
- 10 Aguib, Y. *et al.* Autophagy induction by trehalose counteracts cellular prion infection. *Autophagy* **5**, 361–369 (2009).
- 11 Schaeffer, V. *et al.* Stimulation of autophagy reduces neurodegeneration in a mouse model of human tauopathy. *Brain* **135**, 2169–2177 (2012).
- 12 Castillo, K. *et al.* Trehalose delays the progression of amyotrophic lateral sclerosis by enhancing autophagy in motoneurons. *Autophagy* **9**, 1308–1320 (2013).
- 13 Sarkar, S. *et al.* Neuroprotective effect of the chemical chaperone, trehalose in a chronic MPTP-induced Parkinson's disease mouse model. *Neurotoxicology* **44**, 250–262 (2014).
- 14 Oku, T. & Nakamura, S. Estimation of intestinal trehalase activity from a laxative threshold of trehalose and lactulose on healthy female subjects. *Eur. J. Clin. Nutr.* **54**, 783–788 (2000).
- 15 Wada, S. *et al.* Structure and biological properties of lentztrehalose: a novel trehalose analog. *J. Antibiot.* **66**, 319–322 (2014).
- 16 Cao, G., Alessio, H. M. & Cutler, R. G. Oxygen-radical absorbance capacity assay for antioxidants. *Free Radic. Biol. Med.* **14**, 303–311 (1993).
- 17 Chan, L. L. *et al.* A novel image-based cytometry method for autophagy detection in living cells. *Autophagy* **8**, 1371–1382 (2012).



- 18 Pandya, R. S. *et al*. Therapeutic neuroprotective agents for amyotrophic lateral sclerosis. *Cell. Mol. Life Sci.* **70**, 4729–4745 (2013).
- 19 Igarashi, M. *et al*. Waldiomycin, a novel WalK-histidine kinase inhibitor from *Streptomyces* sp. MK844-mF10. *J. Antibiot.* **66**, 459–464 (2013).
- 20 Wada, S. *et al*. *Candida glabrata* ATP-binding cassette transporters Cdr1p and Pdh1p expressed in a *Saccharomyces cerevisiae* strain deficient in membrane transporters show phosphorylation-dependent pumping properties. *J. Biol. Chem.* **277**, 46809–46821 (2002).
- 21 Wada, S. *et al*. Phosphorylation of *Candida glabrata* ATP-binding cassette transporter Cdr1p regulates drug efflux activity and ATPase stability. *J. Biol. Chem.* **280**, 94–103 (2005).
- 22 Wada, S. *et al*. Rubratoxin A specifically and potently inhibits protein phosphatase 2A and suppresses cancer metastasis. *Cancer Sci.* **101**, 743–750 (2010).



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Supplementary Information accompanies the paper on The Journal of Antibiotics website (<http://www.nature.com/ja>)