

ORIGINAL ARTICLE

The effect of red blood cell transfusion on intermittent hypoxemia in ELBW infants

EG Abu Jawdeh¹, RJ Martin¹, TE Dick², MC Walsh¹ and JM Di Fiore¹

OBJECTIVE: To test the hypothesis that the effect of red blood cell (RBC) transfusion on intermittent hypoxemia (IH) in extremely low birth weight (ELBW) infants is dependent on postnatal age.

STUDY DESIGN: Oxygen saturation of 130 ELBW infants, who required transfusion, was monitored continuously for the first 8 weeks of life. We compared the characteristics of IH ($\text{SpO}_2 \leq 80\%$ for ≥ 4 s and ≤ 3 min), 24 h before and both 24 h and 24 to 48 h after each RBC transfusion at three distinct time periods: Epoch 1, 1 to 7 days; Epoch 2, 8 to 28 days; and Epoch 3, > 28 days.

RESULT: In Epoch 1, the frequency and severity of IH events were not significantly different before and after transfusion. In both Epochs 2 and 3 there was a decrease in IH frequency and severity 24 h after RBC transfusion that persisted for 48 h. In addition, there was a decrease in the overall time spent with $\text{SpO}_2 \leq 80\%$ which persisted for 24 h after transfusion in Epochs 1 and 3, and for 48 h in Epoch 3.

CONCLUSION: The benefit of RBC transfusion on IH is age dependent as improvement in the frequency and severity of IH after transfusion only occurs beyond the first week of life. These observations will aid clinician's decision making by clarifying the benefit of RBC transfusions on patterns of oxygenation in preterm infants.

Journal of Perinatology (2014) **34**, 921–925; doi:10.1038/jp.2014.115; published online 12 June 2014

INTRODUCTION

Intermittent hypoxemia (IH) during early postnatal life is typically the result of impaired respiratory control often superimposed upon suboptimal lung function.¹ It may be associated with increased perinatal morbidities and poor neurodevelopmental outcome.^{2–4} Red blood cell (RBC) transfusions may ameliorate IH events by enhancing oxygen carrying capacity with resultant improvement in respiratory control and/or oxygen reserves.¹ Previous studies have assessed the relationship between RBC transfusions and apnea of prematurity with and without desaturation events and have reported conflicting results.^{5–16} Zagol *et al.*⁶ showed sustained improvement in apnea associated with oxygen desaturation after RBC transfusions. Seidel *et al.*¹⁶ also showed enhancement in oxygenation after RBC transfusions. In contrast, Westkamp *et al.*,¹⁴ Poets *et al.*¹³ and Stute *et al.*¹⁰ found no change in the incidence of IH events post RBC transfusion. These contradictory results may be due to small sample size, lack of continuous saturation monitoring, or broad postnatal and gestational age ranges.^{6,10,13,14,16}

Recently, we have documented that the incidence of IH changes over the first 2 months of life in a cohort of 79 preterm infants of 24 to 28 weeks gestation.³ During the first week of life IH occurs infrequently; in weeks 2 to 3, IH progressively increases, plateaus around 4 weeks; and decreases thereafter.³ Previous studies addressing the incidence of IH following RBC transfusion did not consider alterations in occurrence of IH with progressing postnatal age, which may have influenced their findings. Therefore, we sought to characterize the association between the effect of RBC transfusions on IH and postnatal age by testing the hypothesis that the effect of transfusion on IH in extremely low birth weight (ELBW) infants is dependent on postnatal age.

METHODS

Study design and data collection

To allow for a large sample size and, thus, an increased chance of detecting an effect of transfusion on the incidence of IH we examined two cohorts of preterm infants (24 to 27 6/7 weeks gestation) followed for the first 8 weeks of life. The cohorts were formed between 2005 and 2009 at Rainbow Babies and Children's Hospital Neonatal Intensive Care Unit. The first cohort comprised 79 infants who received standard clinical care, including the need for respiratory support, oxygen and caffeine.³ Standard patient care included a target SpO_2 between 85 and 95%. An additional cohort included 98 infants enrolled in the SUPPORT trial, a multi-centered randomized trial that compared two different target ranges of oxygen saturation ranges (85 to 89%, $n=49$ and 91 to 95%, $n=49$).^{17,18} Infants with congenital malformations were excluded.

Oxygen saturation was recorded continuously using high-resolution (2 s averaging time and a 0.5 Hz sampling rate) pulse oximetry (Radical; Masimo, Irvine, CA, USA) from day 1 to 8 weeks postnatal age. Intermittent hypoxemia was defined as a drop in $\text{SpO}_2 \leq 80\%$ for ≥ 4 s and ≤ 3 min duration. The lower threshold of 4 s was based on previous data by Poets *et al.*¹³ The upper limit in duration (3 min) was chosen arbitrarily to distinguish intermittent hypoxemia from sustained changes in baseline oxygenation. In addition, to account for all periods of hypoxemia, the total time of $\text{SpO}_2 \leq 80\%$ was calculated, which included both intermittent and sustained (> 3 min) hypoxemic events.

For each RBC transfusion, we documented the timing, volume and change in hematocrit. To avoid the potential effect of multiple transfusions, the RBC transfusion was considered eligible for analysis if there was no other transfusion within the 24 h before and 48 h after the transfusion. The 8-week monitoring period was stratified into three epochs based on the frequency of IH for all 177 infants (Figure 1). As previously published by Di Fiore *et al.*,³ we identified a low incidence of IH during the first week of postnatal age, followed by a progressive increase over weeks 2 to 3, a plateau around 4wks, and a decrease thereafter. Epoch 1 was defined as the first week of life when there are few IH events. Epoch 2 was

¹Division of Neonatology, Department of Pediatrics, Rainbow Babies and Children's Hospital, Case Western Reserve University, Cleveland, OH, USA and ²Division of Pulmonary and Critical Care Medicine, Department of Medicine, Case Western Reserve University, Cleveland, OH, USA. Correspondence: JM Di Fiore, Division of Neonatology, Department of Pediatrics, Rainbow Babies & Children's Hospital, Case Western Reserve University, 11100 Euclid Avenue, Suite 3100, Cleveland, OH 44106-6010, USA.

E-mail: jmd3@case.edu

Received 22 January 2014; revised 17 March 2014; accepted 5 May 2014; published online 12 June 2014

defined as 8 days to 4 weeks of life when there is an increase in the number of IH events. Epoch 3 was defined as beyond 4 weeks of life when the IH frequency plateaus and begins to decrease. For each infant only one RBC transfusion in each epoch was analyzed. In infants with multiple transfusions, the first eligible transfusion in each epoch was selected.

Statistical analysis

One-way analysis of variance with repeated measures was performed separately for each epoch to compare the incidence of hypoxic events in the 24-h period before, and 24 h and 24 to 48 h after the transfusion. *Post hoc* comparisons of pre and post transfusion time points (24 h pre, 24 h post and 24 to 48 h post) were performed by the Tukey–Kramer method. The oxygen saturation target range (wide (85 to 95%), high (91 to 95%) and low (85 to 89%)) was included as a covariate in the analysis of variance model to control for a potential effect of baseline oxygen saturation target. Generalized Estimating Equations (GEEs) logistic model or analysis of variance was used to compare baseline characteristics among epochs for categorical and continuous data, respectively.

Sample size

On the basis of previous data suggesting a 40% decrease¹⁴ in the number of IH events with an average standard deviation of 80 IH per day (Figure 1), we calculated that a sample size of 24 infants would be needed in each epoch to detect a change of this magnitude ($\alpha=0.05$, $1-\beta=0.80$). Accessibility to our large infant cohort enabled us to include 78 infants in Epoch 1, 109 infants in Epoch 2 and 75 infants in Epoch 3. The increased number of infants enrolled in this study enabled us to detect a smaller change of 22% in the incidence of IH ($\alpha=0.05$, $1-\beta=0.80$).

Institutional review board approval

The University Hospitals Institutional Review Board approved the study. As this is an expanded analysis of a previously obtained data set, a waiver of consent was granted.




Figure 1. The frequency of intermittent hypoxemia (IH) events during the first 8 weeks of postnatal life. The 8-week postnatal period was stratified into three epochs. Epoch 1 was defined as 1 to 7 days of life, Epoch 2 between 8 and 28 days of life and Epoch 3 after 28 days of life.

RESULTS

Of the 177 preterm infants included in the cohorts, 130 had eligible RBC transfusions in the first 8 weeks of life. Characteristics of the study population are presented in Tables 1 and 2, based on the epoch in which they were transfused. A total of 262 RBC transfusions met criteria for analysis. There was no difference in gestational age, birth weight, race or gender across epochs (Table 1). More infants in Epoch 1 were on ventilator support compared to Epochs 2 and 3 (Table 2). The majority of infants received supplemental oxygen encompassing approximately 90% of infants during each time period (Table 2). The administration of caffeine was significantly lower in Epoch 1 compared to Epochs 2 and 3 (Table 2). The hematocrit increased comparably after RBC transfusion in each of the three epochs (Figure 2). The median RBC transfusion volume was 20 ml kg⁻¹ administered over a duration of 4 h.

Table 2. Percent of infants receiving respiratory support

	Epoch 1	Epoch 2	Epoch 3
Ventilator support (%)	72	58 ^a	57 ^a
Non-invasive ventilation (%)	27	35	31
Other (nasal cannula, O ₂ hood) (%)	1	3	7
No respiratory support (%)	0	3	5
Respiratory support data not available (%)	0	1	0
Supplemental oxygen (%)	91	92	89
Caffeine (%)	62	81 ^a	79 ^a

^a $P < 0.05$ versus Epoch 1.




Figure 2. Hematocrit data presented in box-and-whisker diagram. There was no difference in pre and post red blood cell (RBC) transfusion hematocrit level across epochs. There was a significant increase in the hematocrit level post RBC transfusion in all three epochs ($*P < 0.001$). Box plot represent median and interquartile ranges. Whiskers represent the hematocrit range (minimum–maximum).

Table 1. Demographic data of the studied infants based on transfusion epoch

	Epoch 1	Epoch 2	Epoch 3
N	78	109	75
Gestational age	26 weeks (IQR 25–26)	26 weeks (IQR 25–27)	26 weeks (IQR 25–27)
Birth weight	730 g (IQR 655–806)	770 g (IQR 667–870)	730 g (IQR 635–840)
Male	51%	47%	48%
Black	58%	58%	60%
Post-natal age	4 days (IQR 2–5)	13 days (IQR 10–18)	33 days (IQR 31–38)
Weight at transfusion	665 g (IQR 610–760)	798 g (IQR 670–927)	1030 g (IQR 898–1260)

Abbreviation: IQR, interquartile range.




Figure 3. Characteristics of intermittent hypoxemia (IH) pre and post red blood cell (RBC) transfusion, including (a) frequency, (b) nadir and (c) duration. Frequency significantly decreased and IH nadir increased at 24 h and 24 to 48 h post RBC transfusion only in Epochs 2 and 3 (* $P < 0.02$ versus pre RBC transfusion). There was no change in IH duration.




Figure 4. Percent time with $SpO_2 < 80\%$ pre and post red blood cell (RBC) transfusion. The percent time with $SpO_2 \leq 80\%$ significantly decreased in Epochs 1 and 3 at 24 h post RBC transfusion persisting for 24 to 48 h in Epoch 3 (* $P < 0.05$). There was a trend toward a decrease in time with $SpO_2 \leq 80\%$ in Epoch 2 that did not reach statistical significance ($^{\dagger}P = 0.058$).

In Epoch 1, where minimal IH events occurred, there was no significant difference in the incidence, severity (nadir) and duration of IH before and after RBC transfusion (Figure 3). However, in both Epoch 2, in a postnatal age range where there is an increase in IH events, and Epoch 3, in a postnatal age range where the high incidence of IH events plateaus, there was a significant decrease in IH frequency 24 h after RBC transfusion that persisted for 48 h (all $P < 0.02$; Figure 3a). In addition, there was a decrease in the severity of IH (higher nadir) in Epochs 2 and 3 that persisted for 48 h (all $P < 0.02$; Figure 3b). There was no significant change in duration of IH with RBC transfusion (Figure 3c). We also examined the percent time spent with $SpO_2 \leq 80\%$ in all three epochs. In both Epochs 1 and 3 there was a significant improvement ($P < 0.05$) in the overall time spent with $SpO_2 \leq 80\%$ at 24 h after RBC transfusion persisting for 48 h in Epoch 3 only (Figure 4). In Epoch 2 the improvement in time with $SpO_2 \leq 80\%$ approached, but did not reach statistical significance ($P = 0.058$).

To achieve a large sample size to detect relatively small alterations in IH associated with transfusion, the cohort included three oxygen saturation target groups (wide, 85 to 95%; high, 91 to 95% or low, 85 to 89%). There was no significant effect of SpO_2 target range on the benefit of RBC transfusion (incidence, severity and duration) in any epoch.

DISCUSSION

Intermittent hypoxemic episodes are generally attributed to unstable respiratory control in preterm infants resulting in respiratory pauses and apnea.¹⁹ Two proposed mechanisms may underlie the beneficial effect of RBC transfusions on IH. The first suggests that anemia decreases oxygen delivery to the respiratory control network leading to hypoxic ventilatory depression.^{7,20–22} Thus, RBC transfusions may decrease apnea frequency by improving oxygen delivery to the immature brainstem. The second underlying mechanism is that RBC transfusions increase oxygen stores resulting in greater stability of oxygenation in the presence of apnea. This is consistent with a recent model analysis showing that the rate of arterial oxygen desaturation during apnea decreased with higher hemoglobin levels.¹ Therefore, the improvement in IH after RBC transfusions may be due to either reduction in the frequency of apnea or increased oxygen stores ameliorating IH in the presence of apnea.

Our results demonstrate a significant improvement in IH after RBC transfusion beyond the first week of life. The stratification of postnatal age into three developmental stages may partially explain the conflicting literature by minimizing the potential confounder of the dynamic natural history of IH. There was no benefit of RBC transfusion on frequency, severity or duration of IH in Epoch 1. This may be due to the already low incidence of IH before RBC transfusion in the first week of life. The mechanism underlying this lower incidence of IH in Epoch 1 is not clear, although higher ventilator use may have stabilized both lung volume and alveolar PO_2 , decreasing the likelihood of oxygen desaturation.¹ This explanation seems unlikely as addition of mechanical ventilator status to the statistical model did not affect the relationship between transfusion and IH events. As caffeine therapy improves respiratory control in preterm infants, the lack of benefit of transfusion on IH in Epoch 1 is unlikely due to caffeine

use as significantly fewer infants received caffeine compared to Epochs 2 and 3. Previous studies addressing the relationship between RBC transfusions and IH have primarily assessed infants with both a wide and/or older postnatal age.^{6,10,13,14,16} The mean postnatal age in studies that showed improvement in IH frequency after RBC transfusion^{6,16} corresponded to Epochs 2 and 3 in the current study. Apart from stratification by postnatal age our study had the additional advantage of a tight gestational age range (24 to 27 6/7 weeks).

Use of high-resolution (2 s averaging time and 0.5 Hz sampling rate) oxygen saturation monitoring optimized our ability to accurately identify IH events of both long and short duration. Vagedes et al.²³ have shown that detection of IH can vary up to fivefold with prolonged averaging times, as commonly used in the clinical setting, resulting in an artificial decrease in IH of 5–10 s and increase in IH of >20 s.²³ This may partially explain discrepancies between studies. Although there was no change in IH duration following RBC transfusion, there was an improvement in the total percent of time with SpO₂ ≤ 80%. This finding may be attributed to a decrease in the frequency of both IH and more prolonged periods (>3 min) of hypoxemia that were not included in the IH analysis. Given no improvement in IH after RBC transfusion in Epoch 1, the benefit in the percent of time with SpO₂ ≤ 80% may be due to a decrease in prolonged periods (>3 min) of hypoxemia. Recent data indicate that RBC transfusion may improve myocardial performance and possibly cardiac output in preterm infants, and we cannot exclude the possibility that this effect of simple volume expansion may have benefited oxygenation status of our infants.^{5,24}

A limitation of this study was the lack of accurate long-term apnea documentation which continues to be a challenge in the clinical setting. Bedside nursing documentation has been shown to underreport apnea events⁶ and impedance monitoring, the current clinical mode of monitoring respiration, cannot distinguish obstructive apnea from normal respiration. Therefore, long-term airflow measurements are needed to differentiate these two potential mechanisms, whereby RBC transfusion benefits IH episodes which is impractical in the clinical setting. However, we feel we have documented the concerning clinical component of apnea, namely resultant IH.¹⁹ Unfortunately available data do not allow us to document the response to IH (that is, supplemental oxygen), which may have affected our findings. The inclusion of three different oxygen saturation target groups may also have affected our study findings, although the effect of RBC transfusions on the hypoxemic events was not altered after adjusting for the oxygen saturation target. Including these multiple oxygen saturation target groups provided a large sample size that presented the ability to stratify the postnatal period and to detect more subtle alterations in IH patterns after RBC transfusion. Finally, we were unable to determine the precise indication for blood transfusion in many of the infants in this retrospective review, although anemia was the most common reason for transfusion. While the hematocrit threshold for transfusing preterm infants remains controversial,^{11,25,26} the hematocrit at time of transfusion in this study (range 20.7 to 38.8%) is comparable to the two largest randomized trials (hematocrit range 22 to 46%) assessing transfusion thresholds.^{11,25} The decision to transfuse varies among physicians and may also be influenced by the patient's daily condition. As the number of IH events was elevated during all epochs when compared to previously published data,³ it is conceivable that more patients may have been transfused on sick days with high number of IH events, followed by random improvement over the subsequent 48 h that is not related to the transfusion; however, this cannot explain the lack of benefit in Epoch 1.

In summary, this study has shown a clear benefit of RBC transfusion on the incidence and severity of IH episodes in preterm infants beyond the first week of life. These beneficial effects of

RBC transfusion in neonates should be weighed against potential risks such as transmitted infections,^{27,28} intraventricular hemorrhage^{29,30} and necrotizing enterocolitis.^{28,31} Future studies may allow us to determine whether there is a threshold of IH that is associated with neonatal morbidity.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We are thankful to the Neonatal Research Network SUPPORT trial. This study was supported by NINDS/NIH Grant R01 NS069220 (to TED).

REFERENCES

- Sands SA, Edwards BA, Kelly VJ, Davidson MR, Wilkinson MH, Berger PJ. A model analysis of arterial oxygen desaturation during apnea in preterm infants. *PLoS Comput Biol*. 2009; **5**: e1000588.
- Martin RJ, Wang K, Koroglu O, Di Fiore J, Kc P. Intermittent hypoxic episodes in preterm infants: do they matter? *Neonatology* 2011; **100**: 303–310.
- Di Fiore JM, Bloom JN, Orge F, Schutt A, Schluchter M, Cheruvu VK et al. A higher incidence of intermittent hypoxemic episodes is associated with severe retinopathy of prematurity. *J Pediatr*. 2010; **157**: 69–73.
- Janvier A, Khairy M, Kokkoti A, Cormier C, Messmer D, Barrington KJ. Apnea is associated with neurodevelopmental impairment in very low birth weight infants. *J Perinatol*. 2004; **24**: 763–768.
- Bifano EM, Smith F, Borer J. Relationship between determinants of oxygen delivery and respiratory abnormalities in preterm infants with anemia. *J Pediatr*. 1992; **120**: 292–296.
- Zagol K, Lake DE, Vergales B, Moorman ME, Paget-Brown A, Lee H et al. Anemia, apnea of prematurity, and blood transfusions. *J Pediatr*. 2012; **161**(417–421): e411.
- Joshi A, Gerhardt T, Shandloff P, Bancalari E. Blood transfusion effect on the respiratory pattern of preterm infants. *Pediatrics* 1987; **80**: 79–84.
- DeMaio JG, Harris MC, Deuber C, Spitzer AR. Effect of blood transfusion on apnea frequency in growing premature infants. *J Pediatr*. 1989; **114**: 1039–1041.
- Sasidharan P, Heimler R. Transfusion-induced changes in the breathing pattern of healthy preterm anemic infants. *Pediatr Pulmonol*. 1992; **12**: 170–173.
- Stute H, Greiner B, Linderkamp O. Effect of blood transfusion on cardiorespiratory abnormalities in preterm infants. *Arch Dis Child Fetal Neonatal Ed* 1995; **72**: F194–F196.
- Bell EF, Strauss RG, Widness JA, Mahoney LT, Mock DM, Seward VJ et al. Randomized trial of liberal versus restrictive guidelines for red blood cell transfusion in preterm infants. *Pediatrics* 2005; **115**: 1685–1691.
- Blank JP, Sheagren TG, Vajaria J, Mangurten HH, Benawra RS, Puppala BL. The role of RBC transfusion in the premature infant. *Am J Dis Child*. 1984; **138**: 831–833.
- Poets CF, Pauls U, Bohnhorst B. Effect of blood transfusion on apnoea, bradycardia and hypoxaemia in preterm infants. *Eur J Pediatr*. 1997; **156**: 311–316.
- Westkamp E, Soditt V, Adrian S, Bohnhorst B, Groneck P, Poets CF. Blood transfusion in anemic infants with apnea of prematurity. *Biol Neonate*. 2002; **82**: 228–232.
- Keyes WG, Donohue PK, Spivak JL, Jones MD Jr., Oski FA. Assessing the need for transfusion of premature infants and role of hematocrit, clinical signs, and erythropoietin level. *Pediatrics* 1989; **84**: 412–417.
- Seidel D, Blaser A, Gebauer C, Pulzer F, Thome U, Knapfer M. Changes in regional tissue oxygenation saturation and desaturations after red blood cell transfusion in preterm infants. *J Perinatol*. 2012; **33**: 282–287.
- Finer NN, Carlo WA, Walsh MCRich W, Gantz MG, Laptook AR et al. Early CPAP versus surfactant in extremely preterm infants. *N Engl J Med*. 2010; **362**: 1970–1979.
- Carlo WA, Finer NN, Walsh MC, Rich W, Gantz MG, Laptook AR et al. Target ranges of oxygen saturation in extremely preterm infants. *N Engl J Med*. 2010; **362**: 1959–1969.
- Abu-Shaweesh JM, Martin RJ. Neonatal apnea: what's new? *Pediatr Pulmonol* 2008; **43**: 937–944.
- Cross KW, Oppe TE. The effect of inhalation of high and low concentrations of oxygen on the respiration of the premature infant. *J Physiol*. 1952; **117**: 38–55.
- Rigatto H, Brady JP. Periodic breathing and apnea in preterm infants. II. Hypoxia as a primary event. *Pediatrics* 1972; **50**: 219–228.
- Rigatto H, Brady JP, de la Torre Verdusco R. Chemoreceptor reflexes in preterm infants: I. The effect of gestational and postnatal age on the ventilatory response to inhalation of 100% and 15% oxygen. *Pediatrics* 1975; **55**: 604–613.

- 23 Vagedes J, Poets CF, Dietz K. Averaging time, desaturation level, duration and extent. *Arch Dis Child Fetal Neonatal Ed* 2013; **98**: F265–F266.
- 24 Saleemi MS, El-Khuffash A, Kirkham C, Franklin O, Corcoran JD. Myocardial assessment using tissue doppler imaging in preterm very low-birth weight infants before and after red blood cell transfusion. *J Perinatol* 2013; **33**: 681–686.
- 25 Kirpalani H, Whyte RK, Andersen C, Asztalos EV, Heddle N, Blajchman MA *et al*. The Premature Infants in Need of Transfusion (PINT) study: a randomized, controlled trial of a restrictive (low) versus liberal (high) transfusion threshold for extremely low birth weight infants. *J Pediatr* 2006; **149**: 301–307.
- 26 Crowley M, Kirpalani H. A rational approach to red blood cell transfusion in the neonatal ICU. *Curr Opin Pediatr*. 2010; **22**: 151–157.
- 27 Martin RJ, Fanaroff AA, Walsh MC. *Fanaroff and Martin's neonatal-perinatal medicine: diseases of the fetus and infant* 9th ed. Mosby/Elsevier: St. Louis, MO, 2011.
- 28 Christensen RD, Ilstrup S. Recent advances toward defining the benefits and risks of erythrocyte transfusions in neonates. *Arch Dis Child Fetal Neonatal Ed* 2012; **98**: 365–372.
- 29 Baer VL, Lambert DK, Henry E, Snow GL, Butler A, Christensen RD. Among very-low-birth-weight neonates is red blood cell transfusion an independent risk factor for subsequently developing a severe intraventricular hemorrhage? *Transfusion* 2011; **51**: 1170–1178.
- 30 Christensen RD. Associations between 'early' red blood cell transfusion and severe intraventricular hemorrhage, and between 'late' red blood cell transfusion and necrotizing enterocolitis. *Semin Perinatol* 2012; **36**: 283–289.
- 31 Kirpalani H, Zupancic JA. Do transfusions cause necrotizing enterocolitis? The complementary role of randomized trials and observational studies. *Semin Perinatol* 2012; **36**: 269–276.