CORRIGENDUM

Transcutaneous bilirubin levels in newborns < 35 weeks' gestation

MJ Maisels, MP Coffey and E Kring

Journal of Perinatology (2015) **35**, 776. doi:10.1038/jp.2015.94**Correction to:** *Journal of Perinatology* advance online publication, 25 June 2015; doi:10.1038/jp.2015.34.

Following publication of the above article, the authors noticed an error in the postmenstrual ages in rows three and four of Table 4. The correct table appears here.

Table 4. TcB cutpoints that provide ≥ 98 –99% probability that the TSB is below the phototherapy level for each PMA category

Postmenstrual age	TcB (mg dl^{-1})	Phototherapy level (mg dl^{-1})	Probability (confidence interval)
28 0/7–29 6/7	≤ 2	6	98.1 (90.0–99.6)
	≤ 3	7	99.1 (93.0–99.9)
	≤ 4	7	98.2 (90.4–99.7)
30 0/7–31 6/7	≤ 5	8	99.6 (97.8–99.9)
	≤ 6	8	98.7 (95.5–99.6)
32 0/7–33 6/7	≤ 7	10	99.4 (97.5–99.8)
	≤ 8	10	98.5 (95.6–99.5)
34 0/7–34 6/7	≤ 10	12	99.0 (96.9–99.7)
	≤ 10	13	99.5 (97.3–99.9)
	≤ 11	13	99.0 (95.9–99.8)

Abbreviations: PMA, postmenstrual age; TcB, transcutaneous bilirubin; TSB, total serum bilirubin.