

In vivo and *in vitro* propagation of intraductal papillary mucinous neoplasms

Hirohiko Kamiyama^{1,4}, Mihoko Kamiyama^{1,4}, Seung-Mo Hong¹, Collins A Karikari¹, Ming-Tseh Lin¹, Michael W Borges¹, Margaret Griffith¹, Angela Young¹, Alexis Norris-Kirby¹, Conrad Lubek¹, Masamichi Mizuma¹, Georg Feldmann¹, Chanjuan Shi¹, Hong Liang¹, Michael G Goggins^{1,2}, Anirban Maitra^{1,3}, Ralph H Hruban^{1,3} and James R Eshleman^{1,3}

Intraductal papillary mucinous neoplasms (IPMNs) are one of the three known curable precursor lesions of invasive pancreatic ductal adenocarcinoma, an almost uniformly fatal disease. Cell lines from IPMNs and their invasive counterparts should be valuable to identify gene mutations critical to IPMN carcinogenesis, and permit high-throughput screening to identify drugs that cause regression of these lesions. To advance the study of the biological features of IPMNs, we attempted *in vivo* and *in vitro* growth of selected IPMNs based on the hypothesis that IPMNs could be grown in the most severely immunodeficient mice. We examined 14 cases by implanting them into nude, severe combined immunodeficient (SCID), and NOD/SCID/IL2R γ^{null} (NOG) mice, in addition to direct culture, to generate tumor xenografts and cell lines. One sample was directly cultured only. Thirteen tumors were implanted into the three types of mice, including 10 tumors implanted into the triple immunodeficient NOG mice, in which the majority (8 of 10) grew. This included five IPMNs lacking an invasive component. One of the explanted IPMNs, with an associated invasive carcinoma, was successfully established as a cell line. Tumorigenicity was confirmed by growth in soft agar, growth in immunodeficient mice, and the homozygous deletion of *p16/cdkn2a*. Epithelial differentiation of the cell line was documented by cytokeratin expression. Patient origin was confirmed using DNA fingerprinting. Most non-invasive IPMNs grow in NOG mice. We successfully established one IPMN cell line, and plan to use it to clarify the molecular pathogenesis of IPMNs.

Laboratory Investigation (2010) 90, 665–673; doi:10.1038/labinvest.2010.51; published online 15 March 2010

KEYWORDS: cell lines; immunodeficient mice; intraductal papillary mucinous neoplasm (IPMN); pancreatic cancer; precursor lesions

Pancreatic ductal adenocarcinoma (PDAC) is a notorious deadly cancer that affected an estimated 37 680 Americans in 2008, and resulted in approximately 34 290 deaths (case: fatality ratio = 91%).¹ Approximately 230 000 patients per year develop PDAC worldwide and the 5-year survival rate for these patients is expected to be only 4%.^{2,3} Effective early detection and treatment can improve these statistics, but require a full understanding of the molecular biology of the precursor lesions that give rise to invasive cancer. There are three documented morphologic precursors to pancreatic cancer: *pancreatic intraepithelial neoplasias* (PanINs), *intraductal papillary mucinous neoplasms* (IPMNs), and *mucinous cystic neoplasms*.⁴

IPMNs are mucin-producing epithelial neoplasms that, by definition, involve the main and/or branch pancreatic ducts and often, although not always, have a papillary architecture.^{4,5} It is clear that some IPMNs progress to invasive adenocarcinoma over time;^{6–9} however, several fundamental unanswered questions remain. The complete molecular pathogenesis, of invasive pancreatic cancers arising from IPMNs, is not established, as it is for those arising from PanINs.^{10,11} For example, although the progressive accumulation of mutations in the *KRAS2*, *TP53*, *p16/cdkn2a*, and *DPC4* genes has been well established for PanINs, a similar progression is not as well delineated for the various subtypes of IPMNs.^{12–17} In addition, different genes are sometimes

¹Department of Pathology, The Johns Hopkins University School of Medicine, Baltimore, MD, USA; ²Department of Medicine, The Johns Hopkins University School of Medicine, Baltimore, MD, USA and ³Department of Oncology, The Sol Goldman Pancreatic Cancer Research Center, The Johns Hopkins University School of Medicine, Baltimore, MD, USA

Correspondence: Dr JR Eshleman, MD, PhD, Department of Pathology, The Johns Hopkins University, Suite 341, David Koch CRB-II, 1550 Orleans Street, Baltimore, MD 21231 USA.

E-mail: jeshlema@jhmi.edu

⁴These two authors contributed equally to this work.

Received 1 July 2009; revised 22 December 2009; accepted 28 December 2009

targeted in PanINs and IPMNs. For example, the loss of the *STK11* gene is observed in up to one-third of IPMNs, but is rarely found in PanINs and PDACs.¹⁸ Similarly, activating mutations of the *PIK3CA* gene have been observed in IPMNs, but not in PanIN lesions.¹⁹ The natural history of IPMNs, such as the time and frequency of progression to PDAC, is also not well defined.^{9,20,21} Although the size of the lesion is associated with progression, it is unclear whether a specific size can be used as a clinical cutoff for surgical resection, although one has been proposed.²² In addition, no model exists in which to evaluate potential chemopreventative agents. IPMN cell lines would be valuable to clarify these issues. Thus, we used the techniques that have been used to establish cell lines from invasive pancreatic cancers to attempt to make IPMN cell lines.²³

In this report, we used triple immunodeficient NOG mice to propagate IPMNs, the majority of which grew. From one of these xenografted tumors, we aspirated the fluid from the cystic component, harvested the solid component separately, and generated cell lines from both of them. We conclude that IPMNs can be grown both *in vivo* and *in vitro* provided that the mice are sufficiently immunodeficient.

MATERIALS AND METHODS

Patients, Histopathology, and Tissue Harvest

On the basis of frozen section diagnoses, surgically resected samples were classified as IPMNs with or without an associated invasive carcinoma. Frozen section diagnoses were also confirmed with permanent sections. This work was performed with human subjects and animal committee approval. Freshly harvested IPMNs were implanted into mice and/or directly cultured within 2 h of resection.

In Vivo Growth as Mouse Xenografts

IPMNs were subcutaneously implanted in nude, severe combined immunodeficient (SCID) (*prkdc*^{null}), or NOG (NOD/*Prkdc*^{null}/IL-2R γ ^{null}) mice. Mice were monitored at regular intervals and killed when tumors reached about 1 cm³. Tumors were removed under sterile conditions and used for reimplantation, cryopreserved in DMSO, fixed in formaldehyde, and plated for tissue culture growth.

In Vitro Cell Culture

In a laminar flow biosafety cabinet, tumors explanted from mice or harvested directly in the surgical pathology suite were finely minced (<2 mm) and digested using minimum essential medium (MEM, GIBCO, Carlsbad, CA, USA) containing collagenase type 1 (1 mg/ml, GIBCO) and hyaluronidase (0.7 mg/ml, Sigma-Aldrich, St Louis, MO, USA). The single cells were cultured as described earlier,²³ with the following modifications. Cells were plated on 25 cm² flasks coated with rat-tail type 1 collagen (BD Biosciences, San Jose, CA, USA) and maintained in MEM containing 20% fetal bovine serum (FBS, GIBCO), 1% penicillin-streptomycin (GIBCO), 5 ng/ml EGF (GIBCO), and 0.2 U/ml human

recombinant insulin (GIBCO). In addition, to screen the optimal growth condition for each tumor, we varied several combinations for both the medium and substrates on 24-well plates for five cases.

To overcome fibroblast overgrowth, some cultures were treated periodically with selective trypsinization to remove the fibroblasts. Using a phase microscope, fibroblast-rich and tumor-rich regions were identified and marked on the bottom of the flask. Trypsin was added at room temperature and the culture was monitored. The reaction was stopped when the unwanted cells had detached, whereas the tumor cells were still attached, by aspirating the supernatant and quenching the trypsin by the addition of complete media.

Characterization of Cell Lines

Immunohistochemistry

Immunohistochemistry was used to detect the expression of cytokeratin (AE1 + AE3) (Ventana, Tucson, AZ, USA, PCK26 mouse monoclonal, predilute), vimentin (Ventana, V9 mouse monoclonal, predilute), MUC1 (Novocastra, New Castle on Tyne, UK, Ma552 mouse monoclonal, 1:200), MUC2 (Santa Cruz Biotechnology, Santa Cruz, CA, USA, H-300 rabbit polyclonal, 1:200), p16/CDKN2A (CINtec, Inc, Westborough, MA, USA, E6H4, mouse monoclonal, predilute), TP53 (Dako, Carpinteria, CA, USA, DO-7 mouse monoclonal, predilute), and DPC4/SMAD4 (Santa Cruz Biotechnology, B-8 mouse monoclonal, 1:200). The expression patterns in cell lines were compared with those observed in the primary IPMN and its xenograft. All the immunohistochemical reactions, except DPC4/SMAD4, were performed on a Ventana Benchmark XT, and all detection kits were from Ventana. Labeling for DPC4/SMAD4 protein was performed on a Dako autostainer using the DAKO Envision Plus detection kit.

Sequencing and multiple ligation-dependent probe amplification

The protein coding exons of *Kras*, *TP53*, *p16/CDKN2A*, and *DPC4/SMAD4*, in addition to exons 9 and 20 of the *PIK3CA* gene, were sequenced in the IPMN-1T and IPMN-1Asp cell lines, as described earlier.¹¹ Multiple ligation-dependent probe amplification (MLPA) was performed for *p16/CDKN2A* and *DPC4/SMAD4* using the SALSA MLPA kit (MRC-Holland, Amsterdam, Netherlands) as per manufacturer's instructions, and products were resolved on a Beckman-Coulter CEQ 8000 capillary electrophoresis instrument.

Fingerprinting

Microsatellite DNA fingerprinting was performed using the PowerPlex 1.2 system (Promega Corporation, Madison, WI, USA) at the Johns Hopkins University DNA core facility. The germline pattern was established using normal duodenum from the patient.

Soft agar assay

Anchorage independence was tested using an agar cloning assay with 0.5% agar with 100 000 suspended cells on 1% hard agar in six-well plates using six replicates for each sample. Spherical colonies, whose size was $>50\mu\text{m}$, were counted in 10 random fields using phase microscopy weekly. After 4 weeks, colonies were stained with crystal violet. HPDE is an immortalized normal epithelial pancreatic duct cell line and was used as a negative control.²⁴ Panc1 is a PDAC cell line and was used as the positive control.²⁵

Matrigel invasion assay

Matrigel invasion assays were performed using the BioCoat matrigel invasion chamber (BD Biosciences). Invasive cell derivatives, from largely non-invasive cells, were selected by plating 2.5×10^5 cells in 2 ml of MEM in the top chamber of a six-well plate and selecting cells from the bottom chamber at various times, as described in Results. The process was repeated to obtain second and third passage invasive cells.

Quantitative measurements were performed in 24-well plates to assess invasive capacity of IPMN-1T cells, IPMN-1T cells twice selected for invasion, and Panc1 control cells. For these assays, 2.5×10^4 cells, in 0.5 ml of serum-free MEM, were applied to the top chamber. The lower well, coated with collagen, was filled with complete culture medium containing 20% FBS. After 6–48 h of incubation, the non-invasive cells on the upper surface of the filter were removed with a cotton swab. The filters were fixed in methanol, stained with hematoxylin, and the cells were counted under a microscope at $\times 60$ magnification. The cells that had invaded through the matrigel were counted in 10 randomly selected fields, and the count numbers were averaged. The percent invasion was calculated as the number of cells that had invaded through the matrigel divided by the number of cells that had migrated through the filter without matrigel.

Tumor xenograft

Tumorigenicity in mice was confirmed by subcutaneously injecting approximately 20 million cells, from culture, bilaterally into two mice. Tumor volume (TV) was calculated according to the formula: $\text{TV (mm}^3\text{)} = \text{length} \times \text{width}^2 \times 0.5$. Explanted tumor xenografts were used for reimplantation and fixed in formalin for histology.

RESULTS

Case Selection and Primary Tumors

Owing to the need to fully understand the pathogenesis of IPMNs and their derivative invasive cancers, we attempted *in vivo* and *in vitro* propagation of these lesions. We initially attempted to grow IPMNs with and without an invasive component, but after initial success with *in vivo* propagation, we focused on those IPMNs without an invasive component (Table 1). Nine of 14 patients were male (64.2%), and the mean age was 66 years old. Five of the IPMNs that we

xenografted included an associated invasive cancer elsewhere in the lesion, and nine of the IPMNs were exclusively non-invasive. The non-invasive IPMN components were classified as high-grade dysplasia in seven cases (50%), moderate dysplasia in five (36%), and low-grade dysplasia in two (14%). The associated invasive component was a poorly differentiated adenocarcinoma in one case (20%), a moderately differentiated adenocarcinoma in three (60%), and a well-differentiated adenocarcinoma in one case (20%). The invasive adenocarcinomas were categorized as colloid type in one case and tubular type in four cases.

The results of the IHC labeling for cytokeratin, MUC1, MUC2, p16/CDKN2A, TP53, and DPC4/SMAD4, on the primary neoplasms, are shown in Table 2. Most of the invasive IPMNs were the pancreatobiliary subtype and produced associated tubular carcinomas (cases 1–4). Case 5 was the only intestinal subtype and formed a colloid carcinoma. In contrast, non-invasive IPMNs were either gastric or intestinal subtypes, except for case 8. Cytokeratin was positive and vimentin was negative for all cases. MUC expression patterns matched with the subtype, in which MUC1 was positive for the pancreatobiliary type, and MUC2 was positive for the intestinal type.²⁶ All non-invasive IPMNs expressed p16/CDKN2A and DPC4/SMAD4, but were negative for TP53. The invasive IPMNs showed a loss of p16/CDKN2A expression in three out of five cases, while TP53 was positive in one out of five cases. All five cases had intact DPC4/SMAD4.

In Vivo Growth of Human IPMNs in Immunodeficient Mice

We studied 14 total cases, of which only one was directly cultured (Table 3, case 11). Of the 13 IPMNs implanted in mice, one was implanted into a nude mouse, two were implanted into SCID mice, eight were implanted into triple immunodeficient NOG mice (Table 3, cases 2, 3, 4, 5, 6, 7, 9, and 14), and two were implanted into both SCID and NOG mice (Table 3, cases 8 and 12). Of the 13 implanted IPMNs, 11 grew as tumors in mice, whereas 2 did not grow. We attempted to culture, after explanting, 4 of the 11 tumors that grew as xenografts (Table 3, cases 1, 5, 8, and 9). The other seven mice died of infection or other causes before their tumors could be harvested.

In Vitro Growth of IPMNs After Explanting from Mice

After expansion in the immunodeficient mice, we explanted four cases for *in vitro* growth (Table 3). Another three cases were directly cultured from the surgical pathology suite (two of which were also grown in mice).

For case 1, the xenografted neoplasm formed a cyst. The fluid in the cyst was aspirated and cultured, and the resultant cell line was designated IPMN-1Asp, whereas the solid component of the same neoplasm was cultured and this cell line was designated IPMN-1T. Fibroblasts, which normally

Table 1 Clinicopathological findings of patients with IPMNs

Case	Sex	Age ^a	Diag ^b	Location/size(cm)/type ^c	Dysplasia	Invasion ^d	Pancreatectomy ^e
1	M	40s	IPMN/ml	Ph/2.5/Main	High	Mod/Tub	W
2	M	50s	IPMN/I	Ph,Pt/5.5,5.0/Main	High	Poor/Tub	TP
3	M	50s	IPMN/I	Ph/2.0/Main	High	Mod/Tub	W
4	M	80s	IPMN/I	Ph/6.0/Main	High	Mod/Tub	W
5	F	70s	IPMN/I	Ph/4.0/Branch	High	Well/Coll	W
6	F	80s	IPMN	Ph/3.0/Main	High	—	W
7	M	60s	IPMN	Ph/3.0/Main	Moderate	—	W
8	F	80s	IPMN	Ph/3.0/Main	Moderate	—	W
9	F	70s	IPMN	Ph/2.5/Main	High	—	W
10	F	40s	IPMN	Ph/1.5/Main	Moderate	—	W
11	M	60s	IPMN	Ph/3.0/Branch	Moderate	—	W
12	M	50s	IPMN	Pt/1.6/Main	Moderate	—	DP
13	M	60s	IPMN	Ph,Pb/1.5,1.0/Main	Low	—	W
14	M	70s	IPMN	Pb/6.0/Branch	Low	—	MP

^aAge in decade of life.

^bIPMN, IPMN without invasion; IPMN/I, IPMN with invasion; IPMN/ml, IPMN with microscopic invasion, <1 mm.

^cPh, pancreas head; Pb, pancreas body; Pt, pancreas tail; Main, main duct type; Branch, Branch duct type.

^dDifferentiation and type of invasive carcinoma. Mod, moderately differentiated; Poor, poorly differentiated; Well, well differentiated; Tub, tubular type; Coll, colloid type; —, not applicable.

^eW, whipple; TP, total; DP, distal; MP, middle.

Table 2 Immunohistochemical profile of primary IPMNs

Case	Invasion	Subtype ^a	Cytokeratin ^b	MUC1 ^b	MUC2 ^b	<i>p16</i>	<i>TP53</i> ^b	<i>DPC4/SMAD4</i>
1	Yes/MI ^c	PB/ONC	Exp	Exp	No exp	Lost ^d	No exp	Intact
2	Yes	PB	Exp	Exp	No exp	Lost	Exp	Intact
3	Yes	PB	Exp	Exp ^e	No exp	Lost	No exp	Intact ^f
4	Yes	GAST/PB	Exp	Exp ^g	No exp	Intact	No exp	Intact
5	Yes	INT	Exp	No exp	Exp	Intact	No exp	Intact
6	No	GAST/INT	Exp	No exp	Exp	Intact	No exp	Intact
7	No	GAST/INT	Exp	No exp	Exp ^g	Intact	No exp	Intact
8	No	GAST/PB	Exp	Exp ^g	No exp	Intact	No exp	Intact
9	No	INT	Exp	No exp	Exp	Intact	No exp	Intact
10	No	INT	Exp	No exp	Exp	Intact	No exp	Intact
11	No	GAST	Exp	No exp	No exp	Intact	No exp	Intact
12	No	INT	Exp	No exp	Exp	Intact	No exp	Intact
13	No	GAST	Exp	No exp	No exp	Intact	No exp	Intact
14	No	GAST	Exp	No exp	No exp	Intact	No exp	Intact

^aONC, oncocytic; PB, pancreatobiliary; GAST, gastric; INT, intestinal.

^bNo exp, no expression; exp, expression.

^cMI, microscopic <1 mm invasion.

^dFocally lost.

^eNon-invasive IPMN was negative, whereas invasive adenocarcinoma was positive.

^fNon-invasive IPMN was intact, whereas lost in invasive adenocarcinoma.

^gFocally positive expression.

Table 3 Strategy and results of *in vivo* and *in vitro* propagation

Case	Strategy (DC/mouse)	<i>In vivo</i> growth	<i>In vitro</i> growth
1	Nude	Growth	IPMN-1T, IPMN-1Asp
2	NOG	No growth	NC
3	NOG	Growth	NC
4	NOG	Growth	NC
5	NOG	Growth	IG
6	NOG	Growth	NC
7	NOG	Growth	NC
8	NOG, SCID	Growth	IG
9	NOG	Growth	IG
10	SCID	Growth	NC
11	DC	NI	IG
12	NOG, SCID	No growth	NC
13	DC, SCID	Growth	IG
14	DC, NOG	Growth	IG

DC, direct culture; NC, not cultured; IG, initial growth but failed to establish as a cell line; NI, not implanted.

overgrow in such cultures, were removed using selective trypsinization.²³ Neoplastic cells were successfully purified to homogeneity in the first passage for both IPMN-1T and IPMN-1Asp.

We had several problems producing cell lines from the other six cases. In case 5, the neoplastic cells did not attach, although fibroblasts attached and grew. In cases 9 and 11, some neoplastic cells attached and grew initially, but the growth rate was too slow and fibroblasts overgrew the culture in 2 weeks. In cases 13 and 14, neoplastic cells grew poorly and gradually died in primary cultures.

In five cases, we varied the basal media and substrate. Four types of basal medium were used, including MEM, DMEM, RPMI, and a 1:1 (volume:volume) mixture of MEM and RPMI supplemented as described in Materials and methods. For basal medium, the results using MEM and DMEM were equivalent and consistently superior to RPMI. Five different substrates were used: uncoated tissue culture plastic, glass cover slips, tissue culture plastic coated with rat-tail collagen, matrigel, or polylysine. In all cases, rat-tail collagen-coated flasks were much better than matrigel, polylysine, or uncoated flasks. Glass cover slips were the worst substrate. Fibroblast growth seemed to be stimulated on both matrigel




Figure 1 Histology and immunohistochemistry of matched primary tumor and corresponding xenograft for case 1. Hematoxylin and eosin (HE) and immunohistochemical labeling for cytokeratin, MUC1, MUC2, P16/cdkn2a, TP53, and DPC4/SMAD4. Negative region of P16/cdkn2a is shown for the primary tumor; however, the staining is heterogeneous as shown in Supplementary Figure 2. Magnification as indicated.

and polylysine. Selective trypsinization (see methods) helped to prevent fibroblast overgrowth.

Histological Comparison of Primary Tumors, Xenografts, and IPMN-1 Cell Lines

For case 1, we compared the histological and immunohistochemical findings in the matched primary tumor, the first passage xenograft, and the cell line (Figures 1 and 2). The primary tumor was a main duct-type IPMN with high-grade dysplasia. The subtype was focal oncocytic mixed with pancreatobiliary. Focally, the neoplastic cells formed small irregular nests in the extensive inflammatory stroma associated with the neoplasm, representing <1 mm microscopic invasion. The pathologic stage was T1N0MX. The first passage xenograft grew as an IPMN without invasion. Immunolabeling for cytokeratin was diffusely positive in all three samples (the primary tumor, the xenograft, and the cell line), whereas labeling for vimentin was consistently negative. The primary IPMN expressed MUC1, but did not express MUC2 (Figure 1), and this pattern was maintained in the xenograft and the cell lines (Figure 2). Immunolabeling for the p16/CDKN2A protein was generally lost in the primary tumor, in the xenograft, and in the cell lines, although there

was some heterogeneous expression in the primary tumor (see Supplementary Figure S1). The primary IPMN, the xenograft, and the cell lines did not stain with antibodies to the TP53 protein. Expression of the DPC4/SMAD4 protein was positive in the primary tumor, xenograft, and cell lines.

Case 8 was an IPMN with moderate dysplasia and was composed of pancreatobiliary and gastric subtypes. It did not have an associated invasive component. For MUC1, the primary IPMN was focally positive, and the xenograft was consistently positive. For MUC2, both the primary tumor and the xenograft were negative. Immunolabeling for the TP53 protein was negative in the primary tumor and the xenograft. In both the primary tumor and xenograft, P16 and DPC4/SMAD4 were intact (Supplementary Figure S2). We were unsuccessful at establishing a cell line for the explanted cells from case 8.

Genetic Characterization of IPMN-1 Cell Lines

Both IPMN-1T and IPMN-1Asp were wild type for *Kras* and *PIK3CA*. For *TP53*, there was only a germline SNP (P72R) and no somatic mutations were detected. For *p16/CDKN2A*, there was a homozygous deletion as shown by MLPA (Figure 3). This homozygous deletion is consistent with our




Figure 2 Phase microscopy and immunohistochemistry for the cell lines, IPMN-1T and IPMN-1Asp, stained for cytokeratin, MUC1, MUC2, p16/cdkn2a, TP53, and DPC4/SMAD4 ($\times 20$).




Figure 3 MLPA electropherograms of *p16/CDKN2A*. Negative wild-type control (**a**), positive control cell line with a homozygous deletion (**b**), and IPMN-1T (**c**).




Figure 4 IPMN-1 reimplantation. The tumor in NOG mouse approximately 12 weeks after injection with 20 million IPMN-1T cells (a). Histology of the reimplanted tumor ($\times 20$) showing a region without invasion (b,c). The third passage tumor in NOG mouse in 12 weeks after injection (d). Tumor shown in (d) after opening the skin and bisecting the tumor, revealing the nodular features of the tumor (e). Histology of the third passage tumor ($\times 20$) without evidence of invasion (f, g).

inability to amplify by PCR any of the exons, despite multiple attempts, and with the loss of immunohistochemical labeling. For *DPC4/SMAD4*, all exons sequenced as wild type, and no abnormalities were detected by MLPA (data not shown). These findings are consistent with retention of expression shown by IHC labeling. The results of cell line IPMN-1T and IPMN-1Asp both matched a non-neoplastic sample from the patient using DNA fingerprinting (Supplementary Table 1).

Tumorigenicity of the IPMN-1 Cell Lines

In anchorage-independent cloning assays, both IPMN-1Asp and IPMN-1T grew in soft agar and formed colonies at approximately equivalent rates. The frequency of colony formation was significantly less than that of the positive control Panc1 cells (Supplementary Figure S3), but higher than the negative control HPDE cells. We used invasion through matrigel to select an invasive subclone from the population of cells by plating them on top of the matrigel-coated filters, and growing the cultures until cells could be recovered from the bottom of the wells. Using this approach, invasive IPMN-1T cells were not detected until after 2 weeks of growth on the matrigel, whereas Panc1 invaded by 6 h. Invasive cells were expanded and the matrigel selection process was repeated.

We then tested the parental IPMN-1T, the second passage matrigel-selected derivative (IPMN-1T-M2), and Panc1 in the standard matrigel assay, which measures cell number on the bottom of a matrigel-coated filter at 6–72 h (Supplementary Figure S4). At 6 and 24 h time points, no cells had invaded from either the IPMN-1T or IPMN-1T-M2 cells, in contrast to the Panc1 cells. After 48 h, a few cells had invaded from both IPMN-1T and IPMN-1T-2M cultures.

In addition, to assess *in vivo* tumorigenicity, two NOG mice were subcutaneously injected with 20 million cells bilaterally, and one tumor was detected approximately 12

weeks after injection that measured 7×6 mm (126 mm^3 , Figure 4a). The tumor formed cysts and papillary structures as typically seen in IPMNs. Histology showed only non-invasive IPMN with high-grade dysplasia, with a pancreatobiliary subtype (Figure 4b and c). Half of this tumor was subcutaneously re-implanted into a third passage NOG mouse. After 14 weeks, the tumor had grown to 20×15 mm (2250 mm^3 , Figure 4d) and was explanted. The tumor formed a cystic mass, from which we aspirated approximately 4.5 ml of mucinous fluid. After aspirating the cyst fluid, the tumor was opened, and papillary nodules were revealed inside the cystic mass (Figure 4e). The histology (Figure 4f and g) showed an IPMN-like papillary structure with high-grade dysplasia, but without invasion.

DISCUSSION

We report that IPMNs can be consistently grown in NOG mice at high frequency (8/10, 80%), including those without an associated invasive component (5/8, 62%). We also successfully established the cell lines, IPMN-1T and IPMN-1Asp, from case 1, possibly because it contained a small invasive component. Tumorigenicity was confirmed by growth in soft agar and tumor production in second and third passage mice. In addition, the *p16/CDKN2A* gene was homozygously deleted. The epithelial differentiation of the line was confirmed by the expression of cytokeratin. DNA fingerprinting confirmed the patient from whom it was derived. Matrigel selection of a derivative line, albeit after 2 weeks of growth on the matrigel, suggests that there is a minor subpopulation of cells capable of invasion. This is consistent with the surgical pathology description of the primary tumor. However, we favor the notion that most cells in this mixed culture are non-invasive for the following reasons: because the original tumor histology was a 2.5 cm IPMN with only microscopic invasion of < 1 mm, as passage of this cell line in NOG mice showed

tumorigenicity without any invasion, and because the original cell line lacked invasion in the matrigel assay as typically performed.

We attempted to propagate six additional cases *in vitro*, but were unable to generate cell lines. The failure to generate cell lines fell into three general patterns: first, the neoplastic cells failed to attach; second, the primary cultures were initially successful, but the cell replication was extremely slow, and the cells gradually died; finally, the fibroblasts overgrew the neoplastic cells and destroyed them. To overcome these problems, we prepared a variety of conditions to find the optimal conditions for each culture. Despite the variety of approaches used, the substrate and growth medium may still lack essential growth factors. Finally, it may be impossible to produce cell lines from some IPMNs, because stromal cells may produce essential paracrine growth factors.²⁷

We further evaluated two of the available cases that grew in mice using immunohistochemical labeling and DNA sequencing. The patterns of protein expression in the primary tumor and xenograft generally matched. The expression of cytokeratin and vimentin in the cell lines IPMN-1T and IPMN-1Asp matched the expression observed in the paired primary tumor and xenograft samples. Expression of MUC was also consistent between the primary, xenograft, and cell lines. The p16/CDKN2A, TP53, and DPC4/SMAD4 expression patterns matched with their genetic characterizations.

Our work with IPMNs is actively ongoing. First, we are attempting to document the frequency of engraftment and rate of growth of IPMNs in nude, SCID, and NOG mice. Second, we are attempting to isolate invasive and non-invasive components from IPMN-1. In addition, we are continuing our efforts to establish IPMN cell lines from purely non-invasive IPMNs, and plan to use them to screen for chemoprevention agents.

Supplementary Information accompanies the paper on the Laboratory Investigation website (<http://www.laboratoryinvestigation.org>)

ACKNOWLEDGEMENTS

We acknowledge Dr Ming Tsao for generously providing the HPDE cell line. We also acknowledge Drs Elizabeth M Jaffee, Robert Anders, and Rajni Sharma for helpful discussions, in addition to Ms Guanglan Mo and Dante Trusty for helpful discussions and technical support. This work was funded, in part, by RO1CA130938 (JRE) and P50CA62924 (PI: Dr Scott Kern), the Stewart Trust, the Michael Rolfe Pancreatic Cancer Foundation, and the Mary Lou Wootton Pancreatic Cancer Research Fund.

DISCLOSURE/CONFLICT OF INTEREST

The authors declare no conflict of interest.

- Society AC. Cancer Facts and Figures 2008. American Cancer Society: Atlanta, 2008.
- Parkin DM, Bray F, Ferlay J, *et al*. Global cancer statistics, 2002. *CA Cancer J Clin* 2005;55:74–108.
- Jemal A, Murray T, Samuels A, *et al*. Cancer statistics, 2003. *CA Cancer J Clin* 2003;53:5–26.
- Hruban RH, Pitman MB, Klimstra DS. Tumors of the pancreas. Armed Forces Institute of Pathology, 4th edn. The American Registry of Pathology: Washington, DC, 2007, pp 75–164.
- Longnecker DS, Adler G, Hruban RH, *et al*. Intraductal Papillary-Mucinous Neoplasms of the Pancreas. WHO classification of tumor digestive system, IARC Press: Lyon, France, 2000, pp 237–240.
- Kimura W, Sasahira N, Yoshikawa T, *et al*. Duct-ectatic type of mucin producing tumor of the pancreas—new concept of pancreatic neoplasia. *Hepatogastroenterology* 1996;43:692–709.
- Nakagohri T, Kenmochi T, Kainuma O, *et al*. Intraductal papillary mucinous tumors of the pancreas. *Am J Surg* 1999;178:344–347.
- Terris B, Ponsot P, Paye F, *et al*. Intraductal papillary mucinous tumors of the pancreas confined to secondary ducts show less aggressive pathologic features as compared with those involving the main pancreatic duct. *Am J Surg Pathol* 2000;24:1372–1377.
- Yamaguchi K, Sugitani A, Chijiwa K, *et al*. Intraductal papillary-mucinous tumor of the pancreas: assessing the grade of malignancy from natural history. *Am Surg* 2001;67:400–406.
- Maitra A, Kern SE, Hruban RH. Molecular pathogenesis of pancreatic cancer. *Best Pract Res Clin Gastroenterol* 2006;20:211–226.
- Jones S, Zhang X, Parsons DW, *et al*. Core signaling pathways in human pancreatic cancers revealed by global genomic analyses. *Science* 2008;321:1801–1806.
- Z'Graggen K, Rivera JA, Compton CC, *et al*. Prevalence of activating K-ras mutations in the evolutionary stages of neoplasia in intraductal papillary mucinous tumors of the pancreas. *Ann Surg* 1997;226:491–498; discussion 498–500.
- Islam HK, Fujioka Y, Tomidokoro T, *et al*. Immunohistochemical study of genetic alterations in intraductal and invasive ductal tumors of the pancreas. *Hepatogastroenterology* 2001;48:879–883.
- Sasaki S, Yamamoto H, Kaneto H, *et al*. Differential roles of alterations of p53, p16, and SMAD4 expression in the progression of intraductal papillary-mucinous tumors of the pancreas. *Oncol Rep* 2003;10:21–25.
- Biankin AV, Biankin SA, Kench JG, *et al*. Aberrant p16(INK4A) and DPC4/Smad4 expression in intraductal papillary mucinous tumours of the pancreas is associated with invasive ductal adenocarcinoma. *Gut* 2002;50:861–868.
- Wada K. p16 and p53 gene alterations and accumulations in the malignant evolution of intraductal papillary-mucinous tumors of the pancreas. *J Hepatobiliary Pancreat Surg* 2002;9:76–85.
- Iacobuzio-Donahue CA, Klimstra DS, Adsay NV, *et al*. Dpc-4 protein is expressed in virtually all human intraductal papillary mucinous neoplasms of the pancreas: comparison with conventional ductal adenocarcinomas. *Am J Pathol* 2000;157:755–761.
- Sahin F, Maitra A, Argani P, *et al*. Loss of Stk11/Lkb1 expression in pancreatic and biliary neoplasms. *Mod Pathol* 2003;16:686–691.
- Schonleben F, Qiu W, Ciau NT, *et al*. PIK3CA mutations in intraductal papillary mucinous neoplasm/carcinoma of the pancreas. *Clin Cancer Res* 2006;12:3851–3855.
- Chari ST, Yadav D, Smyrk TC, *et al*. Study of recurrence after surgical resection of intraductal papillary mucinous neoplasm of the pancreas. *Gastroenterology* 2002;123:1500–1507.
- Salvia R, Fernandez-del Castillo C, Bassi C, *et al*. Main-duct intraductal papillary mucinous neoplasms of the pancreas: clinical predictors of malignancy and long-term survival following resection. *Ann Surg* 2004;239:678–685; discussion 685–687.
- Tanaka M, Chari S, Adsay V, *et al*. International consensus guidelines for management of intraductal papillary mucinous neoplasms and mucinous cystic neoplasms of the pancreas. *Pancreatol* 2006;6:17–32.
- Jaffee EM, Schutte M, Gossett J, *et al*. Development and characterization of a cytokine-secreting pancreatic adenocarcinoma vaccine from primary tumors for use in clinical trials. *Cancer J Sci Am* 1998;4:194–203.
- Furukawa T, Duguid WP, Rosenberg L, *et al*. Long-term culture and immortalization of epithelial cells from normal adult human pancreatic ducts transfected by the E6E7 gene of human papilloma virus 16. *Am J Pathol* 1996;148:1763–1770.
- Lieber M, Mazzetta J, Nelson-Rees W, *et al*. Establishment of a continuous tumor-cell line (panc-1) from a human carcinoma of the exocrine pancreas. *Inter J Canc* 1975;15:741–747.
- Furukawa T, Kloppel G, Volkan Adsay N, *et al*. Classification of types of intraductal papillary-mucinous neoplasm of the pancreas: a consensus study. *Virchows Arch* 2005;447:794–799.
- Ricci F, Kern SE, Hruban RH, *et al*. Stromal responses to carcinomas of the pancreas: juxtatumoral gene expression conforms to the infiltrating pattern and not the biologic subtype. *Cancer Biol Ther* 2005;4:302–307.