

PERSPECTIVE

Perspective on Diamond–Blackfan anemia: lessons from a rare congenital bone marrow failure syndrome

KM Sakamoto and A Narla

Leukemia (2018) 32, 249–251; doi:10.1038/leu.2017.314

Congenital and inherited bone marrow failure syndromes are rare and not exactly public health hazards. However, they have important lessons to teach about hematology in particular and cell biology in general. First, significant advances in scientific knowledge and understanding of pathogenesis can be obtained from studying rare diseases in children. Second, studies on rare syndromes provide insights into the normal physiology at both the cellular and global levels. Finally, as in the case of del(5q) myelodysplastic syndrome (where RPS14 was found to be mutated), discoveries from studying rare diseases such as Diamond–Blackfan anemia (DBA) can have important implications for our understanding of other congenital, inherited and acquired diseases in children and adults. In this perspective we briefly review progress and milestones in DBA research over the past two decades.¹

DBA was first described in 1936 by Hugh Josephs but was ultimately named after Louis Diamond and Kenneth Blackfan who described the hypoplastic anemia syndrome in 1938.² Initially, DBA was thought to be an immune-mediated disease that led to corticosteroid therapy that has become the standard of care. However, after 50 years, it is still unclear how precisely corticosteroids improve erythropoiesis in children with DBA. An elevated eADA has been noted since the 1980s in the majority (>75%) of patients with DBA.^{3–5} Why adenosine deaminase should be elevated and what role it might have in DBA again remains uncertain. In 1997, a mutation was noted on chromosome 19 in a child with DBA,^{6,7} and in 1999 was determined to be within the gene encoding *ribosomal protein S19 (RPS19)*. Mutations in *RPS19* were subsequently detected in ~25% of children with DBA.⁸ Studies of ribosomal protein encoding genes were then extended. We now know that ~70% of individuals with DBA have ribosomal protein (RP) mutations. DBA was the first disease to be linked to impaired ribosome function and is the founding member of a group of disorders now known as ribosomopathies.⁹ Ribosomal insufficiency associated with small ribosomal subunits leads to increased ratio of 28S/18S ribosomal RNA, whereas mutations in large ribosomal subunit results in decreased ratio of 28S/18S ribosomal RNA and nucleolar stress.^{10,11} Ebert and co-workers^{12,13} also identified RPS14 insufficiency in individuals with myelodysplastic syndrome and anemia, suggesting ribosome insufficiency contributes to acquired and congenital bone marrow failure syndromes. Exactly how RP insufficiency results in anemia and bone marrow failure is another unknown fact that is under study.

Progress in DBA research has come from the development of *in vitro* and *in vivo* model systems. The discovery of Yamanaka factors enabled scientists to generate pluripotent stem cells from

children with DBA that could then be differentiated into erythroid progenitor cells.^{14,15}

Mice with ribosomal mutations are smaller, have dark skin and decreased numbers of erythrocytes.¹⁶ p53 protein accumulates in epidermal melanocytes and induces C-Kit ligand expression in these mice. Zebrafish embryos injected with morpholinos recapitulate the congenital abnormalities found in children with DBA such as small size and severe anemia.^{17,18} Like humans, red blood cell adenosine deaminase levels are increased in these mouse and zebrafish models. p53 is upregulated in the zebrafish embryos resulting in apoptosis. Another important *in vivo* model recently described is the inducible RPS19 knockdown mouse that has an anemia phenotype that will be a model in which to test potential therapies.¹⁹

Genome sequencing of RNA and DNA has contributed to important advances in DBA research. These studies identified novel RP mutations in children with DBA, highlighting the importance of banking clinical samples.¹ DNA sequencing information has led to identification of certain phenotype–genotype correlations, such as *RPL5/RPL11* mutations and craniofacial abnormalities.¹ *GATA1* mutations are also found in some children with DBA.^{20,21} The inability of *GATA1* to normally regulate erythropoiesis in DBA may result from failure to interact with erythroid-specific promoter elements.²² The DBA Mutation Database and the DBA Registry will undoubtedly help identify new mutations in the future.²³

Going forward, improved genomic sequencing will likely uncover new genes and pathways critical to the pathogenesis of DBA. RNA-sequencing approaches have identified aberrantly regulated genes in bone marrow progenitors of children with DBA.^{24,25} Such studies raise important issues regarding the complexity of differential gene expression in diverse cell populations, culture conditions and bioinformatics data analyses.

Additional insights into DBA come from studies of iron metabolism. For example, an imbalance in heme/globin ratio seems to contribute to impaired erythropoiesis in DBA and myelodysplastic syndrome.²⁶ These data suggested that drugs that suppress heme synthesis might benefit individuals with these disorders. More recently, the use of inducible pluripotent stem cells from individuals with DBA were used to perform an unbiased chemical screen, resulting in the identification of an inducer of autophagy that stimulated erythropoiesis and upregulated globin genes.²⁷ As technology develops the hope is that new drugs will be identified to help individuals with DBA.

Finally, individuals with DBA have an increased risk of developing malignancies, suggesting the need to study

predisposing factors that may have broader implications.^{28–31} Recent analysis from the DBA Registry of North America (DBAR) reported 702 subjects with 12 376 individual-years of follow-up. The incidence of cancer was significantly increased with a 4.75 observed-to-expected ratio for all combined cancers. Significant observed-to-expected ratios were 352 for myelodysplastic syndrome, 45 for colony carcinoma, 42 for osteogenic sarcoma and 29 for acute myelogenous leukemia.³² The overall cancer risk appears to be approaching those observed in dyskeratosis congenital but lower than in Fanconi anemia. Interestingly, the risk of developing specifically gastrointestinal carcinoma and osteogenic sarcoma is higher in DBA. Studies have not found any genotype–phenotype correlation in terms of cancer risk.³²

SUMMARY AND FUTURE DIRECTIONS

Despite substantial advances in our understanding of the molecular aspects of DBA, many questions remain. We do not understand why a germline mutation of RPS19 should affect specific cells, in particular, erythroid progenitors. Future directions will likely focus on the role of inflammation and the bone marrow cytokine milieu that could also contribute to aberrant erythropoiesis.³³ The role of ribosome insufficiency and specific signaling pathways resulting in ineffective erythropoiesis are not well understood. The noncoding RNAs and epigenetic regulation of gene expression in DBA is not well studied.³⁴ Also unstudied is the role of the ubiquitin–proteasome system and its regulation of erythropoiesis.^{35,36}

In summary, therapy of children with DBA has not advanced for decades. Corticosteroids, red blood cell transfusions and hematopoietic cell transplants are the only currently effective therapies. Each is associated with significant morbidity and mortality. We hope future research will uncover new, effective therapies and will improve the quality of life of DBA.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank Bert Glader, Shuo Lin and Mark Wilkes for critical comments on this manuscript. KMS is supported by NIH R01DK107286 and Child Health Research Institute. AN is supported by the American Society of Hematology and Child Health Research Institute.

REFERENCES

- Boria I, Garelli E, Gazda HT, Aspesi A, Quarello P, Pavesi E *et al.* The ribosomal basis of Diamond-Blackfan Anemia: mutation and database update. *Hum Mutat* 2010; **31**: 1269–1279.
- Kaushansky K, Williams WJ. *Williams Hematology*, 8th edn. McGraw-Hill Medical, 2010.
- Fargo JH, Kratz CP, Giri N, Savage SA, Wong C, Backer K *et al.* Erythrocyte adenosine deaminase: diagnostic value for Diamond-Blackfan anaemia. *Br J Haematol* 2013; **160**: 547–554.
- Glader BE, Backer K. Elevated red cell adenosine deaminase activity: a marker of disordered erythropoiesis in Diamond-Blackfan anaemia and other haematologic diseases. *Br J Haematol* 1988; **68**: 165–168.
- Narla A, Davis NL, Lavasseur C, Wong C, Glader B. Erythrocyte adenosine deaminase levels are elevated in Diamond Blackfan anemia but not in the 5q- syndrome. *Am J Hematol* 2016; **91**: E501–E502.
- Gustavsson P, Willing TN, van Haeringen A, Tchernia G, Dianzani I, Donner M *et al.* Diamond-Blackfan anaemia: genetic homogeneity for a gene on chromosome 19q13 restricted to 1.8 Mb. *Nat Genet* 1997; **16**: 368–371.
- Gustavsson P, Skeppner G, Johansson B, Berg T, Gordon L, Kreuger A *et al.* Diamond-Blackfan anaemia in a girl with a de novo balanced reciprocal X;19 translocation. *J Med Genet* 1997; **34**: 779–782.

- Draptchinskaia N, Gustavsson P, Andersson B, Pettersson M, Willig TN, Dianzani I *et al.* The gene encoding ribosomal protein S19 is mutated in Diamond-Blackfan anaemia. *Nat Genet* 1999; **21**: 169–175.
- Raiser DM, Narla A, Ebert BL. The emerging importance of ribosomal dysfunction in the pathogenesis of hematologic disorders. *Leuk Lymphoma* 2014; **55**: 491–500.
- Quarello P, Garelli E, Carando A, Mancini C, Foglia L, Botto C *et al.* Ribosomal RNA analysis in the diagnosis of Diamond-Blackfan Anaemia. *Br J Haematol* 2016; **172**: 782–785.
- Ellis SR. Nucleolar stress in Diamond Blackfan anemia pathophysiology. *Biochim Biophys Acta* 2014; **1842**: 765–768.
- Ebert BL, Pretz J, Bosco J, Chang CY, Tamayo P, Galili N *et al.* Identification of RPS14 as a 5q- syndrome gene by RNA interference screen. *Nature* 2008; **451**: 335–339.
- Dutt S, Narla A, Lin K, Mullally A, Abayasekara N, Megerdichian C *et al.* Haploinsufficiency for ribosomal protein genes causes selective activation of p53 in human erythroid progenitor cells. *Blood* 2011; **117**: 2567–2576.
- Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 2006; **126**: 663–676.
- Garcon L, Ge J, Manjunath SH, Mills JA, Apicella M, Parikh S *et al.* Ribosomal and hematopoietic defects in induced pluripotent stem cells derived from Diamond Blackfan anemia patients. *Blood* 2013; **122**: 912–921.
- McGowan KA, Li JZ, Park CY, Beaudry V, Tabor HK, Sabnis AJ *et al.* Ribosomal mutations cause p53-mediated dark skin and pleiotropic effects. *Nat Genet* 2008; **40**: 963–970.
- Danilova N, Sakamoto KM, Lin S. Ribosomal protein S19 deficiency in zebrafish leads to developmental abnormalities and defective erythropoiesis through activation of p53 protein family. *Blood* 2008; **112**: 5228–5237.
- Taylor AM, Zon LI. Modeling Diamond Blackfan anemia in the zebrafish. *Semin Hematol* 2011; **48**: 81–88.
- Jaako P, Flygare J, Olsson K, Quere R, Ehinger M, Henson A *et al.* Mice with ribosomal protein S19 deficiency develop bone marrow failure and symptoms like patients with Diamond-Blackfan anemia. *Blood* 2011; **118**: 6087–6096.
- Sankaran VG, Ghazvinian R, Do R, Thiru P, Vergilio JA, Beggs AH *et al.* Exome sequencing identifies GATA1 mutations resulting in Diamond-Blackfan anemia. *J Clin Invest* 2012; **122**: 2439–2443.
- Ludwig LS, Gazda HT, Eng JC, Eichhorn SW, Thiru P, Ghazvinian R *et al.* Altered translation of GATA1 in Diamond-Blackfan anemia. *Nat Med* 2014; **20**: 748–753.
- Chlon TM, McNulty M, Goldenson B, Rosinski A, Crispino JD. Global transcriptome and chromatin occupancy analysis reveal the short isoform of GATA1 is deficient for erythroid specification and gene expression. *Haematologica* 2015; **100**: 575–584.
- Mirabello L, Khincha PP, Ellis SR, Giri N, Brodie S, Chandrasekharappa SC *et al.* Novel and known ribosomal causes of Diamond-Blackfan anaemia identified through comprehensive genomic characterisation. *J Med Genet* 2017; **54**: 417–425.
- O'Brien KA, Farrar JE, Vlachos A, Anderson SM, Tsujiura CA, Lichtenberg J *et al.* Molecular convergence in ex vivo models of Diamond-Blackfan anemia. *Blood* 2017; **129**: 3111–3120.
- Ulirsch JC, Lareau C, Ludwig LS, Mohandas N, Nathan DG, Sankaran VG. Confronting in ex vivo models of Diamond-Blackfan anemia. *Blood* 2017; **130**: 1165–1168.
- Yang Z, Keel SB, Shimamura A, Liu L, Gerds AT, Li HY *et al.* Delayed globin synthesis leads to excess heme and the macrocytic anemia of Diamond Blackfan anemia and del(5q) myelodysplastic syndrome. *Sci Transl Med* 2016; **8**: 338ra–367r.
- Doulatov S, Vo LT, Macari ER, Wahlster L, Kinney MA, Taylor AM *et al.* Drug discovery for Diamond-Blackfan anemia using reprogrammed hematopoietic progenitors. *Sci Transl Med* 2017; **9**(pii): eaah5645.
- Savage SA, Dufour C. Classical inherited bone marrow failure syndromes with high risk for myelodysplastic syndrome and acute myelogenous leukemia. *Semin Hematol* 2017; **54**: 105–114.
- Lipton JM, Federman N, Khabbaze Y, Schwartz CL, Hilliard LM, Clark JI *et al.* Osteogenic sarcoma associated with Diamond-Blackfan anemia: a report from the Diamond-Blackfan Anemia Registry. *J Pediatr Hematol Oncol* 2001; **23**: 39–44.
- Vlachos A, Rosenberg P, Kang J, Atsidaftos E, Alter B, Lipton J. Cancer predisposition in Diamond Blackfan Anemia: an update from the Diamond Blackfan Anemia Registry. *Pediatr Blood Cancer* 2016; **63**: S61–S61.
- Vlachos A, Rosenberg P, Kang J, Atsidaftos E, Alter B, Lipton J. Gastrointestinal cancers, osteogenic sarcoma and myelodysplastic syndrome predominate in Diamond Blackfan Anemia. *Pediatr Blood Cancer* 2017; **64**: S45–S45.
- Vlachos A, Rosenberg PS, Kang J, Atsidaftos E, Alter BP, Lipton JM. Myelodysplastic syndrome and gastrointestinal carcinomas characterize the cancer risk in Diamond Blackfan Anemia: a report from the Diamond Blackfan Anemia Registry. *Blood* 2016; **128**. Available at: <http://www.bloodjournal.org/content/128/22/3333/tab-article-info>.

- 33 Schepers K, Campbell TB, Passegue E. Normal and leukemic stem cell niches: insights and therapeutic opportunities. *Cell Stem Cell* 2015; **16**: 254–267.
- 34 Wilkes MC, Repellin CE, Sakamoto KM. Beyond mRNA: the role of non-coding RNAs in normal and aberrant hematopoiesis. *Mol Genet Metab* 2017; e-pub ahead of print 25 July 2017; doi: 10.1016/j.ymgme.2017.07.008.
- 35 Nguyen AT, Prado MA, Schmidt PJ, Sendamarai AK, Wilson-Grady JT, Min M *et al*. UBE2O remodels the proteome during terminal erythroid differentiation. *Science* 2017; **357**(pii): eaan0218.
- 36 Yanagitani K, Juszkievicz S, Hegde RS. UBE2O is a quality control factor for orphans of multiprotein complexes. *Science* 2017; **357**: 472–475.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>

© The Author(s) 2018