


**CORRIGENDA****Highly effective combination of LSD1 (KDM1A) antagonist and pan-histone deacetylase inhibitor against human AML cells**

W Fiskus, S Sharma, B Shah, BP Portier, SGT Devaraj, K Liu, SP Iyer, D Bearss and KN Bhalla

*Leukemia* (2017) 31, 1658; doi:10.1038/leu.2017.77; published online 21 March 2017**Correction to:** *Leukemia* (2014) 28, 2155–2164; doi:10.1038/leu.2014.119; published online 25 April 2014

neither changes the interpretation of the data nor does it alter the overall conclusions of the study. The corrected Figure 7b appears below:

Following the publication of this article, the authors noted that the loading control ( $\beta$ -actin) for the primary AML cells data presented in Figure 7b was incorrect. The correct western blot for the loading control ( $\beta$ -actin) has been added to Figure 7b. This correction**Figure 7.** Treatment with SP2509 and/or PS significantly enhances PS-mediated loss of viability of CD34+ primary AML cells and improves the survival of mice bearing AML xenografts and primagrafts. **(b)** Primary AML cells were treated with SP2509 and/or PS as indicated for 24 hours. Total cell lysates were prepared and immunoblot analyses were conducted for the expression levels of c-Myc, p21, p27, BIM, and  $\beta$ -actin in the lysates.