See ARTICLE page 646
See ARTICLE page 670

IL-17-producing innate lymphoid cells are restricted to mucosal tissues and are depleted in SIV-infected macaques

H Xu¹, X Wang¹, DX Liu¹, T Moroney-Rasmussen¹, AA Lackner¹ and RS Veazey¹

Innate lymphoid cells (ILCs) are an emerging subset of lymphocytes involved in surveillance against virally infected cells. Here, we show CD3⁺CD8^{high} lymphocytes in macaque blood include major subsets of ILCs including natural killer (NK) cells expressing CD16, NKp46, and NKG2A, but also populations of ILCs in mucosal tissues having different properties. One ILC subset secreted interleukin (IL)-17 (ILC17), but these were restricted to mucosal tissues. Some mucosal ILC17 cells expressed classical NK-cell markers, but little NKG2A or NKG2D. Some ILC17 cells secreted IL-22 and tumor necrosis factor- α , but few produced interferon (IFN)- γ or contained granzyme B. IL-17 production by ILCs was induced by IL-6, transforming growth factor- β , and IL-23. Further, simian immunodeficiency virus (SIV) infection resulted in a significant loss of ILC17 cells, especially in the jejunum, which persisted throughout SIV infection. These findings indicate that ILC17 cells may be involved in innate mucosal immune responses, and their loss may contribute to loss of intestinal mucosal integrity and disease progression in human immunodeficiency virus (HIV)/SIV infection.

Innate lymphoid cells (ILCs) represent a novel family of effector lymphocytes representing the first line of defense against virally infected cells and neoplastic cells. ILCs include natural killer (NK) cells, lymphoid tissue-inducer (LTi) cells, and other subsets that produce interleukin 17 (IL-17), IL-22 (NK-22/ILC22), IL-5, and/or IL-13, which have different functions, but are developmentally related.^{1,2} Thus, ILCs are a heterogeneous subset of CD3-negative lymphocytes with diverse functions, some of which may overlap with T-cell subsets such as Th17, Th22, or Th2 cells.^{1,2} However, the phenotypical characterization of, and distinctions between, NK cells and ILCs, and their relative distribution and function in tissues are not yet clear.

As a major subset of ILCs, NK cells are thought to have an important role in controlling human immunodeficiency virus (HIV) infection and disease progression.^{3–5} Recent studies demonstrate dramatic changes in the phenotype and function of NK cells that occur in HIV/simian immunodeficiency virus (SIV) infection.^{6–8} Moreover, individuals with “protective” KIR/HLA genotypes and/or robust NK-cell responses are overrepresented in HIV controllers.^{5,9–11} This is consistent with studies of SIV-infected macaques in which the magnitude of

NK-cell responses inversely correlates with viremia and disease progression.^{12–14}

Among NK subsets, expression of distinctive surface molecules are highly polymorphic and vary between species.^{15–17} For example, mouse NK cells express NK1.1 and DX5, whereas human NK cells are generally defined by CD56 expression, which is not expressed by mouse NK cells. In humans, CD56^{bright} NK cells possess higher capacity for proliferation and cytokine production, but CD56^{dim} cells display higher cytotoxicity.^{15,18,19} In rhesus macaques (*Macaca mulatta*), prior studies of NK cells and now ILCs have been complicated by different gating strategies and phenotypic definitions. For example, efforts to identify subsets of NK cells in nonhuman primates have included CD3⁺CD16⁺, CD3⁺CD8 $\alpha\alpha$ ⁺, CD3⁺CD20⁺HLA-DR⁺, CD16^{+/–}CD56^{+/–}, etc. However, CD56, often used as a primary marker for human NK cells, is also expressed on macaque monocytes, and may influence data interpretation when gating on CD3⁺ populations.²⁰ Another NK marker, CD16, also known as Fc γ RIII, is important in antibody-dependent cell-mediated cytotoxicity, and is among the first of activating receptors expressed on NK cells. However, detection of CD16

¹Tulane National Primate Research Center, Tulane University School of Medicine, 18703 Three Rivers Road, Covington, Louisiana 70433, USA.
Correspondence: RS Veazey (rveazey@tulane.edu)

Received 2 December 2011; accepted 13 April 2012; published online 6 June 2012. doi: 10.1038/mi.2012.39

with the widely used 3G8 monoclonal antibody clone has been shown to be “masked” by anti-SIV immunoglobulin G immune complexes in SIV-infected rhesus macaques, at least when using whole-blood staining techniques.²¹ Therefore, some prior studies may reflect misleading or incomplete definitions of NK cells in SIV-infected macaques. Thus, more detailed phenotyping of NK cells, and now ILCs, needs to be performed, especially in tissues of SIV-infected rhesus macaques to adequately track and characterize innate immune responses.

NK cells have been described in blood and tissues of various species,^{8,22} yet they are negative for the transcription factor RAR-related orphan γ receptor (ROR γ t).¹ In contrast, the ROR γ t is required for the generation of LT α i cells, which are CD117 (c-Kit)^{high} and CD127 (IL-7R α)^{high}, and other subsets characterized by the production of high amounts of IL-22 including NK-like NKp46⁺ cells in the intestinal lamina propria that are CD117^{low}CD127^{low}.^{23–26} LT α i cells are clustered in cryptopatches between crypts of the intestinal lamina propria, where they, together with stromal cells, could direct the formation of isolated lymphoid follicles by recruitment of dendritic cells and B cells, and promote T-helper cell survival and immunoglobulin A production.^{27,28} In contrast, IL-22 + NKp46⁺ cells are found within intestinal villi.²⁹ In general, NK cells, LT α i, ILC22, and other ILC subsets are all characterized by secretion of proinflammatory and cytolytic mediators, and may have critical roles in lentiviral infection.^{2,30–32} However, little is known about the distribution of IL-17-producing ILC subsets and their responses during SIV infection in rhesus macaques.

To better understand the characteristic of ILCs including NK cells in the peripheral blood and mucosal lymphoid tissues, here we defined ILCs as CD8^{high} lymphocytes, which were lineage negative for B cells (CD20), T cells (CD3), and monocytes (CD14), and examined their anatomical distribution, and changes in tissues after SIV infection. As previously shown for NK cells,^{8,33} the data suggest that ILCs in peripheral blood are distinct from mucosal-derived ILCs, and IL-17-producing ILCs were mostly restricted to mucosal tissues. Further, although some ILC17 expressed classical NK-cell markers, few expressed NKG2A or NKG2D. Finally, there was a significant loss of ILC17 cells in SIV infection, which may contribute to the loss of intestinal epithelial mucosal integrity.

RESULTS

CD3[−]CD8^{high} populations define ILCs subsets in rhesus macaques

To better define the phenotype of macaque ILCs by flow cytometry, we first gated on CD3[−]CD8^{high} lymphocytes (**Figure 1a**), confirmed they were lymphocytes by CD45 expression, showed they lacked CD20 (B cells), CD14 (monocytes), CD11c (dendritic cells), CD163, and CD11b + CD11c^{int} (macrophage/eosinophil) expression, and then examined their expression of typical NK markers on these lineage-negative (Lin[−]) CD3[−]CD8^{high} ILC populations for comparison with CD4⁺ and CD8⁺ T-cell subsets (**Figure 1b** and **c**). Lin[−]CD3[−]CD8^{high} cells were essentially all CD45⁺ lymphocytes, whereas the Lin[−]CD8[−] population was CD45 negative (**Figure 1a**). The CD3[−]CD8^{high} population

was thus distinguished from B cells, monocytes, dendritic cells, macrophage, and eosinophils, and expressed low levels of HLA-DR (**Figure 1b**). Analysis of surface NK markers showed that circulating (blood) CD3[−]CD8^{high} cells expressed CD16 (90.9%), CD56 (11.5%), NKp46 (82.4%), and NKG2A (87.2%), compared with T-cell subsets and/or CD3[−]CD8[−] cells, which either lacked or had much lower expression of these markers (**Figure 2c** and **d**). However, as phenotyping NK cells in rhesus macaques is difficult and as none of these markers alone could definitively distinguish NK cells, these are referred to here as ILCs.

Distribution of NK subsets in macaque tissues

It was previously reported that detection of CD16 on NK cells in whole-blood preparations can be “masked” by non-covalent binding of SIV/immunoglobulin G immune complexes when using the anti-CD16 antibody clone 3G8, which is not observed in fractionated (washed) peripheral blood mononuclear cells.²¹ To identify a better antibody for detecting CD16, we compared four different anti-human CD16 antibody clones, which target different CD16 epitopes, which were reported to cross-react with rhesus macaques, and compared percentages of CD16⁺ ILCs (CD3[−]CD8^{high}) cells of whole-blood and isolated (washed) peripheral blood mononuclear cells (PBMCs) from chronically SIV-infected macaques. Consistent with the previous report, the binding of anti-CD16 antibody was significantly reduced in whole blood when using clones 3G8, NKp15, and VEP13, compared with isolated PBMCs ($P < 0.05$). However, staining with clone DJ130c was not affected, and showed consistent results between whole blood and PBMCs. Moreover, this clone detected even higher levels of CD16 than clone 3G8, even on isolated PBMCs (**Figure 2a**). Thus, in this study, we used this anti-CD16 clone to detect CD16⁺ cells.

In macaques, NKG2A⁺ NK cells are identified as medium-to-large lymphocytes, which display specific functional profiles/properties in SIV infection.⁸ Here, CD16⁺, CD56⁺, or NKG2A⁺ NK or ILC cell subsets were compared in different tissues of normal macaques, and results showed ILC subpopulations were differentially distributed in lymphoid and mucosal tissues. High percentages of CD16⁺ ILCs were detected in peripheral blood, but proportions of CD56⁺ ILCs were markedly higher in other tissues including bone marrow, spleen, liver, and intestine (**Figure 2b**). In contrast, percentages of NKG2A⁺ ILC/NK cells were variable in different tissues but their highest frequency was in the liver followed by peripheral blood (**Figure 2b**). These results indicate ILC/NK subpopulations are differentially distributed in various tissues as heterogeneous lymphocyte populations.

ILCs in blood are distinct from ILCs in mucosal lymphoid tissues

To identify distinct ILCs subpopulations between the peripheral blood and mucosal tissues, we compared the expression of various surface molecules on ILCs. Expression of migration/homing markers including CCR7, CCR6, and CD103 were differentially expressed on ILCs in peripheral and mucosal tissues. Circulating ILCs expressed more CCR7 and CCR6, whereas

Figure 1 Phenotypic characterization of innate lymphoid cells in peripheral blood of normal rhesus macaques. **(a)** Representative gating strategy to define innate lymphoid cells (ILCs) in peripheral blood mononuclear cells (PBMCs). Macaque ILCs were defined as CD3⁺CD8^{high}-gated lymphocytes. **(b)** Representative dot plot depicting the expression of CD20, HLA-DR, CD14, CD11c, CD163, and CD11b on CD3⁺CD8^{high} (R1) as compared with CD3⁺CD8^{low/-} (R3) subsets. **(c)** Representative histogram showing expression of classical natural killer (NK) markers, CD16, CD56, NKp46, NKG2A, and NKG2D, on CD3⁺CD8^{high} (R1), CD3⁺CD8⁺ (R2), CD3⁺CD8⁻/intermediate (R3), and CD3⁺CD4⁺ (R4) subsets from peripheral blood. **(d)** Expression of classical NK markers on CD3⁺CD8^{high} (R1), CD3⁺CD8⁺ (R2), CD3⁺CD8⁻/intermediate (R3), and CD3⁺CD4⁺ (R4) subsets from peripheral blood ($n=8$). Examples are representative of eight naive animals. * $P<0.001$; ** $P<0.05$, compared between CD3⁺CD8^{high} and other three subsets, respectively.

jejunum-derived ILCs expressed higher levels of CD103, which promotes lymphocyte migration to mucosal epithelium. Further, the expression of CXCR5, CD62L, CD27, and CD90 were also differentially expressed on ILCs in tissues (**Figure 3a**). In rhesus macaques, other molecules such as CD27 and CD62L were also differentially expressed on ILCs between peripheral blood and mucosal gut tissues. Mucosal ILCs had higher percentages of CD27^{low} and CD62L^{low} populations than peripheral blood (**Figure 3b**). Nonetheless, despite their common CD3⁺CD8^{high} phenotype, ILCs in different tissues appear to be phenotypically diverse.

IL-17-producing ILCs are predominantly restricted to mucosal tissues

To further investigate and compare ILCs in blood and intestinal tissues, total lymphocyte preparations were stimulated with phorbol 12-myristate-13-acetate (PMA) plus ionomycin. Among the IL-17-producing lymphocytes, there was a substantial population of CD3⁺ lymphocytes in the jejunum as shown in **Figure 4a**. Further analyzing these IL-17⁺CD3⁺ populations in tissues showed they were predominantly CD3⁺CD8^{high} ILCs, and were essentially restricted to mucosal tissues including the jejunum, colon, tonsil, and duodenum, and absent

Figure 2 Comparison of anti-CD16 antibody clones for cross-reactivity with rhesus macaques, and distribution of natural killer (NK) cells in lymphoid tissues. (a) Cross-reactivity and comparison of whole-blood (WB) and washed peripheral blood mononuclear cell (PBMC) from the same animals using different anti-CD16 antibody clones in chronically simian immunodeficiency virus-infected rhesus macaques ($n=8$). Note only the DJ130c clone does not demonstrate differential staining between washed PBMC and WB. (b) Tissue distribution of classical NK-cell subsets (CD16+, CD56+, and NKG2A+) in naïve rhesus macaques ($n=5$). * $P<0.05$ between using 3G8 and DJ130c clones in PBMC samples (a) or blood compared with other tissues (b); ** $P<0.01$ between using 3G8 and DJ130c clones in WB; # $P<0.01$ compared with peripheral blood; ## $P<0.05$ between liver and other tissues (except blood), respectively.

in blood or spleen. Similarly, the vast majority of the Th17 cells (CD3+CD4+IL-17+) and CD3+CD8+ T cells-producing IL-17 were detected in intestinal tissues, but in contrast to the ILCs, there were substantial populations of these in blood (PBMC) as well (Figure 4b). Moreover, and similar to Th17 and CD8+ T cells, much higher percentages of IL-17-producing ILC cells (defined here as ILC17 cells) were found in the small intestine compared with the colon or any other tissue. A similar distribution of Th17 cells has been previously reported in mice,³⁴ but to our knowledge these have not been examined in mucosal tissues of humans. As the small intestine is a major immune effector site, these data suggest that ILC cells and other IL-17-producing cells are activated and effector cells involved in maintaining mucosal immunity.

To further identify the distribution of ILC17 in intestinal tissues in macaques, jejunum sections were analyzed by multi-fluorescent immunohistochemistry to determine their location and distribution *in situ*. By simultaneously labeling tissues with CD3, CD8, and IL-17, we could show ILC17 cells (CD3negCD8+IL-17+) were present in the jejunum lamina propria, as well as Tc17 (CD3+CD8+IL-17+) and Th17 (CD3+CD8negIL-17+) cells in normal, uninfected animals (Figure 5a and b). After SIV infection, IL-17+ cells were gradually depleted with disease progression, and were practically undetectable in AIDS macaques (data not shown), consistent

with previous reports documenting loss of Th17 and Tc17 cells in chronic SIV infection.^{35,36}

ILC17 cells differentially express classical NK markers and produce an array of cytokines

As we found that ILC17 cells were predominantly distributed in mucosal tissues, we further phenotyped these for expression of classical NK-cell markers. As shown in Figure 6a, ILC17 cells in the jejunum variably expressed NK molecules including CD16, CD56, Nkp44, and Nkp46 when gating on CD3+CD8^{high} populations. These proportions appear very different than those reported for IL-22-producing NK cells, most of which have been shown to co-express Nkp46 (in mice) or Nkp44 (in humans). The distribution of ILC17 cells was similar between intestinal tissues. In contrast, little to no production of IL-17 was detected from NKG2A+ or NKG2D+ ILC cells in the intestines or from any peripheral blood-derived ILCs (Figure 6a).

As NK cells are characterized by their cytotoxicity and/or regulation of inflammatory responses, we examined cytokine secretion of ILC17 cells in response to mitogens or cytokines. Following mitogen stimulation, ILC17 cells also secreted IL-22 and tumor necrosis factor (TNF)- α . In fact, almost all ILC17 cells co-expressed TNF- α - and ~30% co-expressed IL-22 suggesting these were NK-22 cells now believed to have a major role in mucosal defense.^{32,37,38} However, little interferon (IFN)- γ and granzyme B was secreted by ILC17 cells (Figure 6b). Combined, these data suggest that ILC17 cells might be involved in regulating innate inflammatory responses in gut-associated lymphoid tissues, rather than having cytotoxic properties.

Although PMA and ionomycin stimulation resulted in IL-17 production, it was not clear if similar results could be elicited by stimulation with more physiologically relevant cytokines. Thus, we examined the production of IL-17 from jejunum lamina propria-derived ILC after stimulation with IL-1 β (20 ng ml⁻¹), IL-2 (20 U ml⁻¹), IL-6 (20 ng ml⁻¹), IL-15 (20 ng ml⁻¹), transforming growth factor (TGF)- β (20 ng ml⁻¹), IL-23 (50 ng ml⁻¹), and lipopolysaccharide (LPS; 10 ng ml⁻¹). To some extent, IL-23, TGF- β , and IL-6 all induced IL-17 secretion from ILC17 cells *in vitro* (Figure 6c). However, other cytokines or stimuli such as IL-2, IL-15, IL-1 β , or LPS had little to no effect on IL-17 production from ILC17 cells, compared with PMA/IO stimulation (Figure 6c). Combined, these data demonstrate that ILC17 cells are a specialized subset localized within mucosal tissues.

Dynamics of CD3+CD8^{high} ILCs in peripheral and mucosal intestinal tissues after SIV infection

NK cells are reported to have a crucial role in the early defense against many viral infections including HIV/SIV and herpes-viruses.^{33,39,40} Here, we monitored changes in distribution and proliferation of ILCs (CD3+CD8^{high}) after SIV infection in rhesus macaques. In three animals followed prospectively in SIV infection, both absolute numbers and percentages of ILCs in blood were reduced in acute infection and then began to increase after 14 dpi (Figure 7a and b). Consistently, when additional animals were examined, circulating ILCs were significantly reduced 7–14 days post SIV infection (compared

Figure 3 Distinct expression of surface molecules on innate lymphoid cells between mucosal tissues and peripheral blood. **(a)** Representative histogram displaying different expression of molecules on innate lymphoid cells (ILCs) from peripheral blood mononuclear cells (PBMCs) and jejunum lamina propria. **(b)** Statistical comparison of expression between ILCs from PBMCs and jejunum lamina propria in naïve rhesus macaques ($n=12$). * $P<0.05$.

with healthy macaques, $P<0.05$), but gradually increased and were sometimes even higher than baseline in chronic infection (Figure 7c). On average, SIV infection resulted in significant increases in circulating ILCs at 14–21 dpi (Figure 7d). Owing to the disparate phenotypes of ILCs in peripheral blood and mucosal tissues (Figure 2–4), we also examined mucosal ILCs after SIV infection. To further identify possible subpopulations of ILCs, we assessed the expression of various NK-cell-associated molecules including CD16, CD56, NKG2A, NKp30, and NKp46 on total ILC populations. Circulating CD16⁺, CD56⁺, or NKG2A⁺ ILCs showed no significant changes during SIV infection (Figure 7e–g), and displayed similar proliferation profiles (Ki-67⁺) as total ILCs in blood (Figure 7d and h). However, percentages of NKp30⁺ and NKp46⁺ ILCs were significantly reduced post SIV infection (Figure 7i), which was not consistent with the dynamics of total circulating ILCs. Thus, selective reduction in NKp30⁺ and NKp46⁺ ILC subpopulations occurs in acute SIV infection.

In contrast to blood, percentages of intestinal ILCs increased in early SIV infection, peaked at day 21, and then gradually decreased. In chronic infection, percentages of ILCs in gut-associated

lymphoid tissues were significantly reduced compared with normal macaques (Figure 7j). Further analysis indicated that intestinal CD16⁺, CD56⁺, or NKG2A⁺ ILCs increased in response to acute SIV infection, but NKp46⁺ ILCs significantly decreased after infection. In chronic infection, losses of CD56⁺ and NKp46⁺ mucosal ILCs were detected, compared with milder losses of NKG2A⁺ cells and maintenance of CD16⁺ populations (Figure 7k). Similar to circulating ILCs, proliferation rates of total intestinal ILCs and various subpopulations were also markedly increased in acute SIV infection, yet returned to baseline levels in chronically SIV-infected macaques (Figure 7l). These data demonstrate that intestinal-derived ILCs are indeed distinct from peripheral blood ILCs, and various subpopulations display different dynamics in response to SIV infection. As larger changes in these populations occur in the intestine compared with blood, we hypothesize this may correlate with the higher levels of CD4⁺ T-cell infection and viral replication in intestinal tissues in early SIV infection.⁴¹ If ILCs are involved in effector and regulatory functions, dysfunction and/or deletion of these cells may be associated with failure of local viral control during SIV infection.

Figure 4 Innate lymphoid cells that secrete interleukin (IL)-17 are restricted to mucosal tissues in rhesus macaques. **(a)** Jejunum dot plots gated on IL-17-secreting lymphocytes (left) contain both CD3⁺ and CD3⁻ subsets in jejunum after phorbol 12-myristate-13-acetate/ionomycin stimulation. **(b)** Comparison of IL-17-producing CD4⁺ (Th17), CD8⁺ T cells, and CD3⁻CD8^{high} (innate lymphoid cells (ILCs)) in various lymphoid tissues, including peripheral blood, jejunum, colon, spleen, tonsil, and duodenum. Noted that IL-17-secreting ILCs are restricted to mucosal tissues, with little to no expression of IL-17 from peripheral blood mononuclear cell (PBMC) or spleen-derived ILCs.

Loss of mucosal ILC17 cells correlates with disease progression in SIV-infected macaques

As ILC17 cells appear to be restricted to mucosal tissues, we further investigated the effects of SIV infection on ILC17 cells in intestinal tissues of rhesus macaques. The baseline percentage of ILC17 cells was $\sim 13.27 \pm 2.386\%$ of jejunum lamina propria lymphocytes that was higher than in colon ($5.639 \pm 1.124\%$) in SIV naïve/negative animals. However, ILC17 cells in the jejunum were significantly reduced at all stages of SIV infection, and remained at significantly lower levels in animals with symptomatic AIDS ($1.654 \pm 0.603\%$, $P < 0.05$). Similarly, percentages of ILC17 cell in the colon also showed decreases in acute infection, but there was a slight (but not significant) increase over baseline in chronic infection. However, ILC17 cells were also significantly reduced in the colon in animals with AIDS as compared with chronically SIV-infected animals (**Figure 8a and b**).

Owing to the marked loss of ILC17 cells in the jejunum after SIV infection, we asked whether levels of ILC17 cells in this tissue correlated with intestinal CD4⁺ T-cell or Th17 cell loss in the jejunum. The data showed that ILC17 cell loss positively

correlated with loss of both total CD4⁺ T cells ($R^2 = 0.5906$, $P = 0.0002$) and Th17 cells ($R^2 = 0.5067$, $P < 0.0001$) (**Figure 8c and d**). As HIV/SIV infection results in massive loss of intestinal CD4⁺ T cells,⁴² including loss of Th17 cells,^{43,44} these findings suggest the loss of ILC17 cells may also be involved with the loss of intestinal integrity, and disease progression in AIDS.

DISCUSSION

ILCs are a heterogeneous population of cells thought to have an important role in innate immunity and lentivirus infections. However, more recent evidence suggests they are also involved in regulating adaptive immune responses. Here, we define ILCs as CD3⁻CD8^{high} lymphocytes that lack CD3, CD20, and CD14, and show they are differentially distributed in blood, lymphoid and non-lymphoid tissues, and display different expression patterns of migration or “memory”-associated molecules in rhesus macaques. Further, we show large proportions of these cells that produce IL-17 upon mitogen stimulation are restricted to mucosal effector sites, and lack features of classical NK-cell properties such as Granzyme B. In normal animals, ILCs may

Figure 5 Detection of innate lymphoid cell (ILC)17 in jejunum tissues of normal rhesus macaques by confocal microscopy. **(a)** Jejunum from a normal macaque showing CD3+ (red), interleukin (IL)-17 (green), and CD8+ (blue) cells. The white arrows show ILC17 cells (CD3-CD8+IL-17+); the yellow arrows demonstrate Th17 cells (CD3+CD8-IL-17+); and the blue arrows show Tc17 cells (CD3+CD8+IL-17-). Scale bar = 10 μ m. **(b)** Relative percentages of Th17, Tc17, and ILC17 cells gated through total IL-17+ lymphocytes in the jejunum of normal animals as assessed by flow cytometry.

be a major source of IL-17, which is known to contribute to the maintenance of intestinal epithelial integrity.⁴⁵ Loss of epithelial integrity is now believed to occur after SIV/HIV infection, resulting in translocation of bacteria and other antigens from the intestinal lumen to the systemic circulation, driving systemic immune activation and SIV/HIV replication.^{46,47} Therefore, the loss of IL-17-producing cells may be responsible for the loss of mucosal epithelial integrity, driving SIV/HIV replication and persistence.

In addition, mucosal ILC17 have variable expression patterns of NK-cell-associated molecules such as CD16, CD56, Nkp44, and Nkp46, but little to no NKG2A and NKG2D expression. However, mucosal ILC17 cells also appear to encompass all of the intestinal NK-22 cells. Intestinal mucosal ILC17 cells were significantly reduced in the jejunum post SIV infection, and their loss directly correlated with intestinal CD4+ T-cell and Th17 cell loss. As loss of Th17 cells in the intestine has been shown to correlate with disease progression,^{43,48} these findings suggest that the loss of ILC17 cells may also contribute to disease progression and the pathogenesis of AIDS.

NK cells as a subset of ILCs may phenotypically differ between humans, mice, and macaques, yet prior definitions and gating strategies preclude direct comparisons. Although CD56 is regarded as a major marker for NK cells in humans, its expression on monocytes in rhesus macaques²⁰ may interfere with gating strategies simply based on CD3-negative cells, which could include small monocytes in the lymphocyte gate. However, CD8 is a member of the immunoglobulin superfamily found on the majority of thymocytes, T cells, and NK cells (which almost exclusively express CD8a homodimers), and there is little to no expression of CD8 on monocytes. Therefore, CD3⁻CD8^{high} populations exclude T cells, monocytes, and other cells, and **Figure 1** (R1) confirms that circulating CD3⁻CD8^{high} lymphocytes express low to undetectable levels of CD20, CD14, CD11c, CD163, or CD11b+CD11c^{int}, but highly express classical NK-cell markers such as CD16, CD56, Nkp46, NKG2A, and/or NKG2D. Moreover, detection of CD16 using the anti-CD16 antibody clone DJ130c showed highly consistent results and even better staining than the 3G8 clone, which is known to be masked by anti-SIV antibody complexes (**Figure 2a**). Thus, the anti-CD16 antibody clone DJ130c appears more suitable for CD16 detection in macaques than antibodies previously used for SIV studies, which could underestimate levels of CD16+ cells in whole-blood staining preparations of chronically infected macaques.

ILCs display different distributions and staining characteristics in tissues, especially between peripheral blood and mucosal tissues. Levels of CD56+ ILCs in tissues are significantly higher than in blood (**Figure 2b**) and ILCs in gut-associated lymphoid tissues that are distinct from those in blood in expression of migration or other molecules including CXCR5, CCR7 and CD103, CD27 and CD62L (**Figure 3**). In macaques, mucosal ILCs had much higher levels of CD62L^{low} and CD27^{low/-} subsets than in blood. These data suggest possible diverse functions and immunological experience of the ILC found in blood and those in tissues.

Previous studies found Th17 cells and Nkp44+ NK-22 cells could be a major source of IL-17 in tissues.^{32,33} Within the CD3⁻CD8^{high} subsets, we found ILCs also produced IL-17 but these were restricted to mucosal tissues such as the jejunum, duodenum, and tonsil, compared with blood in which IL-17 was secreted only by CD4+ or CD8+ T cells after PMA/ionomycin stimulation (**Figure 4**). It is well known that Th17 cells are abundant in intestinal tissues; however, we found more Tc17 cells in the jejunum compared with Th17 cells (**Figure 5b**). We also found more ILC17 cells in the jejunum than the colon, which was not surprising, as the jejunum is purely an effector lymphoid tissue, whereas the colon contains abundant organized lymphoid tissues that consist largely of resting, naïve cells. Further, intestinal IL-17-producing cells partially expressed CD16, CD56, Nkp44, and Nkp46 yet little to no expression of NKG2A and NKG2D (**Figure 6a**). Moreover, ILC17 cells predominantly produced TNF- α , but little secretion of cytolytic IFN- γ or granzyme B. In addition, all IL-22+ cells were contained in the IL-17+ subset, yet only about 30% of the IL-17 cells were IL-22+, suggesting ILC17 cells represent a distinct

Figure 6 Phenotyping interleukin (IL)-17-producing innate lymphoid cells (ILCs) for classical NK-cell markers and cytokine secretion in peripheral blood and mucosal tissues of normal rhesus macaques. **(a)** Expression of classical NK-cell markers on ILC17 cells in the peripheral blood, jejunum, and colon. Note there was little to no production of IL-17 from NKG2A⁺ or NKG2D⁺ ILCs. **(b)** Cytokine secretion of ILC17 cells isolated from jejunum after PMA/ionomycin stimulation. Note ILC17 cells also secrete pro-inflammatory (tumor necrosis factor (TNF)- α) and innate (IL-22) cytokines, but express little to no IFN- γ or Granzyme B. **(c)** Response of jejunum ILC17 cells to various cytokine and lipopolysaccharide stimulation.

subset from IL-22-producing NK cells in macaques (**Figure 6b**). Thus, ILC17 cells may be considered as a heterogeneous family of ILCs including NK cells, lymphoid tissue inducing cells (LTis), and other cells such as NK-22, which share a common lineage.^{1,49} Most ILCs including ILC22 and lymphoid tissue inducer cells (LTis) are distinguished by high levels of the transcription factor ROR γ t.^{30,50,51} In rhesus macaques, we tested several anti-human ROR γ t antibodies, but none reliably cross-reacted with macaques. However, IL-17-producing ILCs in macaques did produce IL-22, and expressed NKp46 and/or NKp44 (**Figure 6a**). ILC17 cells also appeared to be lacking cytotoxic functions, as they did not express killer inhibitory receptors such as NKG2A, and produced little IFN- γ (**Figure 5a and b**). Further examination showed mucosal ILC17 could (to some extent) be regulated by TGF- β , IL-6, and IL-23, but not by IL-2, IL-15, or LPS (**Figure 6c**). Perhaps, robust IL-17 production from ILC17 cells may require costimulatory signals such as toll-like receptor ligands, such as the TLR2 ligand for NK-22.⁵² Nonetheless, these

data indicate ILC17 cells represent a specific subset restricted to mucosal tissues, and suggest they also have a role in maintaining mucosal integrity and innate immunity.

In blood, percentages of circulating ILCs were reduced in early SIV infection (10–14 days) and then dramatically increased, accompanied by higher proliferative capacity (day 14–21), but no significant differences were detected between naïve and chronically SIV-infected animals (**Figure 7a–d**). In contrast to blood, percentages of intestinal ILCs increased and peaked at 21 days post SIV infection, and maintained high proliferation rates through 28 days of infection (**Figure 7j, e and l**), but were gradually “depleted” with disease progression. Within the ILCs, the NKp46⁺ population decreased after SIV infection in both peripheral blood and intestinal mucosal tissues (**Figure 7i and k**), but other subsets expressing classic NK cells markers such as CD16, CD56 (except in mucosal tissues in chronic infection), and NKG2A did not significantly decrease post viral infection (**Figure 7e–j and k**). These results indicate that changes

Figure 7 Effects of simian immunodeficiency virus (SIV) infection on blood and jejunum-derived innate lymphoid cells (CD3⁺CD8^{high}) or its subpopulations in rhesus macaques. (**a** and **b**) Prospective analysis of changes in percentage (**a**) and absolute numbers (**b**) of total CD3⁺CD8^{high} innate lymphoid cells (ILCs) in peripheral blood after SIV infection. (**c**, **e**–**g**, and **i**–**k**) Dynamic of ILCs and their subpopulation in blood and jejunum ILCs after SIV infection (naïve, $n=55$; 7 dpi, $n=30$; 14 dpi, $n=23$; 21 dpi, $n=26$; 28 dpi, $n=26$; 35 dpi, $n=18$; chronic, $n=23$). (**d**, **h**, and **l**) Proliferation (Ki-67⁺) of circulating or intestinal ILCs after SIV infection (naïve, $n=24$; 7 dpi, $n=5$; 14 dpi, $n=5$; 21 dpi, $n=4$; 28 dpi, $n=3$; 42 dpi, $n=4$; chronic, $n=14$). * $P<0.05$, compared with uninfected normal animals.

in circulating ILCs differ from those in mucosal tissues. ILCs may also have an important role in innate immunity in mucosal tissues in early viral infection.

SIV infection resulted in reductions of ILC17 cells, especially in the jejunum. In macaques with AIDS, very few ILC17 cells were detectable (Figure 8a and b). The loss of ILC17 cells directly correlated with the loss of intestinal CD4⁺ T cells and Th17 cells (Figure 8c and d), which has been correlated with breakdown of mucosal integrity, resulting in microbial translocation and chronic immune activation.^{36,46} IL-17, in combination with IL-22 (also derived from most IL-17-producing cells), is important for protection against mucosal bacterial infections and for maintenance of the mucosal barrier by promoting intestinal epithelial integrity.^{35,36,48,53–55} In the intestinal mucosa, ILCs are in close proximity to the epithelial barrier, and may even be intraepithelial or subepithelial.³² Although no direct evidence currently shows the loss of intestinal ILC17 cells correlates with breakdown of intestinal mucosal integrity in SIV-infected rhesus macaques, it has been shown that loss of other IL-17-producing cells (CD4⁺ Th17 cells) correlates with loss of epithelial integrity.⁵⁶ Therefore, general preservation of IL-

17-producing cells during HIV infection may prevent the immune activation associated with loss of epithelial integrity and microbial translocation that leads to higher levels of SIV/HIV replication.^{56–58}

METHODS

Animals and virus

A total of 71 adult Indian rhesus macaques (*M. mulatta*), which were negative for SIV, type D retrovirus, and STLV-1 infection, were utilized to examine ILCs in tissues and blood. Another 146 macaques were examined for ILC subsets in peripheral blood. All animals were housed at the Tulane National Primate Research Center in accordance with the Association for Assessment and Accreditation of Laboratory Animal Care International standards. All studies were reviewed and approved by the Tulane University Institutional Animal Care and Use Committee. Of the 71 animals, 20 animals were uninfected controls, and others were infected with SIVmac251 and examined in acute ($n=23$) or chronic infection with either no overt signs of disease (chronic asymptomatic, $n=24$) or with symptoms of AIDS ($n=4$). Blood from three animals was prospectively monitored at different time points post SIV infection. To examine cells from tissues such as intestine, spleen, etc., macaques were euthanized for tissue collection as naïve, uninfected controls, or in acute (7–21 days) or chronic infection (SIV infection more than 3 months).

Figure 8 Reduction of intestinal mucosal innate lymphoid cell (ILC)17 cells after simian immunodeficiency virus (SIV) infection of rhesus macaques. (a) Dot plot demonstrating ILC17 cells after gating on CD3⁺CD8^{high} subsets from the jejunum and colon from a representative normal and AIDS macaque. (b) Loss of intestinal ILCs occurs after SIV infection (normal, $n=10$; acute, $n=6$; chronic, 12; AIDS, $n=4$). * $P<0.05$. (c) Correlation between percentages of ILC17 cells and CD4⁺ T cells or (d) Th17 cells in jejunum during SIV infection (c, $n=18$; d, $n=28$).

Cell isolation and processing

Mononuclear cells from peripheral blood, spleen, and intestinal tissues were isolated and processed as previously described.⁵⁹ Briefly, total peripheral blood mononuclear cells were isolated from EDTA-treated venous blood by density gradient centrifuge with Lymphocyte Separation Media (MP Biomedicals, LLC, Santa Ana, CA) as per manufacturers instruction. Tissues were collected from the jejunum and colon within minutes of euthanasia and processed immediately for cell suspensions using enzymatic digestion as previously described.⁶⁰

Phenotyping

Flow cytometry for surface and intracellular staining was performed using standard protocols.⁶¹ Cells were stained with: CD3 (SP34), Ki67 (B56), CD8 α (3B5, Caltag Laboratories, Burlingame, CA), HLA-DR (Immu-357, Beckman Coulter, Brea, CA), TNF- α (MAB11), IFN- γ (B27), CCR6 (11A9), CCR7 (150503, R&D Systems, Minneapolis, MN), CD20 (L27), CD14 (M5E2), CD11c (S-HCL-3), CD11b (ICRF44), CD163 (GH/61), CD45 (TU116), CD56 (NCAM16.2), CD16 (3G8 and NKp15, BD Biosciences, San Diego, CA; VEP13, Miltenyi Biotec, Auburn, CA; DJ130c, Dako, Carpinteria, CA), NKp30 (Z25, Beckman Coulter), NKp46 (BAB281, Beckman Coulter), NKG2A (Z199, Beckman Coulter), NKG2D (1D11, Biolegend, San Diego, CA), NKp44 (2.9, Miltenyi), CXCR5 (NIH reagents), CD62L (SK11), CD103 (2G5, Beckman Coulter), CD27 (M-T271), CD90 (5E10), Granzyme B (GB11), IL-17 (eBio64CAP17, eBioscience, San Diego, CA), and IL-22 (BG/IL22, Biolegend; 142928; R&D), LIVE/DEAD Fixable Aqua Dead Cell Stain Kit (Invitrogen, Grand Island, NY). All antibodies and reagents were purchased from BD Biosciences Pharmingen (San Diego, CA) unless otherwise noted. Samples were resuspended in BD Stabilizing Fixative (BD Biosciences) and acquired on a FACS Calibur or FACS Aria flow cytometer (Becton Dickinson, San Jose, CA). Data were analyzed with Flowjo software (Tree star, Ashland, OR).

Ex vivo tissue culture and multicolor confocal microscopy

Fresh jejunum tissues were obtained from rhesus macaques within 30 min of necropsy and explants prepared and stimulated for detection of IL-17⁺ cell subsets. Tissues were cut into 1-cm² sections and cultured in complete RPMI medium (10% heat inactivated fetal calf serum, l-glutamine, penicillin, and streptomycin; Invitrogen) either alone (unstimulated

media) or with 100 ng ml⁻¹ PMA plus 0.5 mg ml⁻¹ calcium ionophore (stimulation medium) for 4 h in the presence of 2 μ M monensin (Sigma, St Louis, MO) to block protein transport and release. Tissues were then processed and stained as previously described.⁵⁹ In brief, tissues were embedded and snap frozen in optimum cold temperature compound and 7- μ m frozen sections were stained using unconjugated primary antibodies (CD3, CD8, or IL-17) followed by appropriate secondary antibodies conjugated to Alexa 488 (green), Alexa 568 (red), or Alexa 633 (blue) (Molecular Probes, Eugene, OR). Confocal microscopy was performed using a Leica TCS SP2 confocal microscope equipped with three lasers (Leica Microsystems, Exton, PA). Individual optical slices representing 0.2 μ m and 32 to 62 optical slices were collected at 512 $\times$ 512 pixel resolution. NIH Image (version 1.62, Bethesda, MD) and Adobe Photoshop (version 7.0, San Jose, CA) were used to assign colors to the channels collected: HNPP/Fast Red (Roche, Indianapolis, IN), which fluoresces when exposed to a 568-nm wavelength laser, appears red; Alexa 488 (Molecular Probes) appears green; and Alexa 633 (Molecular Probes) appears blue (Figure 5).

Cell stimulation for detection of cytokines

Lymphocytes (10^6) from jejunum, colon, and tonsil were stimulated *in vitro* with 0.1 μ M PMA and 0.5 μ g ml⁻¹ ionomycin (Sigma-Aldrich, St Louis, MO), or 20 U ml⁻¹ IL-2, 10 ng ml⁻¹ LPS (Sigma), 20 ng ml⁻¹ IL-1 β , IL-6, IL-7, and IL-15 (R&D), TGF- β or 50 ng ml⁻¹ IL-23 (Biolegend) for 2 hours at 37°C. Cells were cultured for an additional 4 hours in the presence of 5 μ g ml⁻¹ Brefeldin A (Sigma-Aldrich), then stained for cell surface markers and Annexin V, fixed in 2% paraformaldehyde, permeabilized in Cytofix/Cytoperm solution (BD Biosciences), and intracellularly co-stained with fluorochrome-labelled antibodies for the cytokines, and acquired with a FACS Aria cytometer (Becton Dickinson). Data were analyzed with Flowjo software (Tree star).

Statistics

Graphical presentation and statistical analysis of the data were performed using GraphPad Prism 4.0. (GraphPad Software, San Diego, CA). Comparisons between groups were analyzed by a one-way analysis of variance and a nonparametric Mann-Whitney *T*-test. *P*-values <0.05 were considered statistically significant. Correlations between samples were calculated and expressed using the Spearman's coefficient of correlation.

ACKNOWLEDGMENTS

We thank Julie Bruhn and Calvin Lanclos for flow cytometry support and Megan Gardner, Kelsi Rasmussen, Megan Watkins, and Maury Duplantier for technical support. This work was supported by NIH grants AI49080, AI084793, RR000164, and a Faculty Enhancement Grant from Tulane University. The authors declare no conflict of interests.

DISCLOSURE

The authors declare no conflicts of interest.

© 2012 Society for Mucosal Immunology

REFERENCES

- Spits, H. & Di Santo, J.P. The expanding family of innate lymphoid cells: regulators and effectors of immunity and tissue remodeling. *Nat. Immunol.* **12**, 21–27 (2011).
- Crellin, N.K., Trifari, S., Kaplan, C.D., Cupedo, T. & Spits, H. Human NKp44+IL-22+ cells and LTI-like cells constitute a stable RORC⁺ lineage distinct from conventional natural killer cells. *J. Exp. Med.* **207**, 281–290 (2010).
- Berger, C.T. & Alter, G. Natural killer cells in spontaneous control of HIV infection. *Curr. Opin. HIV AIDS* **6**, 208–213 (2011).
- French, A.R. & Yokoyama, W.M. Natural killer cells and viral infections. *Curr. Opin. Immunol.* **15**, 45–51 (2003).
- Alter, G. *et al.* Differential natural killer cell-mediated inhibition of HIV-1 replication based on distinct KIR/HLA subtypes. *J. Exp. Med.* **204**, 3027–3036 (2007).

6. De Maria, A. *et al.* The impaired NK cell cytolytic function in viremic HIV-1 infection is associated with a reduced surface expression of natural cytotoxicity receptors (Nkp46, Nkp30 and Nkp44). *Eur. J. Immunol.* **33**, 2410–2418 (2003).
7. Mavilio, D. *et al.* Characterization of CD56-/CD16+ natural killer (NK) cells: a highly dysfunctional NK subset expanded in HIV-infected viremic individuals. *Proc. Natl Acad. Sci. USA* **102**, 2886–2891 (2005).
8. Reeves, R.K., Gillis, J., Wong, F.E., Yu, Y., Connole, M. & Johnson, R.P. CD16- natural killer cells: enrichment in mucosal and secondary lymphoid tissues and altered function during chronic SIV infection. *Blood* **115**, 4439–4446 (2010).
9. Vieillard, V., Fausther-Bovendo, H., Samri, A. & Debre, P. Specific phenotypic and functional features of natural killer cells from HIV-infected long-term nonprogressors and HIV controllers. *J. Acquir. Immune Defic. Syndr.* **53**, 564–573 (2010).
10. O'Connor, G.M., Holmes, A., Mulcahy, F. & Gardiner, C.M. Natural killer cells from long-term non-progressor HIV patients are characterized by altered phenotype and function. *Clin. Immunol.* **124**, 277–283 (2007).
11. Martin, M.P. *et al.* Epistatic interaction between KIR3DS1 and HLA-B delays the progression to AIDS. *Nat. Genet.* **31**, 429–434 (2002).
12. Shieh, T.M., Carter, D.L., Blosser, R.L., Mankowski, J.L., Zink, M.C. & Clements, J.E. Functional analyses of natural killer cells in macaques infected with neurovirulent simian immunodeficiency virus. *J. Neurovirol.* **7**, 11–24 (2001).
13. Pereira, L.E., Johnson, R.P. & Ansari, A.A. Sooty mangabeys and rhesus macaques exhibit significant divergent natural killer cell responses during both acute and chronic phases of SIV infection. *Cell Immunol.* **254**, 10–19 (2008).
14. Bostik, P. *et al.* Decreased NK cell frequency and function is associated with increased risk of KIR3DL allele polymorphism in simian immunodeficiency virus-infected rhesus macaques with high viral loads. *J. Immunol.* **182**, 3638–3649 (2009).
15. Freud, A.G. & Caligiuri, M.A. Human natural killer cell development. *Immunol. Rev.* **214**, 56–72 (2006).
16. Carlyle, J.R. *et al.* Evolution of the Ly49 and Nkrp1 recognition systems. *Semin. Immunol.* **20**, 321–330 (2008).
17. Kulkarni, S., Martin, M.P. & Carrington, M. The Yin and Yang of HLA and KIR in human disease. *Semin. Immunol.* **20**, 343–352 (2008).
18. Yokoyama, W.M., Kim, S. & French, A.R. The dynamic life of natural killer cells. *Annu. Rev. Immunol.* **22**, 405–429 (2004).
19. Cooper, M.A., Fehniger, T.A. & Caligiuri, M.A. The biology of human natural killer-cell subsets. *Trends Immunol.* **22**, 633–640 (2001).
20. Carter, D.L. *et al.* CD56 identifies monocytes and not natural killer cells in rhesus macaques. *Cytometry* **37**, 41–50 (1999).
21. Wei, Q., Stallworth, J.W., Vance, P.J., Hoxie, J.A. & Fultz, P.N. Simian immunodeficiency virus (SIV)/immunoglobulin G immune complexes in SIV-infected macaques block detection of CD16 but not cytolytic activity of natural killer cells. *Clin. Vaccine Immunol.* **13**, 768–778 (2006).
22. Chinen, H. *et al.* Lamina propria c-kit+ immune precursors reside in human adult intestine and differentiate into natural killer cells. *Gastroenterology* **133**, 559–573 (2007).
23. Sawa, S. *et al.* Lineage relationship analysis of RORgammat+ innate lymphoid cells. *Science* **330**, 665–669 (2010).
24. Takatori, H. *et al.* Lymphoid tissue inducer-like cells are an innate source of IL-17 and IL-22. *J. Exp. Med.* **206**, 35–41 (2009).
25. Sanos, S.L. *et al.* RORgammat and commensal microflora are required for the differentiation of mucosal interleukin 22-producing Nkp46+ cells. *Nat. Immunol.* **10**, 83–91 (2009).
26. Luci, C. *et al.* Influence of the transcription factor RORgammat on the development of Nkp46+ cell populations in gut and skin. *Nat. Immunol.* **10**, 75–82 (2009).
27. Tsuji, M. *et al.* Requirement for lymphoid tissue-inducer cells in isolated follicle formation and T cell-independent immunoglobulin A generation in the gut. *Immunity* **29**, 261–271 (2008).
28. Bouskra, D. *et al.* Lymphoid tissue genesis induced by commensals through NOD1 regulates intestinal homeostasis. *Nature* **456**, 507–510 (2008).
29. Satoh-Takayama, N. *et al.* Microbial flora drives interleukin 22 production in intestinal Nkp46+ cells that provide innate mucosal immune defense. *Immunity* **29**, 958–970 (2008).
30. Biron, C.A., Nguyen, K.B., Pien, G.C., Cousens, L.P. & Salazar-Mather, T.P. Natural killer cells in antiviral defense: function and regulation by innate cytokines. *Annu. Rev. Immunol.* **17**, 189–220 (1999).
31. Chang, Y.J. *et al.* Innate lymphoid cells mediate influenza-induced airway hyper-reactivity independently of adaptive immunity. *Nat. Immunol.* **12**, 631–638 (2011).
32. Cella, M. *et al.* A human natural killer cell subset provides an innate source of IL-22 for mucosal immunity. *Nature* **457**, 722–725 (2009).
33. Reeves, R.K. *et al.* Gut inflammation and indoleamine deoxygenase inhibit IL-17 production and promote cytotoxic potential in Nkp44+ mucosal NK cells during SIV infection. *Blood* **118**, 3321–3330 (2011).
34. Ivanov, I.I. *et al.* Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell* **139**, 485–498 (2009).
35. Nigam, P., Kwa, S., Velu, V. & Amara, R.R. Loss of IL-17-producing CD8T cells during late chronic stage of pathogenic simian immunodeficiency virus infection. *J. Immunol.* **186**, 745–753 (2011).
36. Raffatellu, M. *et al.* Simian immunodeficiency virus-induced mucosal interleukin-17 deficiency promotes Salmonella dissemination from the gut. *Nat. Med.* **14**, 421–428 (2008).
37. Aujja, S.J. *et al.* IL-22 mediates mucosal host defense against Gram-negative bacterial pneumonia. *Nat. Med.* **14**, 275–281 (2008).
38. Zheng, Y. *et al.* Interleukin-22 mediates early host defense against attaching and effacing bacterial pathogens. *Nat. Med.* **14**, 282–289 (2008).
39. Alter, G. & Altfeld, M. NK cells in HIV-1 infection: evidence for their role in the control of HIV-1 infection. *J. Intern. Med.* **265**, 29–42 (2009).
40. Lee, S.H., Miyagi, T. & Biron, C.A. Keeping NK cells in highly regulated antiviral warfare. *Trends Immunol.* **28**, 252–259 (2007).
41. Veazey, R.S. *et al.* Gastrointestinal tract as a major site of CD4+ T cell depletion and viral replication in SIV infection. *Science* **280**, 427–431 (1998).
42. Mehndru, S. *et al.* Primary HIV-1 infection is associated with preferential depletion of CD4+ T lymphocytes from effector sites in the gastrointestinal tract. *J. Exp. Med.* **200**, 761–770 (2004).
43. Favre, D. *et al.* Critical loss of the balance between Th17 and T regulatory cell populations in pathogenic SIV infection. *PLoS Pathog.* **5**, e1000295 (2009).
44. Brenchley, J.M. *et al.* Differential Th17 CD4T cell depletion in pathogenic and nonpathogenic lentiviral infections. *Blood* **112**, 2826–2835 (2008).
45. Hartigan-O'Connor, D.J., Hirao, L.A., McCune, J.M. & Dandekar, S. Th17 cells and regulatory T cells in elite control over HIV and SIV. *Curr. Opin. HIV AIDS* **6**, 221–227 (2011).
46. Brenchley, J.M. *et al.* Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nat. Med.* **12**, 1365–1371 (2006).
47. Estes, J.D. *et al.* Damaged intestinal epithelial integrity linked to microbial translocation in pathogenic simian immunodeficiency virus infections. *PLoS Pathog.* **6**, e1001052 (2010).
48. Klatt, N.R. & Brenchley, J.M. Th17 cell dynamics in HIV infection. *Curr. Opin. HIV AIDS* **5**, 135–140 (2010).
49. Cua, D.J. & Tato, C.M. Innate IL-17-producing cells: the sentinels of the immune system. *Nat. Rev. Immunol.* **10**, 479–489 (2010).
50. Eberl, G., Marmon, S., Sunshine, M.J., Rennert, P.D., Choi, Y. & Littman, D.R. An essential function for the nuclear receptor RORgamma(t) in the generation of fetal lymphoid tissue inducer cells. *Nat. Immunol.* **5**, 64–73 (2004).
51. Ouyang, W., Kolls, J.K. & Zheng, Y. The biological functions of T helper 17 cell effector cytokines in inflammation. *Immunity* **28**, 454–467 (2008).
52. Crellin, N.K., Trifari, S., Kaplan, C.D., Satoh-Takayama, N., Di Santo, J.P. & Spits, H. Regulation of cytokine secretion in human CD127(+) LTi-like innate lymphoid cells by Toll-like receptor 2. *Immunity* **33**, 752–764 (2010).
53. Ota, N. *et al.* IL-22 bridges the lymphotoxin pathway with the maintenance of colonic lymphoid structures during infection with *Citrobacter rodentium*. *Nat. Immunol.* **12**, 941–948 (2011).
54. Zaph, C. *et al.* Commensal-dependent expression of IL-25 regulates the IL-23-IL-17 axis in this intestine. *J. Exp. Med.* **205**, 2191–2198 (2008).
55. Klatt, N.R. *et al.* Compromised gastrointestinal integrity in pigtail macaques is associated with increased microbial translocation, immune activation, and IL-17 production in the absence of SIV infection. *Mucosal Immunol.* **3**, 387–398 (2010).
56. Douek, D. HIV disease progression: immune activation, microbes, and a leaky gut. *Top HIV Med.* **15**, 114–117 (2007).

57. Favre, D. *et al.* Tryptophan catabolism by indoleamine 2,3-dioxygenase 1 alters the balance of TH17 to regulatory T cells in HIV disease. *Sci. Transl. Med.* **2**, 32–36 (2010).
58. Rice, L. *et al.* CpG oligodeoxynucleotide protection in polymicrobial sepsis is dependent on interleukin-17. *J. Infect. Dis.* **191**, 1368–1376 (2005).
59. Xu, H. *et al.* Increased B7-H1 expression on dendritic cells correlates with programmed death 1 expression on T cells in simian immunodeficiency virus-infected macaques and may contribute to T cell dysfunction and disease progression. *J. Immunol.* **185**, 7340–7348 (2010).
60. Wang, X. *et al.* Monitoring alpha4beta7 integrin expression on circulating CD4+ T cells as a surrogate marker for tracking intestinal CD4+ T-cell loss in SIV infection. *Mucosal Immunol.* **2**, 518–526 (2009).
61. Wang, X. *et al.* Simian immunodeficiency virus selectively infects proliferating CD4+ T cells in neonatal rhesus macaques. *Blood* **116**, 4168–4174 (2010).