Gastrointestinal stromal tumors: what do we know now?

Christopher L Corless

Department of Pathology (L471) and Knight Diagnostic Laboratories, Oregon Health and Science University, Portland, OR, USA

Gastrointestinal stromal tumors (GISTs) are the most common mesenchymal tumors of the GI tract, arising from the interstitial cells of Cajal, primarily in the stomach and small intestine. They manifest a wide range of morphologies, from spindle cell to epithelioid, but are immunopositive for KIT (CD117) and/or DOG1 in essentially all cases. Although most tumors are localized at presentation, up to half will recur in the abdomen or spread to the liver. The growth of most GISTs is driven by oncogenic mutations in either of two receptor tyrosine kinases: KIT (75% of cases) or PDGFRA (10%). Treatment with tyrosine kinase inhibitors (TKIs) such as imatinib, sunitinib, and regorafenib is effective in controlling unresectable disease; however, drug resistance caused by secondary *KIT* or *PDGFRA* mutations eventually develops in 90% of cases. Adjuvant therapy with imatinib is commonly used to reduce the likelihood of disease recurrence after primary surgery, and for this reason assessing the prognosis of newly resected tumors is one of the most important roles for pathologists. Approximately 15% of GISTs are negative for mutations in *KIT* and *PDGFRA*. Recent studies of these so-called wild-type GISTs have uncovered a number of other oncogenic drivers, including mutations in neurofibromatosis type I, *RAS* genes, *BRAF*, and subunits of the succinate dehydrogenase complex. Routine genotyping is strongly recommended for optimal management of GISTs, as the type and dose of TKI used for treatment is dependent on the mutation identified.

Modern Pathology (2014) 27, S1–S16; doi:10.1038/modpathol.2013.173

Keywords: GIST; KIT; PDGFRA

Gastrointestinal stromal tumors (GISTs) occur primarily in older patients of either sex, with annual incidences between 11 and 19.6 per million population worldwide.^{1–4} This corresponds to between 3300 and 6000 new cases per year in the United States. Following surgical resection, GISTs often recur locally, spread diffusely throughout the serosal surfaces of the abdomen, and/or metastasize to the liver. Advanced disease is associated with metastases to distant sites, including the lung and bone. Before the advent of targeted therapies, the prognosis for advanced GISTs was poor owing to their inherent resistance to both chemotherapy and radiation therapy.

During the past decade, GISTs have served as an important model in the emerging field of molecularly targeted therapies for solid tumors. The nearly simul-

taneous discovery of oncogenic kinase mutations in GISTs and the introduction of kinase inhibitors have led to a rapid evolution in our understanding of these tumors and the biology that defines them.

Clinical and pathological features

GISTs are most commonly present in the stomach (60%) and small intestine (25%), but they also arise in the colon, rectum, esophagus, mesentery, and omentum (15% together).^{5,6} Clinical symptoms associated with GISTs include fatigue, abdominal pain, dysphagia, satiety, and obstruction. The workup often reveals anemia related to mucosal bleeding or intratumoral hemorrhage.

The tumors are generally well circumscribed, have a fleshy pink or tan cut surface, and may show areas of hemorrhagic necrosis and cystic degeneration. They range from 1 cm to more than 40 cm, with an average of ~5 cm. Morphologically, GISTs show a wide spectrum of morphologies, from bland spindle cell proliferations to highly cellular epithelioid tumors with significant nuclear pleomorphism (Figure 1). For these reasons, the morphologic

Correspondence: Dr CL Corless, MD, PhD, Department of Pathology (L471) and Knight Diagnostic Laboratories, Oregon Health and Science University, 3181 Southwest Sam Jackson Park Road, Portland, OR 97239, USA.
Email: corlessc@ohsu.edu

Received 23 May 2013; revised 15 June 2013; accepted 17 June 2013

Figure 1 Examples of gastrointestinal stromal tumor (GIST) morphology. Spindle cell GISTs (left panels) often harbor a mutation in *KIT* exon 9 or 11. Epithelioid GISTs (right panels) vary in their genotype, having either (or neither) a *KIT* or *PDGFRA* mutation.

differential diagnosis is necessarily broad (Table 1). Skeinoid fibers are associated with lower-grade lesions as are foci of calcification.

CD34 is expressed in 70% of GISTs and was the first immunohistochemical marker that helped to distinguish these tumors from leiomyomas and leiomyosarcomas of the GI tract. In 1998, reports by two groups that GISTs commonly express KIT (CD117) led to more reliable diagnosis.^{7,8} The staining may be membranous, diffusely cytoplasmic, or concentrated in a dot-like perinuclear pattern. The addition of DOG1 (ANO1) as another GIST marker has made the diagnosis quite routine,⁹ as DOG1 and CD117 each stain >95% of GISTs and, between them, serve to mark essentially all cases. These antigens are only rarely expressed in other mesenchymal tumors. Smooth muscle actin and muscle-specific actin are variably expressed in GISTs, whereas desmin is usually absent. Immunostaining for SDHA and SDHB has recently been shown to be effective in identifying tumors deficient

in succinate dehydrogenase activity,¹⁰ the significance of which is discussed below.

Oncogenic mutations in GISTs

KIT

In 1998, Hirota *et al*^{7,8} published their breakthrough discovery of KIT mutations in GISTs. Approximately 75% of GISTs harbor a *KIT* gene mutation, and these mutations lead to constitutive activation of the kinase. KIT is a member of the type III receptor tyrosine kinase family that includes platelet-derived growth factor receptors- α and - β (PDGFRA and PDGFRB), as well as the macrophage colony-stimulating-factor receptor (CSF1R) and the Fl cytokine receptor (FLT3).¹¹ Binding of the KIT ligand (stem cell factor, SCF) to KIT results in receptor homodimerization and kinase activation.

Table 1. Differential diagnosis of GIST

<i>Spindle cell tumors</i>	<i>Epithelioid tumors</i>
Fibromatosis (can be weakly CD117 positive)	Epithelioid leiomyoma
Leiomyoma (beware of intermixed CD117-positive ICC cells)	Neuroendocrine tumors
Leiomyosarcoma	Malignant mesothelioma
Schwannoma	Metastatic melanoma (can be CD117 positive)
Malignant peripheral nerve sheath tumor	Angiosarcoma (can be CD117 positive)
Inflammatory myofibroblastic tumor	
Inflammatory fibroid polyp (may have <i>PDGFRA</i> mutations)	
Solitary fibrous tumor	
Synovial sarcoma	
Dedifferentiated liposarcoma	
Endometrial stromal sarcoma	
Sarcomatoid carcinoma	

Abbreviations: GIST, gastrointestinal stromal tumor; ICC, interstitial cells of Cajal; *PDGFRA*, platelet-derived growth factor receptors- α .

Oncogenic *KIT* mutations cause ligand-independent kinase activation. The most common mutations in *KIT* affect the juxtamembrane domain encoded by exon 11 (Figure 2). Two-thirds of GISTs harbor mutations in exon 11 that disrupt the normal juxtamembrane secondary structure that keeps the kinase activation loop from swinging into an active conformation. These mutations include in-frame deletions, insertions, substitutions, or combinations thereof. The deletions are associated with a shorter progression-free and overall survival in comparison with the other exon 11 mutations.^{12–18} In particular, deletions involving codons 557 and/or 558 are associated with malignant behavior.^{19–21}

Aside from exon 11 mutations, between 7 and 10% of GISTs have a mutation in an extracellular domain encoded by exon 9.²² These mutations are thought to mimic the conformational change that the extracellular *KIT* receptor undergoes when ligand is bound. Importantly, the kinase domain in exon 9-mutant *KIT* is essentially the same as in wild-type *KIT*, and this influences inhibitor sensitivity. Interestingly, these mutations occur in tumors arising in the small and large intestine, but are very rarely seen in gastric GISTs.

Mutations in the activation loop (encoded by exon 17) of the kinase are uncommon. They stabilize the active conformation.²³ Primary mutations, such as K642E in the ATP-binding region (encoded by exon 13), are also uncommon.²³ The biological basis of kinase activation by this mutation is unknown, but it is speculated that it interferes with normal auto-inhibitory function of the juxtamembrane domain.

The functional importance of *KIT* mutations in GIST development is supported by several lines of evidence. First, phosphorylated (activated) *KIT* is detectable in GIST tumor extracts. Second, mutant *KIT* is oncogenic, supporting the growth of stably

Figure 2 *KIT* and *PDGFRA* mutations in gastrointestinal stromal tumor (GIST).

transfected BA/F3 cells in nude mice.^{7,24} Third, when expressed in transfected cell lines, mutant forms of *KIT* show constitutive kinase activity in the absence of SCF, as evidenced by autophosphorylation and activation of downstream signaling pathways.^{7,25} Finally, mice engineered to express *KIT* with mutations of the type found in human GISTs develop diffuse ICC hyperplasia within the muscular wall of the stomach and intestine.^{26,27} These mice also develop GIST-like tumors. This histologic picture is similar to that seen in individuals with inherited *KIT*-activating mutations.^{28,29}

Tumor extracts from *KIT*-mutant GISTs demonstrate evidence of activation of downstream signaling pathways, including the MAP kinase pathway (RAF, MEK, and ERK), the PI3 kinase/AKT pathway, and STAT3 (Figure 3a).^{24,30–32} The MAP kinase pathway upregulates important transcriptional regulators such as MYC, ELK, and CREB, and can stimulate the cell cycle through FOS. AKT activation through PI3 kinase and PDK1 leads to increased protein translation, downregulation of the cell cycle inhibitor p27^{KIP}, and anti-apoptotic effects. Recent studies have shown that ETV1 is an important regulator of GIST-specific gene expression during tumorigenesis.³³ *KIT* signaling through the MAP kinase pathway serves to maintain ETV1 activity.

In general, GISTs are heterozygous for a given mutation; however, in ~15% of tumors the remaining wild-type *KIT* allele is lost and this is associated with malignant behavior.^{29,34–36} In serial samples from individual patients, Chen *et al*³⁶ have provided evidence that this occurs through mitotic non-dysjunction, ie, failure of a chromosome 4 pair bearing the wild-type *KIT* allele to separate during mitosis, leaving one daughter cell with a single chromosome 4 containing the mutant *KIT* allele (uniparental monosomy). This correlates with increased mitotic activity and topoisomerase II expression.

Figure 3 (a) *KIT* and *PDGFRA* cell signaling pathways. Dimerization of *KIT* or platelet-derived growth factor receptors- α (*PDGFRA*) leads to signaling through the mitogen-activated protein (MAP) kinase pathway (RAF, MEK and ERK) and the phosphoinositide 3-kinase (PI3K) pathway (AKT, mammalian target of rapamycin (mTOR), S6 kinase). In addition, signal transducer and activator of transcription 3 (STAT3) is activated. The collective impact favors an increase in cell metabolism, cell cycle progression, and a decreased sensitivity to apoptosis. (b) Cell signaling in 'wild-type' gastrointestinal stromal tumors (GISTs). Mutations in neurofibromatosis type I (NF1), *RAS* genes, or *BRAF* lead to increased signaling through the MAP kinases MEK and ERK, promoting cell growth. Loss of the mitochondrial succinate dehydrogenase complex through mutations in *SDHA*, *SDHB*, *SDHC*, or *SDHD*, leads to an accumulation of succinate, which inhibits prolyl hydroxylase-mediated degradation of hypoxia-inducible factor-1 α (HIF1 α) through the proteasome complex. This results in upregulated transcription of a number of genes, including *VEGF*, *IGF1*, and *IGF2*. Succinate also inhibits demethylation of DNA by TET2 through increasing α -ketoglutarate levels.

Platelet-Derived Growth Factor Receptor- α

Immunoblots of GIST extracts from tumors lacking *KIT* gene mutations sometimes show high levels of phosphorylation of the α -receptor for platelet derived growth factor (*PDGFRA*), which is a close homolog of *KIT*.³⁷ *PDGFRA* is activated in GISTs harboring mutations in the juxtamembrane domain (exon 12), the ATP-binding domain (exon 14), or the activation loop (exon 18) (Figure 2 and Table 2). Consistent with their extensive functional

overlap, *KIT* and *PDGFRA* mutations are mutually exclusive in GISTs.

Observations supporting the significance of *PDGFRA* mutations in GIST parallel to those for *KIT* mutations. When expressed in transfected cell lines, mutant forms of *PDGFRA* have constitutive kinase activity in the absence of their ligand, *PDGF-A*,^{37,38} the activated downstream pathways are identical to those in *KIT*-mutant GISTs,^{37,39} and *PDGFRA* is also stabilized by HSP90.⁴⁰ In addition, both types of tumors are immunopositive

Table 2 Molecular Classification of GISTs

<i>Genetic type</i>	<i>Relative frequency (%)</i>	<i>Anatomic distribution</i>	<i>Germline example</i>
<i>KIT</i> mutation	75		
Exon 8	Rare	Small bowel	One kindred
Exon 9 ins AY502–503	8	Small bowel, colon	None
Exon 11 (deletions, single nucleotide substitutions, and insertions)	65	All sites	Several kindreds
Exon 13 K642E	1	All sites	Three kindreds
Exon 17 D820Y, N822K, and Y823D	1	All sites	Five kindreds
<i>PDGFRA</i> mutation	10		
Exon 12 (eg, V561D)	1	All sites	Two kindreds
Exon 14 N659K	Rare	Stomach	None
Exon 18 D842V	6	Stomach, mesentery, omentum	None
Exon 18 (eg, del IMHD 842-846)	2	All sites	One kindred
<i>KIT</i> and <i>PDGFRA</i> wild type	15	All sites	
<i>BRAF</i> V600E	~2		None
<i>SDHA/B/C/D</i> mutations	~6	Stomach and small bowel	Carney–Stratakis
<i>HRAS</i> , <i>NRAS</i> , and <i>PIK3CA</i> mutation	<1		None
Pediatric/Carney triad	~1	Stomach	Not heritable
NF1-related	<1	Small bowel	Numerous

Abbreviations: GIST, gastrointestinal stromal tumor; ICC, interstitial cells of Cajal; PDGFRA, platelet-derived growth factor receptors- α ; NF1, neurofibromatosis type I.

for the markers DOG1 and protein kinase C- θ (PKC θ).^{9,41,42} These markers are highly selective for GISTs over other mesenchymal tumors. Further, as discussed below, both genotypes are associated with cytogenetic changes that are distinctive for GIST.^{37,43}

Despite these molecular similarities, *PDGFRA*-mutant GISTs do show features distinctive from *KIT*-mutant GISTs, including differences in gene expression profile,^{39,44} a striking predilection for the stomach, variable (sometimes negative) expression of KIT,^{20,41,45–48} and a generally lower potential for malignancy.⁴⁹ Morphologically, however, *PDGFRA*-mutant GISTs are not reliably distinguishable from *KIT*-mutant GISTs (Figure 1).

Other Driver Mutations

Approximately 15% of GISTs do not have a detectable mutation in either *KIT* or *PDGFRA*. In other respects, these so-called ‘wild-type’ GISTs are clinically indistinguishable from *KIT*- or *PDGFRA*-mutant GISTs, having identical morphology, expressing high levels of KIT, and occurring anywhere in the GI tract. Phosphorylated KIT is detectable in these tumors, suggesting that KIT is still activated,³⁰ but the mechanism of this activation is unclear. However, recent studies have revealed that wild-type GISTs are a heterogeneous group and display various oncogenic mutations (Table 2). For example, the *BRAF* V600E substitution common in papillary thyroid carcinoma and melanoma is present in up to 13% of wild-type GISTs.⁵⁰ *HRAS*, *NRAS* and *PIK3CA* gene mutations also occur, but are much more rare. As *BRAF* and the RAS proteins are constituents of the MAP kinase signaling cascade,

they can result in *KIT*-independent growth stimulation (Figure 3b) and are possible causes for resistance to *KIT*/*PDGFRA* kinase inhibitors.

It is estimated that 7% of patients with neurofibromatosis type I (NF1) develop one or more GISTs.^{51–56} Most arise in the small intestine and they do not readily metastasize.⁵¹ The majority of these GISTs are wild type for *KIT* and *PDGFRA*, but as expected they do show either somatic mutation or loss of the remaining wild-type *NF1* allele, resulting in signaling through the MAP kinase cascade (Figure 3b).^{51,54,55,57}

SDH-Deficient GISTs

Approximately half of all wild-type GISTs show loss of respiratory chain complex II enzymatic activity.^{10,58} This complex comprises four subunits (SDHA, SDHB, SDHC, and SDHD) and serves to oxidize succinate to fumarate as part of the mitochondrial Krebs cycle. Loss of any of these subunits through gene mutation or post-transcriptional downregulation destabilizes the complex and causes the accumulation of succinate. This results in increased transcription of HIF1 α -regulated genes and decreased DNA demethylation (Figure 3b).⁵⁹ Indeed, SDH-deficient GISTs show a global increase in DNA methylation similar to that seen in gliomas and leukemias with IDH1 and IDH2 mutations.⁶⁰

Germline mutations in *SDHA*, *SDHB*, or *SDHC* increase the risk not only for the development of one or more SDH-deficient GISTs but also for paragangliomas (Carney–Stratakis syndrome).^{10,61} The GISTs in affected patients show either loss or somatic mutation (second hit) of the remaining

wild-type allele. The absence of immunostaining for SDHA, as well as SDHB, is helpful in identifying GISTs with possible SDH gene mutations.^{58,58,62–66}

Approximately 50% of wild-type GISTs demonstrate high expression of IGF1R.⁶⁷ In SDH-deficient GISTs, upregulation of IGF1 and IGF2 may activate IGF1R in an autocrine manner (Figure 3b), resulting in signaling through both the MAP kinase and PI3 kinase/AKT pathways.⁶⁸

GISTs that arise in pediatric patients (~1–2% of all GISTs) are predominantly SDHB immunonegative, but rarely harbor an *SDH* gene mutation. These tumors, which often metastasize but tend to grow slowly, have a different gene expression signature from *KIT*/*PDGFRA*-mutant GISTs.^{69–71} Some pediatric-type GISTs are accompanied by pulmonary chondromas and/or paragangliomas, referred to as Carney triad, a non-heritable syndrome, the genetic cause for which has yet to be determined.⁷²

The origin of GISTs

Interstitial Cells Of Cajal

During the 1990s, a number of investigators noted similarities between GISTs and a population of cells in the GI tract called the interstitial cells of Cajal (ICCs), which serve as pacemakers for peristaltic contractions. These observations led to the hypothesis that ICCs could be the cell-of-origin of GISTs. Mice engineered to express *KIT* with mutations of the type found in human GISTs develop diffuse ICC hyperplasia within the muscular wall of the stomach and intestine.^{26,27} These mice also develop GIST-like tumors. Diffuse ICC hyperplasia has been described in several kindreds with heritable mutations in *KIT* (Table 2), and is associated with dysphagia and the development of multiple GISTs,^{26,29,53,73–78} although many of the tumors do not follow a malignant course.

The relationship between GISTs and ICCs is further supported by parallels in gene expression. For example, high levels of *PKCθ*, *nestin*, and *DOG1* are expressed in both GISTs and ICCs. In addition, the ETS family transcription factor *ETV1* is highly expressed in both GISTs and in the specific subpopulations of ICCs (myenteric and intramuscular, as opposed to submucosal) thought to give rise to GISTs.³³

The observation that some *KIT* and *PDGFRA* mutations in GISTs correlate closely with anatomical location (Table 2) might be explained by their ICC origin. For example, GISTs with a *KIT* exon 9 mutation, which arise primarily in the intestines, may derive from a different subgroup of ICCs than those with a *PDGFRA* D842V mutation, which occur only in the stomach, mesentery, and omentum. The more common *KIT* mutations, by contrast, can be found in GISTs throughout the GI tract, perhaps deriving from a more ubiquitous ICC subtype.

Micro-GISTs

Minute growths (1–10 mm) of ICC/GIST-like cells are present in between 2.9 and 35% of the stomachs thoroughly examined after surgical removal or at autopsy.^{79–82} These so-called micro-GISTs are mitotically inactive and often partially calcified, suggesting tumorigenic arrest. In contrast to the diffuse ICC hyperplasia observed in the presence of a germline *KIT* mutation, micro-GISTs appear to represent a nodular form of ICC overgrowth caused in most cases by local, somatic acquisition of a *KIT* mutation. The type and frequency of *KIT* mutations in micro-GISTs is essentially the same as in clinically significant tumors.⁸³ Subcentimeter GISTs with *PDGFRA* mutations have also been reported.⁷⁹ These observations on micro-GISTs suggest that kinase gene mutations occur very early in GIST tumorigenesis; however, the mutations are probably not sufficient for progression to an oncologically threatening lesion (Figure 4). The large pool of micro-GISTs in the general population likely explains the multiple reported cases in which two or more genotypically distinct GISTs are found in a patient during a single surgical procedure.^{53,79,84}

Chromosomal and Molecular Alterations During GIST Progression

Although oncogenic kinase mutations have a significant role in the development of GISTs, other genetic events are important in their clinical progression (Figure 4). Approximately two-thirds

Figure 4 Origin and progression of gastrointestinal stromal tumors (GISTs). Under the influence of a *KIT* or platelet-derived growth factor receptors- α (*PDGFRA*) mutation, a clonal outgrowth of interstitial cells of Cajal (ICCs) may form a discrete nodule (micro-GIST) measuring <1 cm and showing essentially no mitotic activity. A conservative estimate for the prevalence of micro-GISTs in the US population would exceed 10 million lesions. With the accumulation of additional mutations and chromosomal changes, a micro-GIST may progress to a larger lesion that is clinically significant. Approximately 5000 such lesions are identified each year in the United States, of which 2000 are of high risk or are malignant.

of GISTs demonstrate either monosomy of chromosome 14, or partial loss of 14q.^{37,85–89} Interestingly, these chromosome 14 abnormalities are observed in both *KIT*-mutant and *PDGFRA*-mutant GISTs.^{37,43} Deletions of 14q11.2 include the genes *PARP2*, *APEX1*, and *NDRG2*, whereas deletions of 14q32 include the *SIVA* gene.⁹⁰ Loss of the long arm of chromosome 22 is observed in ~50% of GISTs.^{37,43,85,86,89,91}

Losses on chromosomes 1p, 9p, 11p, and 17p are successively less common than 14q and 22q losses, but are more significantly associated with malignancy (Figure 4).^{37,43,85,89,91–94} Losses on chromosomes 10, 13q, and 15q have also been reported in GISTs.^{43,91} Gains on chromosomes 8q (including *MYC*), 3q (including *SMARCA3*), and 17q are associated with metastatic behavior.^{86,95}

In a recent array-based analysis of gene copy number in 42 GISTs (23 with recurrence or metastasis), the tumors were separated into four groups reflecting their accumulated chromosomal changes. The overall survival of group 1 (loss of 22q, 19, and 1p distal) and group 2 (additional loss of 14q) was significantly better than that of group 3 (added losses of 15q and 1p proximal) and group 4 (additional loss of 10). Specific genes implicated in this analysis included *OXA1L* on 14q, as well as *AKAP13* and *C15orf5* on 15q.

None of the above karyotypic changes is present in pediatric-type GISTs, which remain near-diploid, again emphasizing the different biology of these tumors. In contrast, GISTs arising in NF1 patients often show losses of 14q and 22q.⁹⁶

On the basis of gene expression profiling of high-risk versus low-risk GISTs, the high-risk tumors show significant changes in genes regulating the cell cycle, including genes influenced by the PI3 kinase pathway and genes involved in the G2/M checkpoint.⁹⁷ A significant fraction of malignant GISTs show inactivation of the tumor suppressor gene *CDKN2A* (which encodes the cell cycle regulatory protein p16^{INK4A}) through chromosome 9p21 deletion, either biallelic or in combination with mutation or promoter methylation.^{98–101} *TP53* mutations and decreased p53 immunostaining also correlate with a poor prognosis.^{102–104} Likewise, amplifications of *MDM2* and *CCND1* (cyclin D1), although uncommon in GISTs, are associated with malignancy.¹⁰⁵

Kinase mutations and tyrosine kinase inhibitor therapy

Until the year 2000, treatment options for patients with advanced GIST were poor. The response rate to conventional chemotherapy was <5% and median survival for patients with advanced disease was ~18 months.

The tyrosine kinase inhibitor (TKI) imatinib was developed in the early 1990s as a treatment for chronic myelogenous leukemia because of its

ability to inhibit the fusion oncoprotein BCR–ABL.¹⁰⁶ The observation that ABL shares structural similarity with KIT, and several other tyrosine kinases led to experiments showing that imatinib can inhibit the growth of cells expressing mutant forms of KIT.²⁵ In addition, imatinib showed potent activity against a KIT-mutant GIST cell line. Imatinib inhibits KIT kinase by directly binding to the ATP pocket and competitively inhibiting ATP binding. The KIT receptor is normally in equilibrium between active and inactive conformations. The latter is favored by steric hindrance conferred by the juxtamembrane domain (exon 11), which prevents the activation loop from assuming the conformation required for kinase activation.

With the knowledge that imatinib inhibits KIT signaling, imatinib was first used clinically to treat a 50-year-old female with metastatic GIST, and a dramatic response was observed.¹⁰⁷ Promising results from phase I and II trials led to two international phase III trials, each using similar protocols that would allow a subsequent meta-analysis. The phase III trials compared 400 versus 800mg daily doses of imatinib. Overall, imatinib achieved disease control in 70–85% of patients with advanced KIT-immunopositive GISTs, and the median progression-free survival was 20–24 months.^{108–112} Currently, the median survival for patients with advanced disease treated with frontline imatinib is 5 years, with 34% of patients surviving more than 9 years.¹¹³

A meta-analysis of the phase III trials proved that patients with exon 9-mutant GIST had a significantly longer progression-free survival if they were treated with 800mg of imatinib; hence, this is now the target dose for these patients. In contrast, only 400mg of imatinib is needed to manage exon 11-mutant tumors.¹¹²

The success with imatinib in patients with metastatic disease quickly led to trials of the drug in the adjuvant setting. It was initially shown that 12 months of imatinib after primary resection of a GIST significantly delayed disease recurrence versus a placebo.¹¹⁴ A subsequent trial proved that 36 months of adjuvant imatinib was superior to 12 months.¹¹⁵

Assessing GIST prognosis

FDA approval for the use of imatinib in the adjuvant setting has made it imperative that the prognosis of a newly resected GIST be predicted as accurately as possible. In a 2-year prospective study of sarcomas in France, >85% of GISTs presented as localized lesions.¹¹⁶ Thus, the question of whether to use adjuvant therapy must be considered for the great majority of newly diagnosed GISTs. Kinase genotype does not factor into overall survival once a GIST becomes metastatic. Thus, a *KIT* or *PDGFRA* mutation may set the initial course of a GIST, but the prognosis at the time of clinical presentation is

Table 3 Assessing GIST prognosis

	Size (cm)	Gastric (n = 1055), %	Jejunum/ileum (n = 629), %	Duodenum (n = 144), %	Rectum (n = 111), %
Mitotic index $\leq 5/5 \text{ mm}^2$	≤ 2	0	0	0	0
	$> 2 \leq 5$	1.9	4.3	8.3	8.5
	$> 5 \leq 10$	3.6	24	Insufficient data	Insufficient data
	> 10	10	52	34	57
Mitotic index $> 5/5 \text{ mm}^2$	≤ 2	None	High	Insufficient data	54
	$> 2 \leq 5$	16	73	50	52
	$> 5 \leq 10$	55	85	Insufficient data	Insufficient data
	> 10	86	90	86	71

Abbreviations: GIST, gastrointestinal stromal tumor.

The risk of disease recurrence or metastasis can be estimated based on three parameters defined in this table: (1) the mitotic index (either ≤ 5 mitoses/ 5 mm^2 or > 5 mitoses/ 5 mm^2); (2) tumor size (largest diameter); and (3) tumor site of origin. The table is based on data published by Miettinen *et al.*^{117,118,150}

clearly influenced by other genetic events. Unfortunately, our knowledge of these additional mutations remains limited, and current recommendations for assessing prognosis are based on three simple parameters: tumor size, tumor location, and mitotic index (mitoses/ 5 mm^2). A number of risk assessment schemes have been published; the most widely used scheme was developed at the Armed Forces Institute of Pathology by Miettinen *et al.*^{117–119}, whose considerable efforts in studying the outcome of patients before the advent TKIs have provided the most complete data available (Table 3). It should be noted that tumor rupture, either before or during surgery, is another important negative prognostic factor. Incomplete resection, particularly in the area of the rectum, is also associated with a higher risk of recurrence.¹²⁰

As a group, PDGFRA-mutant GISTs appear to be less aggressive than KIT-mutant GISTs,⁴⁹ yet PDGFRA-mutant tumors can still progress and kill patients. These tumors are assessed using the same criteria as other GISTs.

Responses to TKI therapy

Clinical Disease Persistence

Clinical data suggest that even long-term TKI treatment fails to eradicate GIST cells, resulting in disease persistence. In an attempt to determine the optimal duration of imatinib therapy for advanced, unresectable GIST, an interesting trial randomized patients who had continuous control of their disease during 3 years of imatinib treatment to either continue or to discontinue treatment.¹²¹ For those continuing treatment progression-free survival over the next 2 years was 80%, but for those who stopped therapy it was only 16%. Patients who relapsed after discontinuation of therapy did so because of persistent disease. In contrast, the progression that developed in some of the patients who maintained therapy was due to resistant disease.

Persistence of GIST cells during TKI treatment could be due to the failure of these drugs to eradicate mature GIST cells and/or the failure to eliminate GIST stem cells. Current evidence suggests that both mechanisms underlie GIST persistence in the face of prolonged TKI therapy. Agaram *et al.*¹²² examined a series of 43 clinically responsive GISTs lesions from 28 patients. Histological responses in these resected tumors after 1–31 months of imatinib treatment ranged from $< 10\%$ to $> 90\%$ reduction in tumor cellularity, but tumor cells persisted in all lesions. Three quarters of the lesions showed an absence of mitoses and a proliferative index of 0% by Ki-67 staining, indicating biological quiescence. Interestingly, some of the tumor cells showed transdifferentiation. Indeed, examples of rhabdomyoblastic, cartilaginous, and osseous differentiation have all been observed in cases of treated GIST (Figure 5). Thus, under imatinib suppression, GIST cells may avoid apoptosis by exiting the cell cycle and expressing genes associated with a differentiated phenotype. Whether these cells proliferate again when a TKI is discontinued, or whether GIST stem cells are the source of renewed growth, has not been determined.

In rare cases, GISTs may progress to high grade, anaplastic sarcomas that lose CD117 expression. This has been observed in both imatinib-treated and TKI-naïve tumors.¹²³

Resistance to TKI therapy

Primary Resistance

Resistance to treatment with KIT/PDGFRA inhibitors such as imatinib can be divided into two types: primary and secondary. Approximately 10% of patients with GIST have primary resistance, defined as progression within the first 6 months of treatment. On the basis of data from phase II and phase III trials, tumor response to imatinib correlates with the underlying kinase genotype.^{34,41,112,124,125} The

Figure 5 Differentiation of gastrointestinal stromal tumor (GIST) cells under suppression of a tyrosine kinase inhibitor (TKI). In some cases, imatinib treatment may lead to rhabdomyoblastic differentiation (left panel), or else chondroid or osseous differentiation (right panels).

probability of primary resistance to imatinib for *KIT* exon 11, *KIT* exon 9, and wild-type GISTs is 5, 16 and 23%, respectively.³⁴ These findings likely reflect the underlying sensitivities of different *KIT* genotypes. Exon 11-mutant *KIT* is highly sensitive to imatinib, with an IC_{50} of <100 nM, whereas exon 9-mutant *KIT* and wild-type *KIT* are less sensitive to the drug ($IC_{50} \sim 1000$ nM for each).¹²⁶ Thus, underdosing of imatinib in patients with exon 9 mutations probably accounts for some of the apparent resistance.¹¹²

On the basis of *in vitro* data, the most common *PDGFRA* mutation in GISTs, D842V, is fully resistant to the effects of imatinib.^{34,38,127,128} This mutation favors the active conformation of the kinase domain and consequently disfavors imatinib binding.^{34,129,130} This is corroborated by clinical results, as patients with *PDGFRA* D842V-mutant GIST have low response rates and very short progression-free and overall survivals during imatinib treatment. Crenolanib is a TKI that has

activity versus D842V and is now being tested in a clinical trial. There are, however, other *PDGFRA* mutations that are sensitive to imatinib *in vitro* and patients with these mutations have shown durable responses to imatinib.

Wild-type GISTs include tumors with mutations downstream of *KIT*,^{10,50,131,132} hence, these subsets of wild-type GISTs might respond better to other targeted agents, such as *VEGFR* inhibitors for pediatric/*SDH*-deficient GIST and *BRAF*/*MEK* inhibitors for *BRAF* and *RAS* mutant GIST.¹³³

Secondary Resistance

After an initial response to imatinib, the vast majority of patients eventually develop disease progression. This secondary resistance may manifest in a number of ways, including growth of a nodule within a pre-existing, clinically quiescent lesion, the development of one or more new lesions, or

widespread disease throughout the liver or abdominal cavity. It is now established that acquired mutations in *KIT* or *PDGFRA* account for most secondary resistance, and that these mutations occur almost exclusively in the same gene and allele as the primary oncogenic driver mutation.^{35,134–140}

In a phase II imatinib study for advanced GIST, 67% of the patients whose tumor showed imatinib resistance had a new, or secondary, mutation in *KIT*. Notably, these mutations were common among tumors with a primary exon 11 mutation, but were not observed in wild-type GIST samples.¹³⁷ Unlike primary mutations that activate *KIT*, which are predominantly in the juxtamembrane regions encoded by exons 9 and 11, the secondary mutations were concentrated in two regions of the *KIT* kinase domain. One is the ATP-binding pocket, encoded by exons 13 and 14, mutations of which directly interfere with drug binding. The second is the activation loop (exons 17 and 18), where mutations can stabilize *KIT* in the active conformation and thereby hinder drug interaction. Drug resistance has also been observed in *PDGFRA*-mutant GISTs, in which the most common one is an acquired D842V mutation (activation loop).^{135,137}

Additional studies using more sensitive assays have identified secondary mutations in >80% of drug-resistant GIST lesions.^{141–143} More sobering is that there is significant heterogeneity of resistance across different lesions, and even within different areas of the same lesion. For example, there are reports of up to five different drug resistance mutations in different portions of an individual lesion and up to seven different secondary resistance mutations across multiple tumors in the same patient.¹⁴¹ This heterogeneity of resistance significantly impacts the efficacy of salvage TKI therapy after frontline imatinib, because the diversity of resistant, minority clones precludes the systemic suppression of GIST cells by any particular TKI.

Approaches to imatinib-resistant GIST

Most GIST patients who develop secondary resistance to imatinib will not respond to dose escalation, forcing a switch to an alternative *KIT*/*PDGFRA* TKI. Such salvage agents include sunitinib, sorafenib, regorafenib, vatalanib, masatinib, nilotinib, and dasatinib, as well as other investigational inhibitors. Although all of these agents are *KIT*/*PDGFRA* inhibitors, the majority of them (in contrast to imatinib) also target *VEGFR1/2*.¹⁴⁴ Whether the disease stabilization that can be seen with these salvage agents is related to *VEGFR1/2* inhibition and a consequent decrease in angiogenesis remains unclear.

Sunitinib is FDA-approved for the treatment of GIST patients with progression on imatinib,¹⁴⁵ but biochemical evidence suggests that its activity against secondary imatinib-resistant kinase

mutations is suboptimal. *KIT* ATP-binding pocket mutations are extremely sensitive to sunitinib *in vitro*; however, the activation loop mutations are strongly cross-resistant. Given the approximately equal frequency of these different classes of mutations in imatinib-resistant lesions and the multiplicity of lesions in a typical patient, it is not surprising that mixed responses are common during sunitinib therapy.^{124,146}

Most recently, the FDA has approved regorafenib for the third-line treatment of patients with GIST that is resistant to imatinib and sunitinib. In a phase III trial, the progression-free survival on regorafenib was 4.8 months.¹⁴⁷ Thus, even with newer drugs such as regorafenib, resistance develops over time, suggesting that escape from ATP-competitive inhibitors of *KIT* and *PDGFRA* is inevitable. A new class of non-ATP mimetic kinase inhibitors (switch pocket kinase inhibitors, such as DP-2976) have shown high potency when tested *in vitro* and represent hope in the fight against TKI resistance.^{148,149}

Summary and future directions

Achievements in the treatment of GISTs during the past decade are the direct result of a growing understanding of their molecular biology. The high frequency of primary *KIT* and *PDGFRA* mutations in these tumors makes them sensitive to kinase inhibitors such as imatinib, and this drug is FDA-approved for use in both the adjuvant and advanced disease settings. Accurate assessment of the risk of disease recurrence is essential in determining the use of adjuvant imatinib after resection of a primary tumor. Current recommendations for assessing recurrence risk are based on tumor size, tumor location, and mitotic index (mitoses/5 mm²).^{117–119}

Resistance to imatinib develops in the majority of advanced cases of GIST. An immediate research goal is the development of new inhibitors that have activity against secondary mutations in the activation loop. In addition, the development of effective combination therapy is likely to improve tumor control. Ongoing high-throughput genomic studies may identify additional drivers/modifiers of GIST biology that can be targeted.

Clinical genotyping of GISTs is important to identify *KIT* exon 9-mutant tumors that require a higher dose of imatinib for optimal disease control. Further, there are other molecular subtypes of GIST that do not respond well to conventional *KIT* inhibitors, but may be better treated with other agents (eg, *SDH*-deficient GIST and *PDGFRA* D842V GIST). Thus, the GIST genotyping is critical in personalizing the care of patients who need TKI therapy.

In summary, new insights into the origin and progression of GISTs are setting the stage for further therapeutic innovations, with the goal not just to

control disease growth, but to eliminate all tumor cells at the time of initial therapy. Going forward, the challenge will be to move from a paradigm of tumor suppression to true cancer cure.

Acknowledgments

We thank all the members, past and present, of the Corless and Heinrich laboratories at the Oregon Health and Science University for their continuing efforts on GIST research. Some of the works referenced in this article were supported by generous donations from the GIST Cancer Research Fund and the BP Lester Foundation, and by grant support from the LifeRaft Group.

Disclosure/conflict of interest

CLC has received honoraria, consulting fees, and research support from Novartis and Pfizer. CLC has also received a consulting fee from Bayer.

References

- Chan KH, Chan CW, Chow WH, *et al.* Gastrointestinal stromal tumors in a cohort of Chinese patients in Hong Kong. *World J Gastroenterol* 2006;12: 2223–2228.
- Goettsch WG, Bos SD, Breekveldt-Postma N, *et al.* Incidence of gastrointestinal stromal tumours is underestimated: Results of a nation-wide study. *Eur J Cancer* 2005;41:2868–2872.
- Nilsson B, Bummig P, Meis-Kindblom JM, *et al.* Gastrointestinal stromal tumors: the incidence, prevalence, clinical course, and prognostication in the preimatinib mesylate era. *Cancer* 2005;103:821–829.
- Tryggvason G, Gislason HG, Magnusson MK, *et al.* Gastrointestinal stromal tumors in Iceland, 1990–2003: the Icelandic GIST study, a population-based incidence and pathologic risk stratification study. *Int J Cancer* 2005;117:289–293.
- Corless CL, Fletcher JA, Heinrich MC. Biology of gastrointestinal stromal tumors. *J Clin Oncol* 2004;22: 3813–3825.
- Miettinen M, Lasota J. Gastrointestinal stromal tumors (GISTs): definition, occurrence, pathology, differential diagnosis and molecular genetics. *Pol J Pathol* 2003;54:3–24.
- Hirota S, Isozaki K, Moriyama Y, *et al.* Gain-of-function mutations of c-kit in human gastrointestinal stromal tumors. *Science* 1998;279:577–580.
- Kindblom LG, Remotti HE, Aldenborg F, *et al.* Gastrointestinal pacemaker cell tumor (GIPACT): gastrointestinal stromal tumors show phenotypic characteristics of the interstitial cells of Cajal. *Am J Pathol* 1998;152:1259–1269.
- West RB, Corless CL, Chen X, *et al.* The novel marker, DOG1, is expressed ubiquitously in gastrointestinal stromal tumors irrespective of KIT or PDGFRA mutation status. *Am J Pathol* 2004;165: 107–113.
- Janeway KA, Kim SY, Lodish M, *et al.* Defects in succinate dehydrogenase in gastrointestinal stromal tumors lacking KIT and PDGFRA mutations. *Proc Natl Acad Sci USA* 2011;108:314–318.
- Hanks SK, Quinn AM, Hunter T. The protein kinase family: Conserved features and deduced phylogeny of the catalytic domains. *Science* 1988;241:42–52.
- Ernst SI, Hubbs AE, Przygodzki RM, *et al.* KIT mutation portends poor prognosis in gastrointestinal stromal/smooth muscle tumors. *Lab Invest* 1998;78: 1633–1636.
- Lasota J, Jasinski M, Sarlomo-Rikala M, *et al.* Mutations in exon 11 of c-Kit occur preferentially in malignant versus benign gastrointestinal stromal tumors and do not occur in leiomyomas or leiomyosarcomas. *Am J Pathol* 1999;154:53–60.
- Singer S, Rubin BP, Lux ML, *et al.* Prognostic value of KIT mutation type, mitotic activity, and histologic subtype in gastrointestinal stromal tumors. *J Clin Oncol* 2002;20:3898–3905.
- Taniguchi M, Nishida T, Hirota S, *et al.* Effect of c-kit mutation on prognosis of gastrointestinal stromal tumors. *Cancer Res* 1999;59:4297–4300.
- Andersson J, Bummig P, Meis-Kindblom JM, *et al.* Gastrointestinal stromal tumors with KIT exon 11 deletions are associated with poor prognosis. *Gastroenterology* 2006;130:1573–1581.
- Cho S, Kitadai Y, Yoshida S, *et al.* Deletion of the KIT gene is associated with liver metastasis and poor prognosis in patients with gastrointestinal stromal tumor in the stomach. *Int J Oncol* 2006;28: 1361–1367.
- Liu XH, Bai CG, Xie Q, *et al.* Prognostic value of KIT mutation in gastrointestinal stromal tumors. *World J Gastroenterol* 2005;11:3948–3952.
- Martin J, Poveda A, Llombart-Bosch A, *et al.* Deletions affecting codons 557–558 of the c-KIT gene indicate a poor prognosis in patients with completely resected gastrointestinal stromal tumors: a study by the Spanish Group for Sarcoma Research (GEIS). *J Clin Oncol* 2005;23:6190–6198.
- Tzen CY, Mau BL. Analysis of CD117-negative gastrointestinal stromal tumors. *World J Gastroenterol* 2005;11:1052–1055.
- Wardelmann E, Losen I, Hans V, *et al.* Deletion of Trp-557 and Lys-558 in the juxtamembrane domain of the c-kit protooncogene is associated with metastatic behavior of gastrointestinal stromal tumors. *Int J Cancer* 2003;106:887–895.
- Lux ML, Rubin BP, Biase TL, *et al.* KIT extracellular and kinase domain mutations in gastrointestinal stromal tumors. *Am J Pathol* 2000;156: 791–795.
- Lasota J, Corless CL, Heinrich MC, *et al.* Clinicopathologic profile of gastrointestinal stromal tumors (GISTs) with primary KIT exon 13 or exon 17 mutations: a multicenter study on 54 cases. *Mod Pathol* 2008;21:476–484.
- Rubin BP, Singer S, Tsao C, *et al.* KIT activation is a ubiquitous feature of gastrointestinal stromal tumors. *Cancer Res* 2001;61:8118–8121.
- Heinrich MC, Griffith DJ, Druker BJ, *et al.* Inhibition of c-kit receptor tyrosine kinase activity by STI 571, a selective tyrosine kinase inhibitor. *Blood* 2000;96: 925–932.
- Sommer G, Agosti V, Ehlers I, *et al.* Gastrointestinal stromal tumors in a mouse model by targeted

- mutation of the Kit receptor tyrosine kinase. *Proc Natl Acad Sci USA* 2003;100:6706–6711.
- 27 Rubin BP, Antonescu CR, Scott-Browne JP, *et al.* A knock-in mouse model of gastrointestinal stromal tumor harboring kit K641E. *Cancer Res* 2005;65: 6631–6639.
 - 28 Chen H, Hirota S, Isozaki K, *et al.* Polyclonal nature of diffuse proliferation of interstitial cells of Cajal in patients with familial and multiple gastrointestinal stromal tumours. *Gut* 2002;51:793–796.
 - 29 O'riain C, Corless CL, Heinrich MC, *et al.* Gastrointestinal stromal tumors: insights from a new familial GIST kindred with unusual genetic and pathologic features. *Am J Surg Pathol* 2005;29: 1680–1683.
 - 30 Duensing A, Medeiros F, McConarty B, *et al.* Mechanisms of oncogenic KIT signal transduction in primary gastrointestinal stromal tumors (GISTs). *Oncogene* 2004;23:3999–4006.
 - 31 Rossi F, Ehlers I, Agosti V, *et al.* Oncogenic Kit signaling and therapeutic intervention in a mouse model of gastrointestinal stromal tumor. *Proc Natl Acad Sci USA* 2006;103:12843–12848.
 - 32 Duensing A, Joseph N, Liu C, *et al.* Protein kinase C theta (PKC(theta)) is selectively expressed and constitutively phosphorylated in gastrointestinal stromal tumors (GISTs). *AACR Meet Abstr* 2004; 2004:708.
 - 33 Chi P, Chen Y, Zhang L, *et al.* ETV1 is a lineage survival factor that cooperates with KIT in gastrointestinal stromal tumours. *Nature* 2010;467: 849–853.
 - 34 Heinrich MC, Corless CL, Demetri GD, *et al.* Kinase mutations and imatinib response in patients with metastatic gastrointestinal stromal tumor. *J Clin Oncol* 2003;21:4342–4349.
 - 35 Antonescu CR, Besmer P, Guo T, *et al.* Acquired resistance to imatinib in gastrointestinal stromal tumor occurs through secondary gene mutation. *Clin Cancer Res* 2005;11:4182–4190.
 - 36 Chen LL, Holden JA, Choi H, *et al.* Evolution from heterozygous to homozygous KIT mutation in gastrointestinal stromal tumor correlates with the mechanism of mitotic nondisjunction and significant tumor progression. *Mod Pathol* 2008;21:826–836.
 - 37 Heinrich MC, Corless CL, Duensing A, *et al.* PDGFRA activating mutations in gastrointestinal stromal tumors. *Science* 2003;299:708–710.
 - 38 Hirota S, Ohashi A, Nishida T, *et al.* Gain-of-function mutations of platelet-derived growth factor receptor alpha gene in gastrointestinal stromal tumors. *Gastroenterology* 2003;125:660–667.
 - 39 Kang HJ, Nam SW, Kim H, *et al.* Correlation of KIT and platelet-derived growth factor receptor alpha mutations with gene activation and expression profiles in gastrointestinal stromal tumors. *Oncogene* 2005;24:1066–1074.
 - 40 Matei D, Satpathy M, Cao L, *et al.* The platelet-derived growth factor receptor alpha is destabilized by geldanamycins in cancer cells. *J Biol Chem* 2007;282:445–453.
 - 41 Debiec-Rychter M, Wasag B, Stul M, *et al.* Gastrointestinal stromal tumours (GISTs) negative for KIT (CD117 antigen) immunoreactivity. *J Pathol* 2004;202: 430–438.
 - 42 Duensing A, Joseph NE, Medeiros F, *et al.* Protein kinase C theta (PKC(theta)) expression and constitutive activation in gastrointestinal stromal tumors (GISTs). *Cancer Res* 2004;64:5127–5131.
 - 43 Wozniak A, Sciort R, Guillou L, *et al.* Array CGH analysis in primary gastrointestinal stromal tumors: cytogenetic profile correlates with anatomic site and tumor aggressiveness, irrespective of mutational status. *Genes Chromosomes Cancer* 2007;46:261–276.
 - 44 Subramanian S, West RB, Corless CL, *et al.* Gastrointestinal stromal tumors (GISTs) with KIT and PDGFRA mutations have distinct gene expression profiles. *Oncogene* 2004;23:7780–7790.
 - 45 Wasag B, Debiec-Rychter M, Pauwels P, *et al.* Differential expression of KIT/PDGFRA mutant isoforms in epithelioid and mixed variants of gastrointestinal stromal tumors depends predominantly on the tumor site. *Mod Pathol* 2004;17:889–894.
 - 46 Medeiros F, Corless CL, Duensing A, *et al.* KIT-negative gastrointestinal stromal tumors: proof of concept and therapeutic implications. *Am J Surg Pathol* 2004;28:889–894.
 - 47 Sakurai S, Hasegawa T, Sakuma Y, *et al.* Myxoid epithelioid gastrointestinal stromal tumor (GIST) with mast cell infiltrations: a subtype of GIST with mutations of platelet-derived growth factor receptor alpha gene. *Hum Pathol* 2004;35:1223–1230.
 - 48 Wardelmann E, Hrychyk A, Merkelbach-Bruse S, *et al.* Association of platelet-derived growth factor receptor alpha mutations with gastric primary site and epithelioid or mixed cell morphology in gastrointestinal stromal tumors. *J Mol Diagn* 2004;6: 197–204.
 - 49 Lasota J, Dansonka-Mieszkowska A, Sobin LH, *et al.* A great majority of GISTs with PDGFRA mutations represent gastric tumors of low or no malignant potential. *Lab Invest* 2004;84:874–883.
 - 50 Hostein I, Faur N, Primois C, *et al.* BRAF mutation status in gastrointestinal stromal tumors. *Am J Clin Pathol* 2010;133:141–148.
 - 51 Andersson J, Sihto H, Meis-Kindblom JM, *et al.* NF1-associated gastrointestinal stromal tumors have unique clinical, phenotypic, and genotypic characteristics. *Am J Surg Pathol* 2005;29:1170–1176.
 - 52 de Raedt T, Cools J, Debiec-Rychter M, *et al.* Intestinal neurofibromatosis is a subtype of familial GIST and results from a dominant activating mutation in PDGFRA. *Gastroenterology* 2006;131:1907–1912.
 - 53 Kang DY, Park CK, Choi JS, *et al.* Multiple gastrointestinal stromal tumors: clinicopathologic and genetic analysis of 12 patients. *Am J Surg Pathol* 2007;31:224–232.
 - 54 Maertens O, Prenen H, Debiec-Rychter M, *et al.* Molecular pathogenesis of multiple gastrointestinal stromal tumors in NF1 patients. *Hum Mol Genet* 2006;15:1015–1023.
 - 55 Miettinen M, Fetsch JF, Sobin LH, *et al.* Gastrointestinal stromal tumors in patients with neurofibromatosis 1: a clinicopathologic and molecular genetic study of 45 cases. *Am J Surg Pathol* 2006;30: 90–96.
 - 56 Stewart C, Corless C, Rubin BP, *et al.* Mitotic recombination as evidence of alternative pathogenesis of gastrointestinal stromal tumours in neurofibromatosis type 1. *J Med Genet* 2007;44:e61.
 - 57 Kinoshita K, Hirota S, Isozaki K, *et al.* Absence of c-kit gene mutations in gastrointestinal stromal tumours from neurofibromatosis type 1 patients. *J Pathol* 2004;202:80–85.

- 58 Miettinen M, Killian JK, Wang ZF, *et al.* Immunohistochemical loss of succinate dehydrogenase subunit A (SDHA) in gastrointestinal stromal tumors (GISTs) signals SDHA germline mutation. *Am J Surg Pathol* 2012;37:234–240.
- 59 Mason E, Hornick JL. Succinate dehydrogenase deficiency is associated with decreased 5-hydroxymethylcytosine production in gastrointestinal stromal tumors: implications for mechanisms of tumorigenesis. *Mod Pathol*; advance online publication, 7 June 2013 (e-pub ahead of print).
- 60 Killian JK, Kim SY, Miettinen M, *et al.* Succinate dehydrogenase mutation underlies global epigenomic divergence in gastrointestinal stromal tumor. *Cancer Discov* 2013;3:648–657.
- 61 Pantaleo MA, Astolfi A, Indio V, *et al.* SDHA loss-of-function mutations in KIT-PDGFRα wild-type gastrointestinal stromal tumors identified by massively parallel sequencing. *J Natl Cancer Inst* 2011;103:983–987.
- 62 Wagner AJ, Remillard SP, Zhang YX, *et al.* Loss of expression of SDHA predicts SDHA mutations in gastrointestinal stromal tumors. *Mod Pathol* 2013;26:289–294.
- 63 Dwight T, Benn DE, Clarkson A, *et al.* Loss of SDHA expression identifies SDHA mutations in succinate dehydrogenase-deficient gastrointestinal stromal tumors. *Am J Surg Pathol* 2012;37:226–233.
- 64 Pantaleo MA, Astolfi A, Urbini M, *et al.* Analysis of all subunits, SDHA, SDHB, SDHC, SDHD, of the succinate dehydrogenase complex in KIT/PDGFRα wild-type GIST. *Eur J Hum Genet* 2013 (e-pub ahead of print).
- 65 Oudijk L, Gaal J, Korpershoek E, *et al.* SDHA mutations in adult and pediatric wild-type gastrointestinal stromal tumors. *Mod Pathol* 2013;3:456–463.
- 66 Italiano A, Chen CL, Sung YS, *et al.* SDHA loss of function mutations in a subset of young adult wild-type gastrointestinal stromal tumors. *BMC Cancer* 2012;14:408.
- 67 Belinsky MG, Rink L, Flieder DB, *et al.* Overexpression of insulin-like growth factor 1 receptor and frequent mutational inactivation of SDHA in wild-type SDHB-negative gastrointestinal stromal tumors. *Genes Chromosomes Cancer* 2013;52:214–224.
- 68 Nannini M, Astolfi A, Paterini P, *et al.* Expression of IGF-1 receptor in KIT/PDGFRα wild-type gastrointestinal stromal tumors with succinate dehydrogenase complex dysfunction. *Future Oncol* 2013;9:121–126.
- 69 Prakash S, Sarraf L, Socci N, *et al.* Gastrointestinal stromal tumors in children and young adults: a clinicopathologic, molecular, and genomic study of 15 cases and review of the literature. *J Pediatr Hematol Oncol* 2005;27:179–187.
- 70 Janeway KA, Liegl B, Harlow A, *et al.* Pediatric KIT wild-type and platelet-derived growth factor receptor alpha-wild-type gastrointestinal stromal tumors share KIT activation but not mechanisms of genetic progression with adult gastrointestinal stromal tumors. *Cancer Res* 2007;67:9084–9088.
- 71 Agaram NP, LaQuaglia MP, Ustun B, *et al.* Molecular characterization of pediatric gastrointestinal stromal tumors. *Clin Cancer Res* 2008;14:3204–3215.
- 72 Carney JA. Gastric stromal sarcoma, pulmonary chondroma, and extra-adrenal paraganglioma (Carney Triad): natural history, adrenocortical component, and possible familial occurrence. *Mayo Clin Proc* 1999;74:543–552.
- 73 Beghini A, Tibiletti MG, Roversi G, *et al.* Germline mutation in the juxtamembrane domain of the kit gene in a family with gastrointestinal stromal tumors and urticaria pigmentosa. *Cancer* 2001;92:657–662.
- 74 Chompret A, Kannengiesser C, Barrois M, *et al.* PDGFRα germline mutation in a family with multiple cases of gastrointestinal stromal tumor. *Gastroenterology* 2004;126:318–321.
- 75 Hirota S, Nishida T, Isozaki K, *et al.* Familial gastrointestinal stromal tumors associated with dysphagia and novel type germline mutation of KIT gene. *Gastroenterology* 2002;122:1493–1499.
- 76 Isozaki K, Terris B, Belghiti J, *et al.* Germline-activating mutation in the kinase domain of KIT gene in familial gastrointestinal stromal tumors. *Am J Pathol* 2000;157:1581–1585.
- 77 Maeyama H, Hidaka E, Ota H, *et al.* Familial gastrointestinal stromal tumor with hyperpigmentation: association with a germline mutation of the c-kit gene. *Gastroenterology* 2001;120:210–215.
- 78 Nishida T, Hirota S, Taniguchi M, *et al.* Familial gastrointestinal stromal tumours with germline mutation of the KIT gene. *Nat Genet* 1998;19:323–324.
- 79 Agaimy A, Wunsch PH, Hofstaedter F, *et al.* Minute gastric sclerosing stromal tumors (GIST tumorlets) are common in adults and frequently show c-KIT mutations. *Am J Surg Pathol* 2007;31:113–120.
- 80 Corless CL, McGreevey L, Haley A, *et al.* KIT mutations are common in incidental gastrointestinal stromal tumors one centimeter or less in size. *Am J Pathol* 2002;160:1567–1572.
- 81 Kawanowa K, Sakuma Y, Sakurai S, *et al.* High incidence of microscopic gastrointestinal stromal tumors in the stomach. *Hum Pathol* 2006;37:1527–1535.
- 82 Muenst S, Thies S, Went P, *et al.* Frequency, phenotype, and genotype of minute gastrointestinal stromal tumors in the stomach: an autopsy study. *Hum Pathol* 2011;42:1849–1854.
- 83 Rossi S, Gasparotto D, Toffolatti L, *et al.* Molecular and clinicopathologic characterization of gastrointestinal stromal tumors (GISTs) of small size. *Am J Surg Pathol* 2010;34:1480–1491.
- 84 Gasparotto D, Rossi S, Bearzi I, *et al.* Multiple primary sporadic gastrointestinal stromal tumors in the adult: an underestimated entity. *Clin Cancer Res* 2008;14:5715–5721.
- 85 Bergmann F, Gunawan B, Hermanns B, *et al.* Cytogenetic and morphologic characteristics of gastrointestinal stromal tumors. Recurrent rearrangement of chromosome 1 and losses of chromosomes 14 and 22 as common anomalies. *Verh Dtsch Ges Pathol* 1998;82:275–278.
- 86 Debiec-Rychter M, Lasota J, Sarlomo-Rikala M, *et al.* Chromosomal aberrations in malignant gastrointestinal stromal tumors: correlation with c-KIT gene mutation. *Cancer Genet Cytogenet* 2001;128:24–30.
- 87 Fukasawa T, Chong JM, Sakurai S, *et al.* Allelic loss of 14q and 22q, NF2 mutation, and genetic instability occur independently of c-kit mutation in gastrointestinal stromal tumor. *Jpn J Cancer Res* 2000;91:1241–1249.
- 88 Heinrich MC, Rubin BP, Longley BJ, *et al.* Biology and genetic aspects of gastrointestinal stromal tumors: KIT activation and cytogenetic alterations. *Hum Pathol* 2002;33:484–495.

- 89 Kim NG, Kim JJ, Ahn JY, *et al.* Putative chromosomal deletions on 9P, 9Q and 22Q occur preferentially in malignant gastrointestinal stromal tumors. *Int J Cancer* 2000;85:633–638.
- 90 Assamaki R, Sarlomo-Rikala M, Lopez-Guerrero JA, *et al.* Array comparative genomic hybridization analysis of chromosomal imbalances and their target genes in gastrointestinal stromal tumors. *Genes Chromosomes Cancer* 2007;46:564–576.
- 91 Gunawan B, von HA, Sander B, *et al.* An oncogenetic tree model in gastrointestinal stromal tumours (GISTs) identifies different pathways of cytogenetic evolution with prognostic implications. *J Pathol* 2007;211:463–470.
- 92 El Rifai W, Sarlomo-Rikala M, Miettinen M, *et al.* DNA copy number losses in chromosome 14: an early change in gastrointestinal stromal tumors. *Cancer Res* 1996;56:3230–3233.
- 93 O'Leary T, Ernst S, Przygodzki R, *et al.* Loss of heterozygosity at 1p36 predicts poor prognosis in gastrointestinal stromal/smooth muscle tumors. *Lab Invest* 1999;79:1461–1467.
- 94 Schurr P, Wolter S, Kaifi J, *et al.* Microsatellite DNA alterations of gastrointestinal stromal tumors are predictive for outcome. *Clin Cancer Res* 2006;12:5151–5157.
- 95 el-Rifai W, Sarlomo-Rikala M, Andersson LC, *et al.* High-resolution deletion mapping of chromosome 14 in stromal tumors of the gastrointestinal tract suggests two distinct tumor suppressor loci. *Genes Chromosomes Cancer* 2000;27:387–391.
- 96 Yamamoto H, Tobo T, Nakamori M, *et al.* Neurofibromatosis type 1-related gastrointestinal stromal tumors: a special reference to loss of heterozygosity at 14q and 22q. *J Cancer Res Clin Oncol* 2009;135:791–798.
- 97 Hur K, Lee HJ, Woo JH, *et al.* Gene expression profiling of human gastrointestinal stromal tumors according to its malignant potential. *Dig Dis Sci* 2010;55:2561–2567.
- 98 Perrone F, Tamborini E, Dagrada GP, *et al.* 9p21 locus analysis in high-risk gastrointestinal stromal tumors characterized for c-kit and platelet-derived growth factor receptor alpha gene alterations. *Cancer* 2005;104:159–169.
- 99 Ricci R, Arena V, Castri F, *et al.* Role of p16/INK4a in gastrointestinal stromal tumor progression. *Am J Clin Pathol* 2004;122:35–43.
- 100 Sabah M, Cummins R, Leader M, *et al.* Loss of heterozygosity of chromosome 9p and loss of p16INK4A expression are associated with malignant gastrointestinal stromal tumors. *Mod Pathol* 2004;17:1364–1371.
- 101 Schneider-Stock R, Boltze C, Lasota J, *et al.* Loss of p16 protein defines high-risk patients with gastrointestinal stromal tumors: a tissue microarray study. *Clin Cancer Res* 2005;11(2 Pt 1):638–645.
- 102 Feakins RM. The expression of p53 and bcl-2 in gastrointestinal stromal tumours is associated with anatomical site, and p53 expression is associated with grade and clinical outcome. *Histopathology* 2005;46:270–279.
- 103 Panizo-Santos A, Sola I, Vega F, *et al.* Predicting metastatic risk of gastrointestinal stromal tumors: role of cell proliferation and cell cycle regulatory proteins. *Int J Surg Pathol* 2000;8:133–144.
- 104 Romeo S, Debiec-Rychter M, Van Glabbeke M, *et al.* Cell cycle/apoptosis molecule expression correlates with imatinib response in patients with advanced gastrointestinal stromal tumors. *Clin Cancer Res* 2009;15:4198.
- 105 Tornillo L, Duchini G, Carafa V, *et al.* Patterns of gene amplification in gastrointestinal stromal tumors (GIST). *Lab Invest* 2005;85:921–931.
- 106 Druker BJ, Tamura S, Buchdunger E, *et al.* Effects of a selective inhibitor of the Abl tyrosine kinase on the growth of Bcr-Abl positive cells. *Nat Med* 1996;2:561–566.
- 107 Joensuu H, Roberts PJ, Sarlomo-Rikala M, *et al.* Effect of the tyrosine kinase inhibitor STI571 in a patient with a metastatic gastrointestinal stromal tumor. *N Engl J Med* 2001;344:1052–1056.
- 108 Van Oosterom AT, Judson I, Verweij J, *et al.* STI571, an active drug in metastatic gastrointestinal stromal tumors (GIST) an EORTC phase I study [abstract]. *Proc Am Soc Clin Oncol* 2001;20:1a.
- 109 Demetri GD, von Mehren M, Blanke CD, *et al.* Efficacy and safety of imatinib mesylate in advanced gastrointestinal stromal tumors. *N Engl J Med* 2002;347:472–480.
- 110 Blanke CD, Rankin C, Demetri GD, *et al.* Phase III randomized, intergroup trial assessing imatinib mesylate at two dose levels in patients with unresectable or metastatic gastrointestinal stromal tumors expressing the kit receptor tyrosine kinase: S0033. *J Clin Oncol* 2008;26:626–632.
- 111 Verweij J, Casali PG, Zalcberg J, *et al.* Progression-free survival in gastrointestinal stromal tumours with high-dose imatinib: randomised trial. *Lancet* 2004;364:1127–1134.
- 112 Gastrointestinal Stromal Tumor Meta-Analysis Group (MetaGIST). Comparison of two doses of imatinib for the treatment of unresectable or metastatic gastrointestinal stromal tumors: a meta-analysis of 1,640 patients. *J Clin Oncol* 2010;28:1247–1253.
- 113 von Mehren M, Heinrich MC, Joensuu H, *et al.* Follow-up results after 9 years (yrs) of the ongoing, phase II B222 trial of imatinib mesylate (IM) in patients (pts) with metastatic or unresectable KIT⁺ gastrointestinal stromal tumors (GIST). *J Clin Oncol* 2011;29:609s(Suppl) abstr 10016.
- 114 DeMatteo RP, Ballman KV, Antonescu CR, *et al.* Adjuvant imatinib mesylate after resection of localised, primary gastrointestinal stromal tumour: a randomised, double-blind, placebo-controlled trial. *Lancet* 2009;373:1097–1104.
- 115 Joensuu H, Eriksson M, Sundby Hall K, *et al.* One vs three years of adjuvant imatinib for operable gastrointestinal stromal tumor: a randomized trial. *JAMA* 2012;307:1265–1272.
- 116 Cassier PA, Ducimetière F, Lurkin A, *et al.* A prospective epidemiological study of new incident GISTs during two consecutive years in Rhône Alpes region: incidence and molecular distribution of GIST in a European region. *Br J Cancer* 2010;103:165–170.
- 117 Miettinen M, Sobin LH, Lasota J. Gastrointestinal stromal tumors of the stomach: a clinicopathologic, immunohistochemical, and molecular genetic study of 1765 cases with long-term follow-up. *Am J Surg Pathol* 2005;29:52–68.
- 118 Miettinen M, Makhlof H, Sobin LH, *et al.* Gastrointestinal stromal tumors of the jejunum and ileum: a

- clinicopathologic, immunohistochemical, and molecular genetic study of 906 cases before imatinib with long-term follow-up. *Am J Surg Pathol* 2006;30:477–489.
- 119 Miettinen M, Kopczynski J, Makhoulf HR, *et al.* Gastrointestinal stromal tumors, intramural leiomyomas, and leiomyosarcomas in the duodenum: a clinicopathologic, immunohistochemical, and molecular genetic study of 167 cases. *Am J Surg Pathol* 2003;27:625–641.
 - 120 Agaimy A, Vassos N, Märkl B, *et al.* Anorectal gastrointestinal stromal tumors: a retrospective multicenter analysis of 15 cases emphasizing their high local recurrence rate and the need for standardized therapeutic approach. *Int J Colorectal Dis* 2013;28:1057–1064.
 - 121 Le Cesne A, Ray-Coguard I, Bui BN, *et al.* Discontinuation of imatinib in patients with advanced gastrointestinal stromal tumours after 3 years of treatment: an open-label multicentre randomised phase 3 trial. *Lancet Oncol* 2010;11:942–949.
 - 122 Agaram NP, Besmer P, Wong GC, *et al.* Pathologic and molecular heterogeneity in imatinib-stable or imatinib-responsive gastrointestinal stromal tumors. *Clin Cancer Res* 2007;13:170–181.
 - 123 Antonescu CR, Romeo S, Zhang L, *et al.* Dedifferentiation in gastrointestinal stromal tumor to an anaplastic KIT-negative phenotype: a diagnostic pitfall: morphologic and molecular characterization of 8 cases occurring either *de novo* or after imatinib therapy. *Am J Surg Pathol* 2013;37:385–392.
 - 124 Heinrich MC, Owzar K, Corless CL, *et al.* Correlation of kinase genotype and clinical outcome in the North American Intergroup Phase III Trial of imatinib mesylate for treatment of advanced gastrointestinal stromal tumor: CALGB 150105 Study by Cancer and Leukemia Group B and Southwest Oncology Group. *J Clin Oncol* 2008;26:5360–5367.
 - 125 Debiec-Rychter M, Sciort R, Le CA, *et al.* Kit mutations and dose selection for imatinib in patients with advanced gastrointestinal stromal tumours. *Eur J Cancer* 2006;42:1093–1103.
 - 126 Heinrich MC, Maki RG, Corless CL, *et al.* Primary and secondary kinase genotypes correlate with the biological and clinical activity of sunitinib in imatinib-resistant gastrointestinal stromal tumor. *J Clin Oncol* 2008;26:5352–5359.
 - 127 Corless CL, Schroeder A, Griffith D, *et al.* PDGFRA mutations in gastrointestinal stromal tumors: frequency, spectrum and *in vitro* sensitivity to imatinib. *J Clin Oncol* 2005;23:5357–5364.
 - 128 Weisberg E, Wright RD, Jiang J, *et al.* Effects of PKC412, nilotinib, and imatinib against GIST-associated PDGFRA mutants with differential imatinib sensitivity. *Gastroenterology* 2006;131:1734–1742.
 - 129 Gajiwala KS, Wu JC, Christensen J, *et al.* KIT kinase mutants show unique mechanisms of drug resistance to imatinib and sunitinib in gastrointestinal stromal tumor patients. *Proc Natl Acad Sci USA* 2009;106:1542–1547.
 - 130 Cassier PA, Fumagalli E, Rutkowski P, *et al.* Outcome of patients with platelet-derived growth factor receptor alpha-mutated gastrointestinal stromal tumors in the tyrosine kinase inhibitor era. *Clin Cancer Res* 2012;18:4458–4464.
 - 131 Agaimy A, Terracciano LM, Dirnhofer S, *et al.* V600E BRAF mutations are alternative early molecular events in a subset of KIT/PDGFRA wild-type gastrointestinal stromal tumours. *J Clin Pathol* 2009;62:613–616.
 - 132 Agaram NP, Wong GC, Guo T, *et al.* Novel V600E BRAF mutations in imatinib-naïve and imatinib-resistant gastrointestinal stromal tumors. *Genes Chromosomes Cancer* 2008;47:853–859.
 - 133 Janeway KA, Albritton KH, Van den Abbeele AD, *et al.* Sunitinib treatment in pediatric patients with advanced GIST following failure of imatinib. *Pediatr Blood Cancer* 2009;52:767–771.
 - 134 Chen LL, Trent JC, Wu EF, *et al.* A missense mutation in KIT kinase domain 1 correlates with imatinib resistance in gastrointestinal stromal tumors. *Cancer Res* 2004;64:5913–5919.
 - 135 Debiec-Rychter M, van OA, Marynen P. Mechanisms of resistance to imatinib mesylate in gastrointestinal stromal tumors and activity of the PKC412 inhibitor against imatinib-resistant mutants. *Gastroenterology* 2005;128:270–279.
 - 136 Grimpen F, Yip D, McArthur G, *et al.* Resistance to imatinib, low-grade FDG-avidity on PET, and acquired KIT exon 17 mutation in gastrointestinal stromal tumour. *Lancet Oncol* 2005;6:724–727.
 - 137 Heinrich MC, Corless CL, Blanke CD, *et al.* Molecular correlates of imatinib resistance in gastrointestinal stromal tumors. *J Clin Oncol* 2006;24:4764–4774.
 - 138 Koyama T, Nimura H, Kobayashi K, *et al.* Recurrent gastrointestinal stromal tumor (GIST) of the stomach associated with a novel c-kit mutation after imatinib treatment. *Gastric Cancer* 2006;9:235–239.
 - 139 Wakai T, Kanda T, Hirota S, *et al.* Late resistance to imatinib therapy in a metastatic gastrointestinal stromal tumour is associated with a second KIT mutation. *Br J Cancer* 2004;90:2059–2061.
 - 140 Wardelmann E, Thomas N, Merkelbach-Bruse S, *et al.* Acquired resistance to imatinib in gastrointestinal stromal tumours caused by multiple KIT mutations. *Lancet Oncol* 2005;6:249–251.
 - 141 Liegl B, Kepten I, Le C, *et al.* Heterogeneity of kinase inhibitor resistance mechanisms in GIST. *J Pathol* 2008;216:64–74.
 - 142 Nishida T, Kanda T, Nishitani A, *et al.* Secondary mutations in the kinase domain of the KIT gene are predominant in imatinib-resistant gastrointestinal stromal tumor. *Cancer Sci* 2008.
 - 143 Lim KH, Huang MJ, Chen LT, *et al.* Molecular analysis of secondary kinase mutations in imatinib-resistant gastrointestinal stromal tumors. *Med Oncol* 2008;25:207–213.
 - 144 Demetri GD. Differential properties of current tyrosine kinase inhibitors in gastrointestinal stromal tumors. *Semin Oncol* 2011;38(Suppl 1):S10–S19.
 - 145 Demetri GD, Van Oosterom AT, Garrett CR, *et al.* Efficacy and safety of sunitinib in patients with advanced gastrointestinal stromal tumour after failure of imatinib: a randomised controlled trial. *Lancet* 2006;368:1329–1338.
 - 146 Nishida T, Takahashi T, Nishitani A, *et al.* Sunitinib-resistant gastrointestinal stromal tumors harbor cis-mutations in the activation loop of the KIT gene. *Int J Clin Oncol* 2009;14:143–149.
 - 147 Demetri GD, Reichardt P, Kang Y, *et al.* Efficacy and safety of regorafenib for advanced gastrointestinal stromal tumours after failure of imatinib and

- sunitinib (GRID): an international, multicentre, randomised, placebo-controlled, phase 3 trial. [abstract]. *Lancet* 2013;381:295–302.
- 148 Eide CA, Adrian LT, Tyner JW, *et al.* The ABL switch control inhibitor DCC-2036 is active against the chronic myeloid leukemia mutant BCR-ABL T315I and exhibits a narrow resistance profile. *Cancer Res* 2011;71:3189–3195.
- 149 Heinrich MC, Wise S, Hood M, *et al.* *In vitro* activity of novel KIT/PDGFRα switch pocket kinase inhibitors against mutations associated with drug-resistant GI stromal tumors. *J Clin Oncol* 2010;28:15s(Suppl) abstr 10007.
- 150 Lasota J, Miettinen M. KIT and PDGFRα mutations in gastrointestinal stromal tumors (GISTs). *Semin Diagn Pathol* 2006;23:91–102.