Aggressive non-alcoholic steatohepatitis following rapid weight loss and/or malnutrition

Jia-Huei Tsai^{1,2}, Linda D Ferrell³, Vivian Tan⁴, Matthew M Yeh⁵, Monika Sarkar⁶ and Ryan M Gill³

¹Department of Pathology, National Taiwan University Hospital, Taipei, Taiwan; ²Graduate Institute of Pathology, College of Medicine, National Taiwan University, Taipei, Taiwan; ³Department of Pathology, University of California, San Francisco, CA, USA; ⁴Genomic Health, Redwood City, CA, USA; ⁵Department of Pathology, University of Washington, Seattle, WA, USA and ⁶Division of Gastroenterology and Hepatology, Department of Medicine, University of California, San Francisco, CA, USA

While non-alcoholic steatohepatitis is a slowly progressive disease, patients may rarely present in acute liver failure. We describe six patients who developed severe hepatic dysfunction following rapid weight loss or malnutrition. Rapid weight loss (18 to 91 kg) occurred after Roux-en-Y gastric bypass in four patients and starvation-like dieting or hypoalbuminemia was noted in two patients. Four patients either died or received an urgent liver transplant. Pathologic findings were characterized by advanced alcoholic steatohepatitis-like features, including extensive/circumferential centrilobular pericellular fibrosis, central scar with perivenular sclerosis/veno-occlusion with superimposed hepatocellular dropout, abundant/prominent hepatocellular balloons, and numerous Mallory–Denk bodies, but there was no history of excess alcohol consumption. This study characterizes clinicopathologic features of aggressive non-alcoholic steatohepatitis following rapid weight loss or malnutrition, which should be included in the differential diagnosis with alcohol when a patient is considered for liver transplantation. The mechanism of liver injury in aggressive steatohepatitis is unknown, but rapid fat mobilization in obese patients may potentially cause oxidative stress to the liver and further study is needed to determine if there is a genetic predisposition to this form of injury and if antioxidants may protect the liver during rapid weight loss/malnutrition.

Modern Pathology (2017) **30**, 834–842; doi:10.1038/modpathol.2017.13; published online 3 March 2017

The term ‘non-alcoholic steatohepatitis’ was first proposed by Ludwig *et al.*¹ in 1980 to denote steatohepatitis in absence of alcohol abuse. Well-established risk factors for non-alcoholic steatohepatitis include obesity, metabolic syndrome, hyperlipidemia, and type 2 diabetes.^{2,3} Non-alcoholic steatohepatitis prevalence is increasing around the world, especially in the United States,⁴ and is expected to be the most common etiology of cirrhosis (and the most common indication for liver transplantation) in coming years.⁵ A multi-hit model has been proposed to explain the pathogenesis of non-alcoholic steatohepatitis.³ Proposed mechanisms of disease include proinflammatory cytokines, oxidative stress, and endoplasmic reticulum stress, as well

as an interplay between gut microbiome, diet, and genetic influences.³ Diagnosis of non-alcoholic steatohepatitis requires a liver biopsy with steatosis, lobular inflammation, and hepatocellular ballooning (with or without characteristic pericellular fibrosis).^{6,7} Other common histologic findings in non-alcoholic steatohepatitis include Mallory–Denk bodies, acidophil bodies, glycogenosis, and glycogenated nuclei. Alcoholic steatohepatitis often manifests with more intense portal and lobular inflammation with neutrophil satellitosis, cholestasis, central vein obliteration, more prominent ballooned hepatocytes, and abundant Mallory–Denk bodies.⁶

Non-alcoholic steatohepatitis typically progresses slowly and may not be clinically apparent.² Up to 20% of patients with non-alcoholic steatohepatitis progress to cirrhosis over a 15-year time period.⁸ Hepatic decompensation in non-alcoholic steatohepatitis is rare and has been associated with drug use, rapid weight loss/malnutrition due to fasting, and intestinal resection or bariatric surgery.^{9–16} Rapid

Correspondence: Dr RM Gill, MD, PhD, Department of Pathology, University of California San Francisco, 505 Parnassus Avenue, M545, Box 0102, San Francisco, CA 94143, USA.
E-mail: Ryan.Gill@ucsf.edu

Received 6 December 2016; revised 25 January 2017; accepted 26 January 2017; published online 3 March 2017

weight loss/malnutrition has been reported to induce hepatic inflammation and exacerbate steatohepatitis with progression to liver failure within a relatively short timeframe.^{17,18} However, liver injury due to rapid weight loss/malnutrition remains poorly understood and pathologic features have not been well-characterized. In this study, we describe six patients with rapid weight loss/malnutrition who developed severe hepatic dysfunction and histologic findings suggestive of an aggressive form of non-alcoholic steatohepatitis.

Materials and methods

Case Selection

To identify cases of potential aggressive non-alcoholic steatohepatitis, and following approval by the Committee on Human Research, the Department of Pathology, University of California, San Francisco (San Francisco, CA, USA) files were searched from 2000 to 2015 for 'steatohepatitis.' All cases suggestive of rapid progression/decompensation were selected for further review. Cases were excluded if there was any history of alcohol or drug abuse. We also received one case from the University of Washington Department of Pathology (Seattle, WA, USA). Demographic information is provided in Table 1.

All cases were evaluated by JHT, LDF, and RMG to document the following pathologic features: centrilobular scarring, perivenular sclerosis, centrilobular arteries,¹⁹ pericellular fibrosis, Mallory–Denk bodies, hepatocellular ballooning, ductular reaction, and cholestasis. Severe centrilobular scarring was defined as broad central-based fibrosis with extensive and circumferential pericellular fibrosis involving a majority of central zones. Trichrome and elastic stains (orcein stain or Elastic Van Gieson stain) were used to assess fibrosis and subacute hepatocellular dropout.^{20,21}

Results

Demographics and Clinical History

All patients were female with a median age of 39 years (range 33–46 years; Table 1) and had a history of rapid weight loss or malnutrition. There was no history of prior liver disease. Patient 2 had detectable Epstein–Barr virus IgM, but clinical and histologic findings did not support acute Epstein–Barr virus hepatitis as a cause of liver failure. Five patients were morbidly obese; three underwent Roux-en-Y gastric bypass, one received long-limb Roux-en-Y gastric bypass, and one followed a low carbohydrate/high protein diet for weight loss. There was no history of alcohol abuse or diet pill/supplement use. An intraoperative liver biopsy from Roux-en-Y gastric bypass was available in patient 1, which

showed only moderate steatosis and early centrilobular pericellular fibrosis. The only non-obese patient in our study had hypoalbuminemia, suggestive of malnutrition, but no further history was available.

Clinical Presentation and Outcome

All patients presented with acute or subacute hepatic dysfunction, ranging from severe jaundice to asterixis, prominent hepatomegaly, and encephalopathy (with symptoms reported from 1 to 12 weeks prior to admission). There were two deaths and two patients received a liver transplant for severe liver failure soon after admission. The liver condition in one patient improved after medical management and one follow-up biopsy was taken three years later. Another patient had no available follow-up information.

Pathologic Findings

Pathologic findings are summarized in Table 2. Specimens included two explants (patients 2 and 3), two autopsies (patients 1 and 4), and two needle biopsies (patients 5 and 6). Massive hepatomegaly with fatty change was found in patients 1 and 4, which weighed 6680 and 5600g, respectively. The other two liver explants, in patients 2 and 3, showed micronodular cirrhosis and weighed 1825 and 1491g, respectively.

All histologic sections showed evidence of steatohepatitis, which was characterized by severe central-based fibrous scars (Figure 1a and b), prominent hepatocellular ballooning, and abundant dense Mallory–Denk bodies (Figure 1c). There were apoptotic hepatocytes in lobular parenchyma and a prominent neutrophil infiltrate around ballooned hepatocytes in all cases. Centrilobular arteries were present to a variable extent in five patients (Figure 1d). There was focal ductular reaction, mostly in a periportal distribution, in five patients, while hepatocanalicular cholestasis was only identified in three patients (Figure 1e). Perivenular sclerosis was observed in five patients and was characterized by fibrous wall thickening, intimal fibrosis, or total venous sclerosis with luminal obliteration. (Figure 1f).

Trichrome stain was used to estimate extent of centrilobular scarring (Figure 2a) and perivenular sclerosis (Figure 2b). Most central zones showed severe fibrosis and were involved by broad fibrous scars with extensive and circumferential pericellular fibrosis throughout the centrilobular region (Zone 3), which also extended into Zone 2 (midzonal) and focally to the periportal region (Zone 1), with associated bridging fibrosis (ie, central-central bridge formation). Centrilobular two-tone trichrome staining intensity was evident in four patients (Figure 2c) with dense aggregates of thick collagen bundles in darker staining areas and, at most, thin collagen fibers dispersed loosely in lighter staining areas,

Table 1 Patient characteristics and laboratory data on admission

	<i>Patient 1</i>	<i>Patient 2</i>	<i>Patient 3</i>	<i>Patient 4</i>	<i>Patient 5</i>	<i>Patient 6</i>
Age	33	37	40	37	37	46
Gender	F	F	F	F	F	F
Ethnicity	Hispanic	Caucasian	Hispanic	Caucasian	Caucasian	Caucasian
Previous history	Morbid obesity	Morbid obesity	Morbid obesity, anemia and pancytopenia	Morbid obesity	Morbid obesity	Previous history of fatty liver and irregular menses
Clinical setting	Weight loss due to long-limb Roux-en-Y bypass	Weight loss due to Roux-en-Y bypass	Weight loss due to low carbohydrate and high protein diet	Weight loss due to Roux-en-Y bypass	Weight loss due to Roux-en-Y bypass	1-week history of abdominal pain and distention
Alcohol	None	None	Occasionally	Occasionally	None	None
Medications	None	Omeprazole	Omeprazole, vitamin K	Benadryl	None	None
Weight loss/time	18 kg over 5 months	91 kg over 3 years	23 kg over 1 year	45 kg over 8 months	40 kg over 7 years	Not available
BMI changes	45.3 to 38.4	59.4 to 26.5	39.7 to 30.2	55.1 to 36.5	46 to 32	Current BMI: 24.5
Symptoms/signs	Icteric skin, nausea, vomiting, jaundice and encephalopathy	Jaundiced, asterixis, ascites, prominent hepatomegaly	Jaundiced, ascites and encephalopathy	Asterixis, icteric skin, prominent hepatomegaly, lower extremity edema, Right upper quadrant abdominal pain	Hepatomegaly, severe malnutrition with peripheral neuropathy and thiamine deficiency, portal hypertension with varices	Hepatomegaly
Initial symptoms prior to admission	19 days	3 months	2 months	Not available	Not available	1 week
Liver tests	AST = 80 U/l ALT = 143 U/l AP = 189 U/l Total bilirubin = 11.7 mg/dl Albumin = 2.4 g/dl Evidence of coagulopathy	AST = 105 U/l ALT = 39 U/l AP = 124 U/l Total bilirubin = 27.6 mg/dl Albumin = 1.8 g/dl INR = 2.4	AST = 115 U/l ALT = 25 U/l AP = 109 U/l Total bilirubin = 35.2 mg/dl Albumin = 2.1 g/dl INR = 2.1	AST = 110 U/l ALT = 25 U/l AP = 98 U/l Total bilirubin = 11 mg/dl Albumin = 1.3 g/dl INR = 1.4	Transaminases: elevated Bilirubin: within normal limits Evidence of coagulopathy	AST = 106 U/l ALT = 22 U/l AP: within normal limit Total bilirubin = 2.8 mg/dl Albumin = 2.3 g/dl
Viral and autoimmune serologies	HAV IgM (–) HBsAg (–) HBsAb (–) HCV IgM (–) CMV Ab (+) ANA < 40	HAV IgM (–) HBsAg (–) HBsAb (–) HCV IgM (–) HCV IgM (–) HIV Ab (–) CMV Ab (+) EBNA Ab (+) EBV IgM (+) ANA < 40 Anti-smooth muscle (–) LKM Ab (–)	HAV IgG (+) HAV IgM (–) HBsAg (–) HBcAb (–) HBsAb (+) HBc IgM (–) HCV Ab (–) HIV Ab (–) CMV Ab (–) EBNA Ab (+) EBV IgM (–) AMA (–)	HAV IgG (–) HAV IgM (–) HBsAg (–) HBcAb (–) HBc IgM (–) HCV Ab (–) HIV Ab (–) CMV Ab (–) EBNA Ab (–) EBV IgM (–) AMA (–)	Not available	HAV, HBV, HCV (–) IgG = 1700, mildly elevated ANA (–) LKM (–)
Other labs	Ceruloplasmin: within normal limits Ferritin = 305 µg/l α-1-antitrypsin: within normal limits	Ceruloplasmin: within normal limits α-1-antitrypsin: within normal limits	Ceruloplasmin: within normal limits α-1-antitrypsin: within normal limits	Ceruloplasmin: within normal limits Ferritin = 735 µg/l		Ceruloplasmin: within normal limits α-1-antitrypsin: within normal limits
Outcome	Death	Liver transplantation	Liver transplantation	Death	Improved after medical treatment	Not available

Abbreviations: Albumin (normal adult: 3.5–4.8 g/dl); ALT, alanine aminotransferase (normal adult female: 11–50 U/l); AMA, anti-mitochondrial antibody; ceruloplasmin (normal: 19–68 mg/dl); ANA, antinuclear antibody (normal: negative titer < 40); α-1-antitrypsin (normal: 79–207 mg/dl); AP, alkaline phosphatase (normal: adult female: 31–95 U/l); AST, aspartate aminotransferase (normal adult: 17–42 U/l); total bilirubin (normal: 0.2–1.3 mg/dl); BMI, body mass index; CMV, cytomegalovirus; EBNA, Epstein–Barr Virus nuclear antigen; EBV, Epstein–Barr virus; ferritin (normal females age 30–39: 12–160 µg/l); HAV, hepatitis A virus; HBsAg, hepatitis B surface antigen; HBsAb, hepatitis B surface antibody; HBc, hepatitis B core; HCV, hepatitis C; HIV, human immunodeficiency virus; INR, international normalized ratio (normal: 0.9–1.3); LKM Ab, liver kidney microsomal antibody.

Table 2 Aggressive non-alcoholic steatohepatitis histologic findings

	Patient 1	Patient 2	Patient 3	Patient 4	Patient 5	Patient 6
Specimen	Autopsy	Explant	Explant	Autopsy	Needle biopsy	Needle biopsy
Gross findings	Massive hepatomegaly with focal infarcts (6680 g)	Micronodular cirrhosis (1825 g)	Early micronodular cirrhosis (1491 g)	Massive hepatomegaly with severe fatty change (5600 g)	Not available	Not available
Fibrosis stage ^a	Stage 3	Stage 4	Stage 4	Stage 3	Stage 3–4	Stage 3
Total macrovesicular steatosis (large and small droplet; %)	80%	10%	40%	40%	40%	80%
Severe centrilobular scarring	Yes	Yes	Yes	Yes	Yes	Yes
Location of steatosis	Panacinar	Azonal	Panacinar	Azonal	Panacinar	Panacinar
Ballooned hepatocytes	Prominent	Prominent	Prominent	Prominent	Prominent	Prominent
Mallory–Denk bodies	Many	Many	Many	Many	Many	Many
Ductular reaction	Yes	Yes	Yes	Yes	No	Yes
Cholestasis	No	Yes	Yes	Yes	No	No
Perivenular sclerosis/occlusion	No/No	Yes/Yes	Yes/Yes	Yes/No	Yes/Yes	Yes/Yes
Centrilobular arteries	No	Yes	Yes	Yes	Yes	Yes
Trichrome staining pattern	Bridging fibrosis	Two-tone staining pattern	Two-tone staining pattern	Bridging fibrosis	Two-tone staining pattern	Two-tone staining pattern
Elastic stain results (Elastic can Gieson and/or orcein stains)	Not available	Lack of elastic fibers in centrilobular scars	Lack of elastic fibers in centrilobular scars	Not available	Lack of elastic fibers in centrilobular scars	Not available

^aAccording to the Non-alcoholic Steatohepatitis Clinical Research Network staging methodology.²²

which suggested subacute injury. Elastic stains were performed on biopsies from three patients and demonstrated a lack of well-defined elastic fibers in pale staining areas, which confirmed our impression of subacute dropout rather than established fibrosis (Figure 2d).

Fibrosis was staged (summarized in Table 2) using Non-alcoholic Steatohepatitis Clinical Research Network methodology.²² In patient 5, there was severe centrilobular scarring (as well as pale staining areas on trichrome stain suggestive of subacute dropout), sufficient for classification as at least bridging fibrosis (Figure 3a and b), which was overall considered stage 3–4 based on degree of pericellular fibrosis (Figure 3c). A follow-up biopsy was available three years after clinical resolution of liver failure, which demonstrated significant reduction in fibrosis with only few thin, bridging, collagenous bands around vaguely nodular hepatic lobules (Figure 3d), to suggest remodeling of prior fibrosis. Notably, no steatosis, ballooned hepatocytes, or Mallory–Denk bodies were seen in this biopsy.

Discussion

In this study, we describe a series of patients with an aggressive form of non-alcoholic steatohepatitis related to rapid weight loss/malnutrition, who had no significant alcohol exposure, and who developed severe liver dysfunction that usually resulted in liver transplantation or death. Liver biopsy demonstrated a characteristic combination of severe centrilobular scarring, abundant hepatocellular ballooning, and numerous Mallory–Denk bodies. Extensive pericellular fibrosis was the most striking finding and was well beyond what is typically seen in non-alcoholic steatohepatitis. Ductular reaction and cholestasis were also particularly prominent in three patients with elevated bilirubin. Aggressive non-alcoholic steatohepatitis shares similar histologic features with alcoholic steatohepatitis, namely abundant Mallory–Denk bodies, prominent neutrophil infiltrate around ballooned hepatocytes, central vein obliteration, pericellular fibrosis, and cholestasis.^{6,7} The finding of subacute injury, which can be best identified on a well-controlled trichrome stain (though occasionally an elastic stain is needed for confirmation of these findings, as elastic fiber deposition supports early fibrosis^{20,21}), further supports consideration of a rapidly progressive or acute/subacute process. Four of six patients had an increased aspartate aminotransferase/alanine aminotransferase ratio (> 2:1), which has been associated with alcoholic steatohepatitis. Although it is not possible to absolutely exclude concealment of significant alcohol consumption, increased aspartate aminotransferase/alanine aminotransferase ratio is not entirely specific for alcoholic steatohepatitis and has been described more generally with advanced fibrosis;²³ also, specificity of this ratio has not been evaluated in rapidly

Figure 1 Pathologic features of aggressive non-alcoholic steatohepatitis in patient 2. (a) and (b) Examples of severe centrilobular scarring with pericellular fibrosis. Asterisks indicate portal tracts. (c) Prominent ballooned hepatocytes with Mallory–Denk bodies. (d) Centrizonal arteries (arrowheads) in fibrous central zones. (e) Ductular reaction and cholestasis. (f) Perivenular sclerosis with central vein occlusion. (a–f: H&E stain; original magnification: (a): $\times 40$; (b): $\times 100$; (c–f): $\times 200$). H&E, hematoxylin and eosin.

progressive steatohepatitis. In our opinion, both alcoholic steatohepatitis and aggressive non-alcoholic steatohepatitis should be included in the histologic differential diagnosis when a patient presents with a rapidly progressive form of steatohepatitis.

It is estimated that 16% of bariatric surgery patients have incidental non-alcoholic steatohepatitis with nearly 4% expected to have bridging fibrosis

or cirrhosis on initial evaluation,²⁴ which suggests slow evolution of a clinically silent disease in this population. In contrast, aggressive non-alcoholic steatohepatitis presents with rapid liver failure before there is evidence of cirrhosis. This scenario was confirmed on liver biopsy from one patient in our study, who had only mild centrilobular pericellular fibrosis on initial biopsy, but after rapid weight loss developed aggressive non-alcoholic

Figure 2 Additional examples of fibrosis in aggressive non-alcoholic steatohepatitis. (a) Broad dense centrizonal fibrosis, H&E stain. (b) Perivenular sclerosis and intimal thickening. Centrizonal arteries are also present to the left of the vein, trichrome stain. (c) Subacute changes with two-tone trichrome staining; dense, thick collagen bundles stain dark blue (asterisks), while regions with subacute dropout stain pale blue (arrows), trichrome stain. (d) Elastic fiber deposition in early fibrosis (arrows—lack of elastic fiber deposition in regions of necrosis; asterisks—elastic fiber deposition in early fibrosis), orcein stain. (Original magnification: (a): $\times 100$; B: $\times 200$; (c, d): $\times 40$). H&E, hematoxylin and eosin.

steatohepatitis. These observations suggest an acute/subacute mechanism of centrizonal liver injury related to rapid weight loss and/or malnutrition. Also, one survivor was found to have resolution of hepatocellular ballooning, with fibrosis remodeling, three years after presentation, to suggest that the liver may return to adequate function after weight stabilizes and/or nutrition returns to normal. Utilization of traditional staging and grading schemes for non-alcoholic steatohepatitis may not be appropriate in the setting of aggressive non-alcoholic steatohepatitis, as the original schemes were not developed to assess rapidly progressive liver injury.²²

A mechanism for aggressive non-alcoholic steatohepatitis in the setting of rapid weight loss/malnutrition is not yet apparent. Since most affected patients, in other reports on rapid weight loss, were morbidly obese, attention has been drawn to an increase in lipolysis and enhanced release of endogenous free fatty acid from adipose deposits.^{18,25,26} Fat is rapidly mobilized from visceral adipose tissue and free fatty acids flood the

portal circulation in rapid weight loss and may perhaps overwhelm hepatic parenchyma.²⁵ Oxidation of free fatty acid by hepatocytes, which first occurs in the central zone, creates reactive oxygen species that may trigger mitochondrial injury and cytokine release (eg, tumor necrosis factor- α and interleukin-6),²⁷ which may correspond to damage in Zone 3 (as with other agents known to increase liver free radical formation (eg, carbon tetrachloride, acetaminophen, ethanol) or in ischemia-reperfusion injury). Also, increased intestinal permeability in malnutrition leads to an influx of endotoxin into the portal circulation, which increases tumor necrosis factor- α production.^{28,29} Tumor necrosis factor- α has been specifically implicated as a biomarker in non-alcoholic steatohepatitis.³⁰ Both fatty acids and endotoxin are hypothesized to aggravate the inflammatory activity of steatohepatitis.²⁷ Furthermore, a fatty liver is less tolerant to oxidative stress than normal liver.^{18,31,32} Malnutrition also depresses hepatic antioxidant reserves and impairs overall metabolic function of mitochondria.^{33,34} Similarly,

Figure 3 Histologic resolution/regression in patient 5. (a) Initial presenting biopsy at the time of liver failure with severe centrizonal scarring with central vein occlusion (asterisk), H&E stain. (b) Initial biopsy with many ballooned hepatocytes and Mallory–Denk bodies, H&E stain. (c) Initial biopsy with two-tone trichrome staining, highlighting centrizonal scar and obliterated vein (asterisk), trichrome stain. (d) Follow-up biopsy, three years after clinical resolution, without steatosis or ballooned hepatocytes. There are thin fibrous septa stretching from the central zones with mild/moderate portal and lobular inflammation, H&E stain. (Original magnification: (a): $\times 100$; (b): $\times 200$; (c, d): $\times 100$). H&E, hematoxylin and eosin.

primary graft dysfunction, which can occur in association with markedly steatotic liver allografts, corresponds to an impaired ability to recover from ischemia reperfusion.³⁵ It seems plausible that sustained delivery of enough free fatty acid (and perhaps associated endotoxin/cytokines), to an already compromised fatty liver, may underlie development of aggressive non-alcoholic steatohepatitis,^{18,31} but mechanistic studies are needed to further investigate this hypothesis and to consider a potential role for antioxidants to protect the liver during rapid weight loss/malnutrition. A genetic predisposition to developing rapidly progressive steatohepatitis is also a consideration and molecular testing could allow for additional insight into the mechanism of disease. For example, genome wide association analyses have identified specific gene polymorphisms (ie, patatin-like phospholipase domain containing 3 (*PNPLA3*, rs738409, encoding I148M) and transmembrane 6 superfamily member 2 (*TM6SF2*, rs58542926, encoding Glu167Lys)) that

may confer increased susceptibility to development of steatohepatitis.^{3,36}

Jejuno–ileal bypass is contraindicated for weight loss due to a high incidence of progressive hepatic disease leading to liver failure and cirrhosis.¹⁴ Such cases are reported to show histologic features similar to alcoholic steatohepatitis.³⁷ Liver complications in jejuno–ileal bypass are complex, with only a subset of patients developing liver failure, and the etiology is considered multifactorial, with ongoing chronic liver injury noted in some patients.^{14,37,38} In addition to marked weight loss and severe malnutrition (related to poor absorption and severe diarrhea), the excluded intestine in this procedure is predisposed to bacterial overgrowth and synthesis of harmful endotoxin/lipopolysaccharide and/or bile acid metabolites.^{14,18,25,26,38} In contrast, Roux-en-Y gastric bypass has a shorter bypassed segment, with exposure to bile and pancreatic juice, which reduces bacterial overgrowth.³⁹ In addition, Roux-en-Y gastric bypass usually achieves more controlled weight loss than jejuno–ileal bypass.^{17,26,40} Liver failure

associated with Roux-en-Y gastric bypass is considered a rare event⁴¹ and cases of aggressive non-alcoholic steatohepatitis seen in our series were specifically associated with rapid weight loss and/or malnutrition rather than a surgical complication. Roux-en-Y gastric bypass is considered a safe procedure if drastic weight loss is avoided in the postoperative course.^{17,25}

In conclusion, we have encountered several cases of aggressive non-alcoholic steatohepatitis in association with rapid weight loss/malnutrition, which can result in over-emphasis of a potential alcohol etiology in pathology reports. While it may not be possible to entirely exclude an alcohol etiology, acute liver failure with extensive pericellular fibrosis, steatosis, hepatocellular ballooning, Mallory–Denk bodies, cholestasis, and central vein obliteration should not be specifically ascribed to alcohol and may represent an acute to subacute response to central zone injury during rapid weight loss/malnutrition. The histologic findings in aggressive non-alcoholic steatohepatitis may point to a similar mechanism of disease between alcoholic steatohepatitis and non-alcoholic steatohepatitis.

Disclosure/conflict of interest

The authors declare no conflict of interest.

References

- Ludwig J, Viggiano TR, McGill DB, *et al*. Nonalcoholic steatohepatitis: Mayo Clinic experiences with a hitherto unnamed disease. *Mayo Clin Proc* 1980;55:434–438.
- Farrell GC, Larter CZ. Nonalcoholic fatty liver disease: from steatosis to cirrhosis. *Hepatology* 2006;43:S99–S112.
- Yu J, Marsh S, Hu J, *et al*. The Pathogenesis of Nonalcoholic Fatty Liver Disease: Interplay between Diet, Gut Microbiota, and Genetic Background. *Gastroenterol Res Pract* 2016;2016:2862173.
- Younossi ZM, Koenig AB, Abdelatif D, *et al*. Global epidemiology of nonalcoholic fatty liver disease-Metanalytic assessment of prevalence, incidence, and outcomes. *Hepatology* 2016;64:73–84.
- Charlton MR, Burns JM, Pedersen RA, *et al*. Frequency and outcomes of liver transplantation for nonalcoholic steatohepatitis in the United States. *Gastroenterology* 2011;141:1249–1253.
- Yeh MM, Brunt EM. Pathological features of fatty liver disease. *Gastroenterology* 2014;147:754–764.
- Yeh MM, Brunt EM. Pathology of nonalcoholic fatty liver disease. *Am J Clin Pathol* 2007;128:837–847.
- Angulo P. Long-term mortality in nonalcoholic fatty liver disease: is liver histology of any prognostic significance? *Hepatology* 2010;51:373–375.
- Geerts A, Darius T, Chapelle T, *et al*. The multicenter Belgian survey on liver transplantation for hepatocellular failure after bariatric surgery. *Transplant Proc* 2010;42:4395–4398.
- Nanki T, Koike R, Miyasaka N. Subacute severe steatohepatitis during prednisolone therapy for systemic lupus erythematosus. *Am J Gastroenterol* 1999;94:3379.
- Capron JP, Delamarre J, Dupas JL, *et al*. Fasting in obesity: another cause of liver injury with alcoholic hyaline? *Dig Dis Sci* 1982;27:265–268.
- Peura DA, Stromeyer FW, Johnson LF. Liver injury with alcoholic hyaline after intestinal resection. *Gastroenterology* 1980;79:128–130.
- Miura H, Ijichi M, Ando Y, *et al*. A rapidly progressive and fatal case of nonalcoholic steatohepatitis following pancreaticoduodenectomy. *Clin J Gastroenterol* 2013;6:470–475.
- Hocking MP, Duerson MC, O'Leary JP, *et al*. Jejunoileal bypass for morbid obesity. Late follow-up in 100 cases. *N Engl J Med* 1983;308:995–999.
- Cotler SJ, Vitello JM, Guzman G, *et al*. Hepatic decompensation after gastric bypass surgery for severe obesity. *Dig Dis Sci* 2004;49:1563–1568.
- Caldwell SH, Hespenheide EE. Subacute liver failure in obese women. *Am J Gastroenterol* 2002;97:2058–2062.
- Shaffer EA. Bariatric surgery: a promising solution for nonalcoholic steatohepatitis in the very obese. *J Clin Gastroenterol* 2006;40:S44–S50.
- Mattar SG, Velcu LM, Rabinovitz M, *et al*. Surgically-induced weight loss significantly improves nonalcoholic fatty liver disease and the metabolic syndrome. *Ann Surg* 2005;242:610–617.
- Gill RM, Belt P, Wilson L, *et al*. Centrizonal arteries and microvessels in nonalcoholic steatohepatitis. *Am J Surg Pathol* 2011;35:1400–1404.
- Ferrell LD, Greenberg MS. Special stains can distinguish hepatic necrosis with regenerative nodules from cirrhosis. *Liver Int* 2007;27:681–686.
- Nakayama H, Itoh H, Kunita S, *et al*. Presence of perivenular elastic fibers in nonalcoholic steatohepatitis Fibrosis Stage III. *Histol Histopathol* 2008;23:407–409.
- Kleiner DE, Brunt EM, Van Natta M, *et al*. Design and validation of a histological scoring system for nonalcoholic fatty liver disease. *Hepatology* 2005;41:1313–1321.
- Williams AL, Hoofnagle JH. Ratio of serum aspartate to alanine aminotransferase in chronic hepatitis. Relationship to cirrhosis. *Gastroenterology* 1988;95:734–739.
- Kleiner DE, Berk PD, Hsu JY, *et al*. Hepatic pathology among patients without known liver disease undergoing bariatric surgery: observations and a perspective from the longitudinal assessment of bariatric surgery (LABS) study. *Semin Liver Dis* 2014;34:98–107.
- Verna EC, Berk PD. Role of fatty acids in the pathogenesis of obesity and fatty liver: impact of bariatric surgery. *Semin Liver Dis* 2008;28:407–426.
- de Freitas AC, Campos AC, Coelho JC. The impact of bariatric surgery on nonalcoholic fatty liver disease. *Curr Opin Clin Nutr Metab Care* 2008;11:267–274.
- Csak T, Ganz M, Pespisa J, *et al*. Fatty acid and endotoxin activate inflammasomes in mouse hepatocytes that release danger signals to stimulate immune cells. *Hepatology* 2011;54:133–144.
- Welsh FK, Farmery SM, MacLennan K, *et al*. Gut barrier function in malnourished patients. *Gut* 1998;42:396–401.
- Nolan JP. The role of intestinal endotoxin in liver injury: a long and evolving history. *Hepatology* 2010;52:1829–1835.
- Ajmera V, Perito ER, Bass NM, *et al*. Novel plasma biomarkers associated with liver disease severity in adults with nonalcoholic fatty liver disease. *Hepatology* 2017;65:65–77.

- 31 Day CP. NASH-related liver failure: one hit too many? *Am J Gastroenterol* 2002;97:1872–1874.
- 32 Day CP, James OF. Steatohepatitis: a tale of two ‘hits’? *Gastroenterology* 1998;114:842–845.
- 33 Robinson MK, Rustum RR, Chambers EA, *et al*. Starvation enhances hepatic free radical release following endotoxemia. *J Surg Res* 1997;69:325–330.
- 34 van Zutphen T, Ciapaite J, Bloks VW, *et al*. Malnutrition-associated liver steatosis and ATP depletion is caused by peroxisomal and mitochondrial dysfunction. *J hepatol* 2016;65:1198–1208.
- 35 Marsman WA, Wiesner RH, Rodriguez L, *et al*. Use of fatty donor liver is associated with diminished early patient and graft survival. *Transplantation* 1996;62:1246–1251.
- 36 Anstee QM, Seth D, Day CP. Genetic factors that affect risk of alcoholic and nonalcoholic fatty liver disease. *Gastroenterology* 2016;150:1728–1744.
- 37 Peters RL. Patterns of hepatic morphology in jejunoileal bypass patients. *Am J Clin Nutr* 1977;30:53–57.
- 38 O’Leary JP. Hepatic complications of jejunoileal bypass. *Semin Liver Dis* 1983;3:203–215.
- 39 Sugerman HJ, Kellum JM, DeMaria EJ. Conversion of proximal to distal gastric bypass for failed gastric bypass for superobesity. *J Gastrointest Surg* 1997;1:517–524.
- 40 Griffen WO Jr, Young VL, Stevenson CC. A prospective comparison of gastric and jejunoileal bypass procedures for morbid obesity. *Surg Obes Relat Dis* 2005;1:163–172.
- 41 Mahawar KK, Parmar C, Graham Y, *et al*. Monitoring of Liver Function Tests after Roux-en-Y Gastric Bypass: An Examination of Evidence Base. *Obes Surg* 2016;26:2516–2522.