

24. Pan, T., Loria, A. & Zhong, K. Probing of tertiary interactions in RNA: 2'-hydroxyl-base contacts between the RNase P RNA and pre-tRNA. *Proc. Natl Acad. Sci. USA* **92**, 12510–12514 (1995).
25. Loria, A. & Pan, T. Recognition of the T stem-loop of a pre-tRNA substrate by the ribozyme from *Bacillus subtilis* ribonuclease P. *Biochemistry* **36**, 6317–6325 (1997).
26. Nolan, J. M., Burke, D. H. & Pace, N. R. Circularly permuted tRNAs as specific photoaffinity probes of ribonuclease P RNA structure. *Science* **261**, 762–765 (1993).
27. Hendrickson, W. A. Determination of macromolecular structures from anomalous diffraction of synchrotron radiation. *Science* **254**, 51–58 (1991).
28. Murshudov, G. N., Vagin, A. A. & Dodson, E. J. Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr. D* **53**, 240–255 (1997).
29. Brunger, A. T. *et al.* Crystallography & NMR system: a new software suite for macromolecular structure determination. *Acta Crystallogr. D* **54**, 905–921 (1998).
30. Haas, E. S., Banta, A. B., Harris, J. K., Pace, N. R. & Brown, J. W. Structure and evolution of ribonuclease P RNA in Gram-positive bacteria. *Nucleic Acids Res.* **24**, 4775–4782 (1996).

Supplementary Information accompanies the paper on *Nature's* website (<http://www.nature.com/nature>).

Acknowledgements We thank X. Liu and Y. Xiao for technical assistance, A. Changela, H. Feinberg, V. Grum and members of DND–CAT for help with data collection, and A. Changela, C. Correll, V. Grum, E. Sontheimer, B. Taneja and J. Wedekind for comments and suggestions. Research was supported by the NIH (to A.M.) and an NIH NRSA Fellowship to A.K. Support from the R.H. Lurie Cancer Center of Northwestern University to the Structural Biology Center is acknowledged. Portions of this work were performed at the DuPont–Northwestern–Dow Collaborative Access Team (DND–CAT) Synchrotron Research Center at the Advanced Photon Source (APS) and at the Stanford Synchrotron Radiation Laboratory (SSRL). DND–CAT is supported by DuPont, Dow and the NSF, and use of the APS is supported by the DOE. SSRL is operated by the DOE, Office of Basic Energy Sciences. The SSRL Biotechnology Program is supported by the NIH and the DOE.

Competing interests statement The authors declare that they have no competing financial interests.

Correspondence and requests for materials should be addressed to A.M. (e-mail: a-mondragon@northwestern.edu). Coordinates have been deposited in the Protein Data Bank under the accession code 1NBS.

retractions

A cytosolic catalase is needed to extend adult lifespan in *C. elegans* *daf-1* and *clk-1* mutants

J. Taub, J. F. Lau, C. Ma, J. H. Hahn, R. Hoque, J. Rothblatt & M. Chalfie

Nature **399**, 162–166 (1999).

We no longer have confidence in our observations associating a reduction in adult lifespan with a putative mutation in the *Caenorhabditis elegans* catalase gene *ctl-1* and therefore retract this paper. With the assistance of J. Liang and C. Keller, we have confirmed that *C. elegans* has multiple catalase genes (actually three in tandem) and that the original strain, TU1061, has decreased transcription of *ctl-1* messenger RNA. However, we have also found several errors, one identifying a single nucleotide deletion as the defect in the putative *ctl-1* mutation and others in the identification of strains carrying mutations in multiple genes. In particular, we have not seen the expected reduction in *ctl-1* mRNA in other

strains tested. The longevity results obtained with these strains are therefore meaningless. We are grateful to our colleagues, particularly C. Kenyon and M. Crowder, for conveying to us their concerns about our results. □

Metal–insulator transition in chains with correlated disorder

Pedro Carpena, Pedro Bernaola-Galván, Plamen Ch. Ivanov & H. Eugene Stanley

Nature **418**, 955–959 (2002).

This Letter reported numerical simulations of one-dimensional disordered binary systems, and found a threshold value for the exponent characterizing the long-range power-law correlations of the system. Below this threshold, the system behaves as an insulator and above it, in the thermodynamic limit, the system behaves as a conductor. Unfortunately, we have now found that this observation was a consequence of the algorithm used to generate long-range correlations in binary chains, because above the threshold value of the exponent only a finite number of segments of atoms of the same type (A or B) exists, even in the thermodynamic limit of an infinitely large system. Thus, the system studied was not truly disordered. As a result, what we observed at the critical threshold value for the correlation exponent was not a transition from insulator to metal behaviour in a disordered system (as reported), but a transition from a disordered to an ordered system. For this reason, the authors retract the claim of a metal–insulator transition in the infinite binary chain with correlated disorder. The results are still valid that relate to the behaviour of a binary chain below the critical threshold value of the correlation exponent, and to large but finite system sizes (as found in the DNA example discussed in the Letter).

We thank L. Hufnagel and T. Geisel for drawing this to our attention. □

erratum

Wave-like properties of solar supergranulation

L. Gizon, T. L. Duvall Jr & J. Schou

Nature **421**, 43–44 (2003).

In Fig. 1, the units of frequency should be microhertz (μHz), not millihertz (mHz). In the US-printed issues, Fig. 3b appeared blurred. □